

THE LOCAL MOTIVIC MONODROMY CONJECTURE FOR SIMPLICIAL NONDEGENERATE SINGULARITIES

MATT LARSON, SAM PAYNE, AND ALAN STAPLEDON

ABSTRACT. We prove the local motivic monodromy conjecture for singularities that are nondegenerate with respect to a simplicial Newton polyhedron. It follows that all poles of the local topological zeta functions of such singularities correspond to eigenvalues of monodromy acting on the cohomology of the Milnor fiber of some nearby point, as do the poles of Igusa's local p -adic zeta functions for large primes p .

CONTENTS

1.	Introduction	1
2.	Examples	13
3.	A nonnegative formula for nearby eigenvalues	16
4.	A necessary condition for the vanishing of the local h -polynomial	26
5.	The local formal zeta function and its candidate poles	41
6.	Fake poles for the local formal zeta function	46
7.	Beyond the simplicial case	65
	References	67

1. INTRODUCTION

Throughout, let \mathbb{k} be a field of characteristic 0, and let $f \in \mathbb{k}[x_1, \dots, x_n]$ be a regular function whose vanishing locus X_f contains $0 \in \mathbb{A}^n$. The coefficients of f are contained in a finitely generated subfield $\mathbb{k}' \subset \mathbb{k}$, so we may choose an embedding $\mathbb{k}' \subset \mathbb{C}$, view f as a holomorphic function on \mathbb{C}^n , and consider the Milnor fiber \mathcal{F}_x , with its monodromy action, for any geometric point $x \in X_f$. The characteristic polynomial of the induced action on $H^*(\mathcal{F}_x, \mathbb{C})$ is independent of all choices and its zeros are the *eigenvalues of monodromy* of f at x . The monodromy is quasi-unipotent, so all such eigenvalues of monodromy are roots of unity. We say that $\exp(2\pi i\alpha)$ is a *nearby eigenvalue of monodromy* of f if 0 lies in the Zariski closure of the locus of points $x \in X_f$ such that $\exp(2\pi i\alpha)$ is an eigenvalue of monodromy of f at x .

The local motivic zeta function is a subtle invariant of the singularity of f at 0, introduced by Denef and Loeser [DL98]. Let $K^{\hat{\mu}}$ be the Grothendieck ring of \mathbb{k} -varieties with good $\hat{\mu}$ -action, where $\hat{\mu} = \varprojlim \mu_m$ is the inverse limit of the groups of m th roots of unity, and let $\mathcal{M}^{\hat{\mu}} := K^{\hat{\mu}}[\mathbb{L}^{-1}]$ be the associated motivic ring obtained by inverting $\mathbb{L} := [\mathbb{A}^1]$. Then the local motivic zeta function $Z_{\text{mot}}(T) \in \mathcal{M}^{\hat{\mu}}[[T]]$ is expressible non-uniquely as the formal power series expansion of a rational function in $\mathcal{M}^{\hat{\mu}}\left[T, \frac{1}{1-\mathbb{L}^a T^b}\right]_{(a,b) \in \mathbb{Z} \times \mathbb{Z}_{>0}, a/b \in \mathcal{P}}$ for some finite $\mathcal{P} \subset \mathbb{Q}$. Any such \mathcal{P} is a *set of candidate poles* for $Z_{\text{mot}}(T)$, as defined in [BV16, BN20].

Local Motivic Monodromy Conjecture. *There is a set of candidate poles $\mathcal{P} \subset \mathbb{Q}$ for $Z_{\text{mot}}(T)$ such that, for every $\alpha \in \mathcal{P}$, $\exp(2\pi i\alpha)$ is a nearby eigenvalue of monodromy.*

Date: May 23, 2024.

Note that the notion of poles is subtle in this context because $K^{\hat{\mu}}$ is not known to be an integral domain; in particular, it is unclear whether the intersection of two sets of candidate poles for $Z_{\text{mot}}(T)$ is necessarily a set of candidate poles. Our first main result (Theorem 1.1.1) confirms the local motivic monodromy conjecture for singularities that are nondegenerate with respect to a simplicial Newton polyhedron.

1.1. Main results. For $u = (u_1, \dots, u_n)$ in $\mathbb{Z}_{\geq 0}^n$, let $x^u := x_1^{u_1} \cdots x_n^{u_n}$, and write $f = \sum_u a_u x^u$. The Newton polyhedron of f , denoted $\text{Newt}(f)$, is the Minkowski sum $\text{conv}\{u : a_u \neq 0\} + \mathbb{R}_{\geq 0}^n$. For each face F of $\text{Newt}(f)$, we consider $f|_F := \sum_{u \in F} a_u x^u$. Then f is *nondegenerate* if, for all compact faces F , the vanishing locus of $f|_F$ has no singularities in the complement of the coordinate hyperplanes in \mathbb{A}^n .

For any face F of $\partial \text{Newt}(f)$ that meets the interior of the orthant $\mathbb{R}_{> 0}^n$, let $C_F := \overline{\mathbb{R}_{\geq 0} F}$ be the closure of the cone spanned by F . The set of all faces of such cones forms a fan Δ whose support is the positive orthant $\mathbb{R}_{\geq 0}^n$. We say that $\text{Newt}(f)$ is *simplicial* if Δ is a simplicial fan.

Theorem 1.1.1. *Suppose that $\text{Newt}(f)$ is simplicial and f is nondegenerate. Then there is a set of candidate poles $\mathcal{P} \subset \mathbb{Q}$ for $Z_{\text{mot}}(T)$ such that, for every $\alpha \in \mathcal{P}$, $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy.*

In other words, the local motivic monodromy conjecture is true for any nondegenerate singularity with a simplicial Newton polyhedron. This was known previously for $n = 2$ [BN20]. Our definition of simplicial Newton polyhedron agrees with that in [JKYS19]. A *convenient* Newton polyhedron, i.e., one that intersects each of the coordinate axes [Kou76], is simplicial if and only if each of its compact faces is a simplex.

Remark 1.1.2. The methods used in the proof of Theorem 1.1.1 are discussed in Section 1.4. Roughly speaking, we have one collection of arguments, presented in Sections 3-4 that prove existence of eigenvalues corresponding to candidate poles associated to many facets of $\text{Newt}(f)$. Another collection of arguments, presented in Section 5, show that certain such candidate poles are *fake* and can be removed to give a smaller set of candidate poles. Each of these arguments is carried out not only for simplicial Newton polyhedra, but in somewhat greater generality. As a result, we are able to prove a range of cases of the local motivic monodromy conjecture where f is nondegenerate and $\text{Newt}(f)$ is not necessarily simplicial, including all such cases for $n = 3$. See Section 7 for details.

The local motivic monodromy conjecture is a motivic analogue of the local p -adic and topological monodromy conjectures, and the following cases of the latter conjectures are consequences of Theorem 1.1.1.

The local motivic zeta function specializes to the local topological zeta function $Z_{\text{top}}(s) \in \mathbb{Q}(s)$ by expanding $Z_{\text{mot}}(T)$ as a power series in $\mathbb{L} - 1$ and setting $T \mapsto \mathbb{L}^{-s}$ and $[Y] \mapsto \chi(Y/\hat{\mu})$ [DL98, Section 2.3]. It follows that the poles of $Z_{\text{top}}(s)$ are contained in every set of candidate poles for $Z_{\text{mot}}(T)$.

Theorem 1.1.3. *Suppose $\text{Newt}(f)$ is simplicial and f is nondegenerate. If α is a pole of $Z_{\text{top}}(s)$, then $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy.*

This confirms the local topological monodromy conjecture [DL92, Conjecture 3.3.2] for singularities that are nondegenerate with respect to a simplicial Newton polyhedron.

If $f \in \mathbb{Z}_p[x_1, \dots, x_n]$ has good reduction mod p , i.e., if $\bar{f} \in \mathbb{F}_p[x_1, \dots, x_n]$ is nondegenerate with $\text{Newt}(\bar{f}) = \text{Newt}(f)$, then $Z_{\text{mot}}(T)$ also specializes to the Igusa local p -adic zeta function $Z_{(p)}(s) \in \mathbb{Q}(p^s)$, which is viewed as a global meromorphic function in the complex variable s . In this case, the real part of any pole of $Z_{(p)}(s)$ is contained in every set of candidate poles for $Z_{\text{mot}}(T)$.

Theorem 1.1.4. *Suppose $f \in \mathbb{Z}_p[x_1, \dots, x_n]$, $\text{Newt}(f)$ is simplicial, and f is nondegenerate with good reduction mod p . If α is a pole of $Z_{(p)}(s)$, then $\exp(2\pi i \Re(\alpha))$ is a nearby eigenvalue of monodromy.*

If $f \in \mathbb{Z}[x_1, \dots, x_n]$ is nondegenerate, then f has good reduction mod p for all but finitely many primes p . In this sense, Theorem 1.1.4 implies that the local p -adic monodromy conjecture holds for nondegenerate singularities with simplicial Newton polyhedra.

1.2. Background and motivation. We now discuss the background and motivation for the local monodromy conjectures. In particular, we recall the definitions of the local motivic, p -adic, and topological zeta functions and how they relate to the geometry of embedded log resolutions. We also recall A'Campo's formula for the zeta function of monodromy at the origin.

1.2.1. Archimedean zeta functions. The motivation for the local monodromy conjectures comes from a theorem of Malgrange concerning the following archimedean analogues of local zeta functions. Suppose $\mathbb{k} = \mathbb{R}$ or \mathbb{C} , and let Φ be a smooth function supported on a compact set that does not contain any critical points of f other than 0. Consider the function

$$Z_{\Phi}(s) := \int_{\mathbb{k}^n} \Phi(x) |f(x)|^{\delta s} dx,$$

where $\delta = 1$ if $\mathbb{k} = \mathbb{R}$ and $\delta = 2$ if $\mathbb{k} = \mathbb{C}$. This integral converges for $s \in \mathbb{C}$ with $\Re(s) > 0$.

Theorem 1.2.1 ([Mal74]). *The function $Z_{\Phi}(s)$ extends to a meromorphic function on \mathbb{C} whose poles are rational numbers. Moreover, if α is a pole of $Z_{\Phi}(s)$, then $\exp(2\pi i\alpha)$ is an eigenvalue of monodromy.*

Furthermore, every eigenvalue of monodromy of f at the origin appears as a pole of $Z_{\Phi}(s)$ for some Φ .

1.2.2. Log resolutions and zeta functions of monodromy. Let $h: \mathcal{F}_x \rightarrow \mathcal{F}_x$ denote the monodromy action on the Milnor fiber of f at $x \in X_f$. The zeta function of monodromy of f at x is then

$$(1) \quad \zeta_x(t) := \frac{\det(1 - th^* | H^{\text{even}}(\mathcal{F}_x, \mathbb{C}))}{\det(1 - th^* | H^{\text{odd}}(\mathcal{F}_x, \mathbb{C}))}.$$

When f is nondegenerate, the zeta function of monodromy at 0 may be expressed in terms of the numerical data of a log resolution and the topological Euler characteristics of the strata in the fiber, as follows.

Let $\pi: Y \rightarrow \mathbb{A}^n$ be a proper morphism that is an isomorphism away from X_f , such that the support of $D := \pi^{-1}(X_f)$ is a divisor with simple normal crossings. Let D_1, \dots, D_r be the irreducible components of D . The associated *numerical data* of this log resolution are the pairs of integers (N_i, ν_i) , where N_i and $\nu_i - 1$ are the orders of vanishing of $\pi^*(f)$ and $\pi^*(dx_1 \wedge \dots \wedge dx_n)$, respectively, along D_i .

For $I \subset \{1, \dots, r\}$, let $D_I := \bigcap_{i \in I} D_i$ and $D_I^{\circ} := D_I \setminus \bigcup_{j \notin I} D_{I \cup \{j\}}$. We then define

$$E_I := D_I \cap \pi^{-1}(0) \text{ and } E_I^{\circ} = D_I^{\circ} \cap \pi^{-1}(0)$$

for the corresponding closed and locally closed strata in the fiber over 0.

Theorem 1.2.2 ([A'C75]). *The zeta function of monodromy acting on the cohomology of \mathcal{F}_0 is*

$$(2) \quad \zeta_0(t) = \prod_{i=1}^r (1 - t^{N_i})^{\chi(E_i^{\circ})},$$

where we omit the corresponding term if E_i° is empty.

Note that the exponents $\chi(E_i^{\circ})$ can be positive or negative, and there can be a great deal of cancellation in simplifying this rational function expression for $\zeta_0(t)$ down to a quotient of two relatively prime polynomials. In particular, it is difficult to determine from the numerical data of the log resolution whether any given root of $1 - t^{N_i}$ is an eigenvalue of monodromy.

1.2.3. *Igusa's local \mathfrak{p} -adic zeta functions.* Let \mathbb{k} be a finite extension of \mathbb{Q}_p , equipped with the unique extension of the p -adic valuation and its associated norm. Let $R \subset \mathbb{k}$ be the valuation ring, with $\mathfrak{p} \subset R$ the maximal ideal. For instance, if $f \in \mathbb{Q}_p[x_1, \dots, x_n]$, then $R = \mathbb{Z}_p$ and $\mathfrak{p} = p\mathbb{Z}_p$. Igusa introduced and studied the local zeta function

$$Z_{\mathfrak{p}}(s) := \int_{\mathfrak{p}^n} |f(x)|^s dx,$$

in his proof of a conjecture of Borewicz and Shafarevich [BS66, p. 63] on rationality of generating functions for the number of solutions mod p^m to a polynomial equation with integer coefficients. Here, dx denotes the normalized Haar measure on the compact additive group \mathfrak{p}^n . Note that $Z_{\mathfrak{p}}(s)$ is a nonarchimedean analogue of the asymptotic integrals $Z_{\Phi}(s)$; the role of Φ is played by the indicator function of the compact subset \mathfrak{p}^n . Igusa proved that $Z_{\mathfrak{p}}(s)$ is a rational function in q^{-s} [Igu75], where $q = |R/\mathfrak{p}|$, and that its real poles other than -1 are all of the form $\alpha_i := -\nu_i/N_i$, where (N_i, ν_i) is the numerical data associated to some exceptional divisor in a given log resolution [Igu78]. Denef gave a second proof of the rationality of $Z_{\mathfrak{p}}(s)$, using p -adic cell decompositions [Den84].

Remark 1.2.3. Note that some sources in the literature define a local \mathfrak{p} -adic zeta function by integrating with respect to the restriction of the normalized Haar measure on R^n ; the result differs from our $Z_{\mathfrak{p}}(s)$ by a factor of q^{-n} . Such renormalizations do not affect the poles of the local zeta functions.

Typically, very few of the rational numbers α_i associated to the numerical data in a log resolution are actually poles of $Z_{\mathfrak{p}}(s)$. In the archimedean setting, this is explained by Malgrange's theorem (Theorem 1.2.1), since many rational numbers that appear in this way do not correspond to eigenvalues of monodromy.

Both Denef [Den85] and Igusa [Igu88] observed that the analogue of Malgrange's theorem seems to hold for $Z_{\mathfrak{p}}(s)$; in all examples that had been computed, whenever α_j is a pole of $Z_{\mathfrak{p}}(s)$, the corresponding root of unity $\exp(2\pi i \alpha_j)$ is an eigenvalue of monodromy. Loeser proved that this is true for $n = 2$ [Loe88] and for certain nondegenerate singularities in higher dimensions [Loe90]. By the early 1990s, the expectation that this nonarchimedean analogue of Malgrange's theorem should hold was known as the monodromy conjecture. See, e.g., [Den91a, Conjecture 4.3], [Den91b, Conjecture 2.3.2], and [Vey93, p. 546–547]. We will follow the usual convention and call this the *local \mathfrak{p} -adic monodromy conjecture* to distinguish it from the topological and motivic variants that followed.

Local \mathfrak{p} -adic Monodromy Conjecture. *Suppose \mathbb{k} is a number field. For all but finitely many primes $\mathfrak{p} \subset \mathcal{O}_{\mathbb{k}}$, if α is a pole of $Z_{\mathfrak{p}}(s)$, then $\exp(2\pi i \Re(\alpha))$ is a nearby eigenvalue of monodromy.*

Interest in this conjecture has persisted through the decades [Nic10, VS22]. Bories and Veys proved it for $n = 3$ when f is nondegenerate [BV16, Theorem 0.12]. There has been little progress in higher dimensions.

1.2.4. *Good reduction.* The local \mathfrak{p} -adic zeta function has a particularly simple expression when $X_f \subset \mathbb{A}^n$ has an embedded log resolution with good reduction mod \mathfrak{p} . Most results showing that poles of $Z_{\mathfrak{p}}(s)$ correspond to nearby eigenvalues of monodromy for $n \geq 3$, including those of [BV16] and our Theorem 1.1.4, have a good reduction hypothesis.

Suppose the log resolution $\pi: Y \rightarrow \mathbb{A}^n$ factors through a closed embedding $Y \hookrightarrow \mathbb{P}^m \times \mathbb{A}^n$ over \mathbb{k} . Let \mathbb{P}_R^m and \mathbb{A}_R^n denote the projective and affine spaces of dimension m and n , respectively, over R . Let X_R and Y_R be the closures of X and Y in \mathbb{A}_R^n and $\mathbb{P}_R^m \times \mathbb{A}_R^n$. Then π extends naturally to a projective morphism $\pi_R: Y_R \rightarrow X_R$. Let \overline{X} and \overline{Y} be the respective special fibers of X_R and Y_R . Base change to $\mathbb{F}_q = R/\mathfrak{p}$ gives a projective morphism $\overline{\pi}: \overline{Y} \rightarrow \overline{X}$. Let \overline{D}_i be the special fiber of the closure of D_i in Y_R .

Definition 1.2.4. *The resolution $\pi: Y \rightarrow \mathbb{A}^n$ has good reduction mod \mathfrak{p} if*

- \overline{Y} is smooth in a neighborhood of $\overline{\pi}^{-1}(0)$;
- $\overline{D}_1, \dots, \overline{D}_r$ are smooth and distinct over \mathbb{F}_q , and they meet each other transversely.

Note that, if f and π are defined over a number field K , then π has good reduction mod \mathfrak{p} for all but finitely many primes \mathfrak{p} in the ring of integers \mathcal{O}_K [Den87, Theorem 2.4].

Any resolution with good reduction mod \mathfrak{p} gives rise to a pleasant formula for $Z_{\mathfrak{p}}(s)$ in terms of the numerical data of the resolution and the number of \mathbb{F}_q -points in the strata of the fiber over 0. Let

$$\overline{E}_I^{\circ} = \{x \in \overline{\pi}^{-1}(0) : x \in \overline{D}_i \text{ if and only if } i \in I\}.$$

Theorem 1.2.5 ([Den87, Theorem 3.1]). *Suppose π has good reduction mod \mathfrak{p} . Then*

$$(3) \quad Z_{\mathfrak{p}}(s) = \sum_{I \subset \{1, \dots, r\}} (q-1)^{|I|} |\overline{E}_I^{\circ}(\mathbb{F}_q)| \prod_{i \in I} \frac{q^{-N_i s - \nu_i}}{1 - q^{-N_i s - \nu_i}}.$$

Note that point counts over finite fields are analogous to topological Euler characteristics over \mathbb{C} ; both are additive with respect to disjoint unions and multiplicative with respect to products. The role of $|\overline{E}_i^{\circ}(\mathbb{F}_q)|$ in (3) is analogous to that of $\chi(E_i^{\circ})$ in (2). When $|\overline{E}_i^{\circ}(\mathbb{F}_q)|$ vanishes, then a term in (3) involving a pole at α_i vanishes, and when the Euler characteristic $\chi(E_i^{\circ})$ vanishes, a term in (2) involving the multiplicity of the corresponding eigenvalue of monodromy vanishes. For more explicit connections, see [Den91a].

1.2.5. *Local topological zeta functions.* The analogy between Euler characteristics and the point counts over finite fields that appear in formulas for the local zeta functions in cases of good reduction leads to the topological zeta functions of Denef and Loeser. Heuristically, these are limits of local \mathfrak{p} -adic zeta functions. The local topological zeta function is defined as follows:

$$Z_{\text{top}}(s) := \sum_{I \subset \{1, \dots, r\}} \chi(E_I^{\circ}) \prod_{i \in I} \frac{1}{N_i s + \nu_i}.$$

It is independent of the choice of resolution [DL92, Theorem 2.1.2].

Suppose f has coefficients in a number field K . Then poles of $Z_{\text{top}}(s)$ give rise to poles of most local \mathfrak{p} -adic zeta functions. More precisely, after clearing denominators, we may assume that f has coefficients in the ring of integers. In this case, if α is a pole of $Z_{\text{top}}(s)$ then, for all but finitely many primes \mathfrak{p} in the ring of integers, there are infinitely many unramified extensions $\mathbb{k} | K_{\mathfrak{p}}$ such that α is a pole of the local \mathfrak{p} -adic zeta function of f over \mathbb{k} . See [DL92, Theorem 2.2].

Local Topological Monodromy Conjecture ([DL92, Conjecture 3.3.2]). *If α is a pole of $Z_{\text{top}}(s)$, then $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy.*

We note that local topological zeta functions have a pleasantly simple expression for singularities that are nondegenerate [DL92, Section 5]. For the corresponding formula for the local \mathfrak{p} -adic zeta function of a nondegenerate singularity with good reduction mod \mathfrak{p} , see [DH01, Theorem 4.2].

One naturally expects that the local topological monodromy conjecture should be easier to prove than the local \mathfrak{p} -adic monodromy conjecture, even in the good reduction case, and experience does bear this out. For instance, both conjectures are known in the special case when f is nondegenerate and $n = 3$. However, the proof of the topological case [LVP11] preceded the proof of the \mathfrak{p} -adic case [BV16] by a few years and is considerably shorter. See [VS22, Exercise 3.65] for an example of a nondegenerate hypersurface (for $n = 5$) with a real pole of its local \mathfrak{p} -adic zeta functions that is not a pole of its local topological zeta function.

1.2.6. *The local motivic zeta function.* The local motivic zeta function of f at 0 is a formal power series with coefficients in a localization of the $\hat{\mu}$ -equivariant Grothendieck ring of varieties and can be defined in terms of an embedded log resolution, as follows.

Let $\mu_m := \text{Spec } \mathbb{k}[t]/(t^m - 1)$ denote the group of m th roots of unity over \mathbb{k} , and let

$$\hat{\mu} := \varprojlim \mu_m.$$

An action of $\hat{\mu}$ on a \mathbb{k} -variety Y is *good* if the action factors through μ_m for some m , and Y is covered by invariant affine opens. The *Grothendieck ring* $K^{\hat{\mu}}$ is additively generated by classes $[Y]$, where Y is a \mathbb{k} -variety with good $\hat{\mu}$ -action, subject to the relations:

- if Z is a closed $\hat{\mu}$ -invariant subvariety, then $[Y] = [Y \setminus Z] + [Z]$;
- if $W \rightarrow Y$ is a $\hat{\mu}$ -equivariant \mathbb{A}^m -bundle, then $[W] = [\mathbb{A}^m \times Y]$.

In the second relation, $\hat{\mu}$ acts trivially on \mathbb{A}^m . Multiplication in the Grothendieck ring $K^{\hat{\mu}}$ is given by $[Y] \cdot [Z] = [Y \times Z]$, with the diagonal $\hat{\mu}$ -action on $Y \times Z$.

For each nonempty subset $I \subset \{1, \dots, r\}$, let $m_I := \gcd\{N_i : i \in I\}$. Then D_I° is covered by Zariski open subsets $U \subset Y$ on which π^*f is of the form ug^{m_I} , where u is a unit on U , and g is a regular function. Consider the Galois cover $\tilde{D}_I^\circ \rightarrow D_I^\circ$, with Galois group μ_{m_I} whose restriction to such an open set $D_I^\circ \cap U$ is

$$\{(z, y) \in \mathbb{A}^1 \times (D_I^\circ \cap U) : z^{m_I} = u^{-1}\}.$$

Then \tilde{D}_I° comes with the evident good $\hat{\mu}$ -action that factors through μ_{m_I} and commutes with the projection to \mathbb{A}^n . Let

$$\tilde{E}_I^\circ := \tilde{D}_I^\circ \times_{\mathbb{A}^n} \{0\}$$

be the induced Galois cover of the fiber of D_I° over 0, with the good $\hat{\mu}$ -action that it inherits from \tilde{D}_I° .

Let $\mathbb{L} := [\mathbb{A}^1]$, and set $\mathcal{M}^{\hat{\mu}} = K^{\hat{\mu}}[\mathbb{L}^{-1}]$. The local motivic zeta function of f at 0 is the formal power series expansion in $\mathcal{M}^{\hat{\mu}}[[T]]$ of the following rational function in $\mathcal{M}^{\hat{\mu}}(T)$:

$$Z_{\text{mot}}(T) = \sum_{I \subset \{1, \dots, r\}} (\mathbb{L} - 1)^{|I|} [\tilde{E}_I^\circ] \prod_{i \in I} \frac{\mathbb{L}^{-\nu_i} T^{N_i}}{1 - \mathbb{L}^{-\nu_i} T^{N_i}}.$$

Note, in particular, that $Z_{\text{mot}}(T)$ is contained in the subring of $\mathcal{M}^{\hat{\mu}}[[T]]$ generated over $\mathcal{M}^{\hat{\mu}}$ by T and $\{\frac{1}{1 - \mathbb{L}^{-\nu_i} T^{N_i}} : 1 \leq i \leq r\}$. This subring depends on the choice of a log resolution, but the power series $Z_{\text{mot}}(T)$ is independent of all choices.

Remark 1.2.6. In the literature, an additional multiplicative factor of \mathbb{L}^{-n} sometimes appears in the definition of the local motivic zeta function. See, e.g. [RV03, (0.1.2)] and [VS22, Theorem 3.18]. Other versions differ from ours by a factor of $\mathbb{L} - 1$ [BN20, Corollary 5.3.2]. These renormalizations are not relevant to the local motivic monodromy conjecture.

Grothendieck rings of varieties are not integral domains [Poo02], so care is required in defining poles of $Z_{\text{mot}}(T)$. Various notions are possible. See, for instance, [RV03, §4]. We follow the now standard convention and state the local motivic monodromy conjecture in terms of sets of candidate poles, as in [BV16, BN20].

Definition 1.2.7. Let \mathcal{P} be a finite set of rational numbers. Then \mathcal{P} is a set of candidate poles for $Z_{\text{mot}}(T)$ if $Z_{\text{mot}}(T)$ is contained in

$$\mathcal{M}^{\hat{\mu}} \left[T, \frac{1}{1 - \mathbb{L}^a T^b} \right]_{(a,b) \in \mathbb{Z} \times \mathbb{Z}_{>0}, a/b \in \mathcal{P}}.$$

Roughly speaking, if α satisfies any reasonable notion of being a pole of $Z_{\text{mot}}(T)$, then it is contained in every set of candidate poles.

Remark 1.2.8. In practice, passing to an embedded log resolution π is not a useful way of computing local zeta functions; this typically introduces many exceptional divisors whose numerical data correspond neither to poles of the zeta function nor to eigenvalues of monodromy. One obtains more efficient expressions for the local zeta functions of nondegenerate singularities by first proving that they can be computed from a log smooth partial resolution [BN20] or a stacky resolution [Que].

1.3. Prior results. The local monodromy conjectures remain wide open in general, despite the persistent efforts of many mathematicians over a period of decades. Perhaps most surprising is that the local topological monodromy conjecture remains open for isolated nondegenerate singularities, even though there are well-known and relatively simple combinatorial formulas for both the characteristic polynomial of monodromy [Var76, Theorem 4.1] and the local topological zeta function [DL92, Theorem 5.3]. Nevertheless, there is a vast literature on the local monodromy conjectures, far more than can reasonably be reviewed here. We give only a brief and largely ahistorical review of prior work closely related to our main theorems, and recommend the excellent survey articles [Nic10, VS22] for more detailed discussions and further references.

1.3.1. Local monodromy conjectures. For $n = 2$, Bultot and Nicaise proved the local motivic monodromy conjecture in full generality [BN20, Theorem 8.2.1].

For nondegenerate singularities when $n = 3$, Lemahieu and Van Proeyen proved the local topological monodromy conjecture [LVP11]. Bories and Veys used the same arguments for existence of eigenvalues and developed new arguments to reduce the size of sets of candidate poles, proving the local \mathfrak{p} -adic monodromy conjecture [BV16]. They also proved a *naive* variant of the local motivic monodromy conjecture for nondegenerate singularities with $n = 3$. In the naive variant, the ring $K^{\hat{\mu}}$ is replaced with the ordinary Grothendieck ring of varieties (without $\hat{\mu}$ -action); the local motivic zeta function specializes to the local naive motivic zeta function by setting $[Y] \mapsto [Y/\hat{\mu}]$.

Esterov, Lemahieu, and Takeuchi introduced new arguments for both existence of eigenvalues and cancellation of poles for local topological zeta functions of nondegenerate singularities, especially for $n = 4$, and stated a conjecture for how these should generalize to higher dimensions [ELT22, Conjecture 1.3]. Recently, while this paper was in the final stages of preparation, Quek produced a naive motivic upgrade for some of the pole cancellation arguments from [ELT22], giving a new proof of the main result of Bories and Veys for $n = 3$. Quek also suggested a different way in which the pole cancellation statements for small n might generalize to higher dimensions [Que, Question 5.1.8]. Neither of these predictions is correct. See Examples 2.2.1 and 2.2.3.

We also note that one of the claimed results on cancellation of poles in [ELT22], namely their Proposition 3.7, which is a special case of their Theorem 4.3, is incorrect. A single facet that is a B_1 -*pyramid not of compact type* in the sense of their Definition 3.1 can contribute a nontrivial pole to the local topological zeta function, and this happens already for $n = 4$. See Example 2.2.4. We use a different notion of non-compact B_1 -facet that agrees with that of Quek [Que, Definition 1.1.7], specializes to that of Lemahieu and Van Proeyen when $n = 3$ [LVP11, Definition 3], and also has the expected pole cancellation property for candidate poles contributed only by a single B_1 -facet in higher dimensions. See Theorem 6.1.2. For compact faces, [ELT22, Proposition 3.7] is correct. From the authors, we understand that the mistake in the corresponding statement for non-compact facets does not seriously affect the other arguments in their paper.

Finally, we note that Budur and van der Veer recently proved the local monodromy conjectures for nondegenerate singularities whose Newton polyhedron is a large dilate of a convenient Newton polyhedron [BvdV22, Theorem 1.10]. Indeed, they show that when $P = \text{Newt}(f)$ is convenient and k is sufficiently large, every candidate eigenvalue corresponding to a facet of kP is an eigenvalue of monodromy. The proof is an application of Varchenko's formula [Var76, Theorem 4.1], and the bound on k depends on P . Here we show, by different arguments that depend on Ehrhart theory and positivity properties of local h -polynomials, that any $k \geq 2$ is large enough. We also prove a generalization of this result when P is not necessarily convenient. See Theorem 3.4.7 and Proposition 3.4.11.

1.3.2. Global zeta functions and strong monodromy conjectures. There are *global* versions of the local motivic, \mathfrak{p} -adic, and topological zeta functions and their associated monodromy conjectures. See, e.g., [DL92] for a discussion of the local and global topological zeta functions. The difference between the local and global

motivic zeta functions is illustrated by [BN20, Theorems 8.3.2 and 8.3.5]. The global zeta functions are invariants of $X_f \subset \mathbb{A}^n$, while the local zeta functions are invariants of its germ at 0.

Replacing “local” by “global” in each of the local monodromy conjectures gives rise to its global counterpart. There are also *strong* versions of the local and global monodromy conjectures proposing that the real parts of the poles of the corresponding zeta functions are zeros of the Bernstein-Sato polynomial b_f . If α is a zero of b_f then $\exp(2\pi i\alpha)$ is an eigenvalue of monodromy, and all eigenvalues of monodromy occur in this way [Mal74]. It is also conjectured that the orders of poles of local zeta functions are bounded by the multiplicities of zeros of b_f [DL92, Conjecture 3.3.1’].

The strong local and global motivic monodromy conjectures are known for $n = 2$ [BN20]. Loeser has given a combinatorial condition on Newton polyhedra that guarantees that each candidate pole associated to a facet is a zero of the Bernstein-Sato polynomial [Loe90]. Nondegenerate polynomials with such Newton polyhedra therefore satisfy the strong local motivic monodromy conjecture.

Aside from this, we note that if X_f is smooth aside from an isolated singularity at 0, then each local monodromy conjecture at 0 implies the corresponding global monodromy conjecture. The Newton polyhedra whose nondegenerate singularities are isolated were classified by Kouchnirenko [Kou76]. Furthermore, if f has such a Newton polyhedron and $f|_F$ has no singularities outside the coordinate hyperplanes for *all* faces F of $\text{Newt}(f)$, not just the compact faces, then X_f is smooth away from 0. Thus the global motivic monodromy conjecture for isolated singularities with simplicial Newton polyhedra that satisfy this stronger nondegeneracy condition follows from Theorem 1.1.1.

1.3.3. Further variants of the local zeta functions and monodromy conjectures. There are also monodromy and holomorphy conjectures for \mathfrak{p} -adic zeta functions *twisted* by a character, and topological analogues of twisted \mathfrak{p} -adic zeta functions. For discussions of these variants, see, e.g., [Den91b]. Another variant is the topological zeta function for a variety equipped with a holomorphic form that plays the role of $\Phi(x)dx$ in Malgrange’s archimedean zeta functions [Vey07]. Our results on eigenvalues of monodromy in Sections 3-4 are applicable to all such variants.

1.4. Methods and structure of the paper. We conclude the introduction with a brief overview of our approach to the local motivic monodromy conjecture and outline the content of each section of the paper.

1.4.1. Key definitions. We first recall the notion of candidate poles and candidate eigenvalues. In the literature, a candidate pole and candidate eigenvalue is associated to each facet of $\text{Newt}(f)$. For our purposes, it is important to extend these notions to a wider class of faces of $\text{Newt}(f)$. To be precise, let G be a proper face of $\text{Newt}(f)$ that contains the vector $\mathbf{1} = (1, \dots, 1)$ in its linear span, denoted $\text{span}(G)$. Let ψ_G be the unique linear function on $\text{span}(G)$ with value 1 on G . Then

$$\alpha_G := -\psi_G(\mathbf{1})$$

is the *candidate pole* associated to G , and $\exp(2\pi i\alpha_G)$ is the corresponding *candidate eigenvalue* of monodromy. We say that G *contributes* α_G as a candidate pole. If G' contains G as a face, then G' also contains $\mathbf{1}$ in its linear span and $\alpha_{G'} = \alpha_G$.

Definition 1.4.1. Let $\text{Contrib}(\alpha)$ be the set of faces of $\text{Newt}(f)$ that contribute the candidate pole α .

Then $\{\alpha \in \mathbb{Q} : \text{Contrib}(\alpha) \neq \emptyset\} \cup \{-1\}$ is a set of candidate poles for $Z_{\text{mot}}(T)$ [BN20, Corollary 8.3.4]. This set of candidate poles is standard in the literature. The key difference here is that we consider faces in $\text{Contrib}(\alpha)$ of arbitrary codimension, not just facets. This change in perspective is crucial in what follows.

Let C be a cone in Δ , the fan over the faces of $\text{Newt}(f)$. The rays of C are the union of rays through vertices in $\text{Newt}(f)$ and rays disjoint from $\text{Newt}(f)$ that contain a coordinate vector e_ℓ for some $1 \leq \ell \leq n$. In particular, for each ray of C , there is a corresponding distinguished generator: either the corresponding

vertex of $\text{Newt}(f)$, or the corresponding coordinate vector e_ℓ . We let $\text{Gen}(C)$ be the set of distinguished generators of the rays of C .

We say that a vertex A in G is an *apex* with *base direction* e_ℓ^* if $\langle e_\ell^*, A \rangle > 0$, and $\langle e_\ell^*, V \rangle = 0$ for all $V \in \text{Gen}(C_G)$ with $V \neq A$. In this case, $G \cap \{V \in \mathbb{R}_{\geq 0}^n : \langle e_\ell^*, V \rangle = 0\}$ is the corresponding *base* of G .

Definition 1.4.2. *A face G of $\text{Newt}(f)$ is B_1 if it has an apex A with base direction e_ℓ^* , and $\langle e_\ell^*, A \rangle = 1$.*

The notion of B_1 was introduced for simplicial facets in [LVP11, Definition 3]. For arbitrary facets, our definition agrees with [Que, Definition 1.1.7] but is more restrictive than [ELT22, Definition 3.1]. All of these definitions of B_1 -facets agree when $\text{Newt}(f)$ is simplicial. Note that the base direction e_ℓ^* determines the apex A . The converse is not true. A B_1 -face may have several apices, and when the face is not a facet, each of those apices can have multiple base directions. We introduce the following definition.

Definition 1.4.3. *A face G of $\text{Newt}(f)$ is UB_1 if it has an apex A with a unique base direction e_ℓ^* , and $\langle e_\ell^*, A \rangle = 1$.*

Theorems 1.4.6 and 1.4.7, show the importance of the notion of UB_1 -faces. Note that every B_1 -facet is UB_1 ; the distinction between B_1 and UB_1 is only relevant when considering higher codimension faces.

1.4.2. *Eigenvalue multiplicities and local h -polynomials.* The starting point for our work is the third author's nonnegative formula for the multiplicities of eigenvalues of monodromy at 0 when f is nondegenerate and $\text{Newt}(f)$ is convenient [Sta17, Section 6.3]. Specializing [Sta17, Theorem 6.20] from equivariant mixed Hodge numbers to equivariant multiplicities, one obtains a combinatorial formula with nonnegative integer coefficients for the multiplicities of the eigenvalues of monodromy on the reduced cohomology of \mathcal{F}_0 .

Assume that $\text{Newt}(f)$ is simplicial. If we forget the lattice structure of the fan Δ , we may view Δ as encoding a triangulation of a simplex, e.g., by slicing with a transverse hyperplane. Then the combinatorial formula for eigenvalues is a sum over cones C in Δ of a contribution that is a product of two nonnegative factors, one coming from Ehrhart theory (the number of lattice points in a polyhedral set). The other factor is the evaluation of the local h -polynomial $\ell(\Delta, C; t)$ at $t = 1$. These local h -polynomials were first introduced and studied by Stanley in the special case where $C = 0$ and later generalized by Athanasiadis, Nill, and Schepers [Ath12a, Nil12]. They have nonnegative, symmetric integer coefficients and naturally appear when applying the decomposition theorem to toric morphisms. See [Sta92, Theorem 5.2], [KS16, Theorem 6.1] and [dCMM18].

This formula for eigenvalue multiplicities in the convenient nondegenerate case offers fundamental advantages over earlier approaches to existence of eigenvalues. Whereas the formulas of A'Campo and Varchenko for zeta functions of monodromy typically involve a great deal of cancellation, the third author's formula is a sum of nonnegative terms. Moreover, for each compact face G in $\text{Contrib}(\alpha)$, there is a canonically associated *essential face* $E \subset G$. See Definition 3.3.1. Then the Ehrhart factor in the summand associated to C_E for the multiplicity of $\exp(2\pi i\alpha)$ is strictly positive. Thus, either $\exp(2\pi i\alpha)$ is an eigenvalue of monodromy or $\ell(\Delta, C_E; t)$ is zero. There are a number of simple sufficient conditions for the nonvanishing of $\ell(\Delta, C_E; t)$; for instance, if E meets the interior of the positive orthant, then $\ell(\Delta, C_E; 0) = 1$. The general problem of classifying when local h -polynomials vanish was posed by Stanley [Sta92, Problem 4.13]. See [dMGP⁺20] for a classification when $n \leq 4$ and $E = \emptyset$ and for partial results in higher dimensions.

1.4.3. *A nonnegative formula for nearby eigenvalues.* In Section 3, we extend the results of [Sta17] to the case where $\text{Newt}(f)$ is simplicial but not necessarily convenient. In this setting, the singularity of X_f at 0 may not be isolated, and the Milnor fibers at 0 and at nearby points may have cohomology in multiple positive degrees.

In this setting, we consider $\tilde{\chi}(\mathcal{F}_x) := \sum_i (-1)^i \tilde{H}^i(\mathcal{F}_x, \mathbb{C})$ as a virtual representation, where \tilde{H} denotes reduced cohomology. Now $\exp(2\pi i\alpha)$ has a multiplicity $\tilde{m}_x(\alpha)$, which may be positive or negative, as an

eigenvalue in this virtual representation. We consider these multiplicities not only at 0 but also at a general point x_I in each coordinate subspace \mathbb{A}^I contained in X_f . The idea of studying the eigenvalues at these points was first introduced when $n = 3$ in [LVP11] and further developed in [ELT22]. We give a nonnegative formula for the alternating sum

$$\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{m}_{x_I}(\alpha).$$

See Theorem 3.2.1 for a precise statement.

Theorem 3.2.1 implies, in particular, the remarkable fact that the corresponding alternating product of monodromy zeta functions is a polynomial, i.e.,

$$(4) \quad \prod_{\mathbb{A}^I \subset X_f} \left(\frac{\zeta_{x_I}(t)}{1-t} \right)^{(-1)^{n-1-|I|}} \in \mathbb{Z}[t].$$

From this perspective, the theorem provides a nonnegative formula for the vanishing order of this polynomial at $\exp(2\pi i\alpha)$. See Remark 3.2.3 for the precise formula.

Just as in the convenient case, this nonnegative formula is a sum over cones C in Δ , and each of the terms is once again an Ehrhart factor times $\ell(\Delta, C; 1)$. Moreover, for each compact $G \in \text{Contrib}(\alpha)$, we have an essential face $E \subset G$, and the Ehrhart factor in the C_E -summand for the multiplicity of $\exp(2\pi i\alpha)$ is strictly positive. We deduce the following corollary. See Corollary 3.3.2 for an equivalent statement.

Corollary 1.4.4. *Suppose $\text{Newt}(f)$ is simplicial and f is nondegenerate. Let G be a compact face in $\text{Contrib}(\alpha)$ with essential face E . If $\ell(\Delta, C_E; t)$ is nonzero, then $\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{m}_{x_I}(\alpha) > 0$. In particular, $\exp(2\pi i\alpha)$ is a nearby eigenvalue of monodromy (for reduced cohomology).*

This motivates a detailed study of necessary conditions for the vanishing of $\ell(\Delta, C_E; t)$ when E is the essential face associated to some compact face $G \in \text{Contrib}(\alpha)$. In this situation, we also have some additional structure which is crucial for our arguments. The face $C_G \setminus C_E \in \text{lk}_\Delta(C_E)$ admits what we call a *full partition*. See Lemma 3.3.4 and Definition 4.1.2.

1.4.4. *A necessary condition for the vanishing of the local h -polynomial.* Motivated by the results of Section 3, in Section 4 we undertake a detailed (and necessarily technical) investigation of the conditions under which $\ell(\Delta, C'; t)$ vanishes, where C' is a cone in Δ that is contained in a cone that admits a full partition. This section is self-contained and applies to any local h -polynomial of a geometric triangulation. See [LPS22] for further work on necessary conditions for the vanishing of the local h -polynomial in a more general setting, for quasi-geometric homology triangulations.

Recall that $\ell(\Delta, C'; t)$ is naturally identified with the Hilbert function of a module $L(\Delta, C')$ [Ath12b, Ath12a], as follows. Consider the ideal in the face ring $\mathbb{Q}[\text{lk}_\Delta(C')]$ generated by monomials x^C such that $C \sqcup C'$ meets the interior of the orthant $\mathbb{R}_{>0}^n$. Then $L(\Delta, C')$ is the image of this ideal in the quotient of $\mathbb{Q}[\text{lk}_\Delta(C')]$ by a special linear system of parameters. Thus $\ell(\Delta, C'; t) = 0$ if and only if every such monomial is contained in the ideal generated by a special linear system of parameters. When C admits a full partition, we can associate a distinguished monomial with image in $L(\Delta, C')$. We show that the image of this distinguished monomial specializes to a top degree cohomology class, given by the refined self-intersection of a compact T -invariant subvariety of half-dimension in a toric variety. An explicit calculation shows that this self-intersection is not zero. This calculation is purely combinatorial, and its proof constitutes the majority of the section.

Using this calculation, we prove the following theorem, which is an immediate consequence of Theorem 4.1.3. A cone C in $\text{lk}_\Delta(C')$ is a *U-pyramid* if it meets the interior of the positive orthant and there is a

ray $r \in C$ such that $(C \sqcup C') \setminus r$ is contained in a unique coordinate hyperplane in \mathbb{R}^n , i.e., C is a pyramid with a *unique* base direction with respect to the apex r in $\text{lk}_\Delta(C')$. See Definition 4.1.1.

Theorem 1.4.5. *Let Δ be a simplicial fan supported on $\mathbb{R}_{\geq 0}^n$. Let C' be a cone in Δ , and let $C \in \text{lk}_\Delta(C')$. If $\ell(\Delta, C'; t) = 0$ and C admits a full partition, then C is a U -pyramid.*

When G is compact and $C = C_G \setminus C_E \in \text{lk}_\Delta(C_E)$, the condition that G is UB_1 is equivalent to the condition that C is a U -pyramid. See Lemma 3.3.3. This leads to the following theorem, which is our main result on existence of eigenvalues.

Theorem 1.4.6. *Suppose $\text{Newt}(f)$ is simplicial and f is nondegenerate. Let $\alpha \in \mathbb{Q}$. Then either every face in $\text{Contrib}(\alpha)$ is UB_1 , or $\exp(2\pi i\alpha)$ is an eigenvalue of monodromy for the reduced cohomology of the Milnor fiber at the generic point of some coordinate subspace $\mathbb{A}^I \subset X_f$.*

Section 3 proves that this theorem follows from Theorem 1.4.5, whose proof is given in Section 4.

1.4.5. *The local formal zeta function and its candidate poles.* In Sections 5 and 6, we prove the following theorem, which is complementary to Theorem 1.4.6.

Theorem 1.4.7. *Suppose $\text{Newt}(f)$ is simplicial and f is nondegenerate. Let*

$$\mathcal{P} = \{\alpha \in \mathbb{Q} : \text{Contrib}(\alpha) \neq \emptyset\} \cup \{-1\}, \text{ and } \mathcal{P}' = \{\alpha \in \mathcal{P} : \alpha \notin \mathbb{Z}, \text{ every face in } \text{Contrib}(\alpha) \text{ is } UB_1\}.$$

Then $\mathcal{P} \setminus \mathcal{P}'$ is a set of candidate poles for $Z_{\text{mot}}(T)$.

Note that Theorem 1.1.1 follows directly from Theorems 1.4.6 and 1.4.7, using the fact that 1 is an eigenvalue of monodromy on $H^0(\mathcal{F}_0, \mathbb{C})$.

Our starting point for the proof of Theorem 1.4.7 is the formula for $Z_{\text{mot}}(T)$ in [BN20, Theorem 8.3.5], which expresses $Z_{\text{mot}}(T)$ as a sum over lattice points in the dual fan to $\text{Newt}(f)$. We introduce the *local formal zeta function* $Z_{\text{for}}(T)$, which is a power series over a polynomial ring that specializes to $Z_{\text{mot}}(T)$. The local formal zeta function depends only on $\text{Newt}(f)$, unlike $Z_{\text{mot}}(T)$ which depends on f . The advantage of working with $Z_{\text{for}}(T)$ is that an intersection of two sets of candidate poles of $Z_{\text{for}}(T)$ is a set of candidate poles (Lemma 5.3.5), so it suffices to show that, for each $\alpha \notin \mathbb{Z}$ such that $\text{Contrib}(\alpha)$ consists entirely of UB_1 -faces, there is a set of candidate poles for $Z_{\text{for}}(T)$ not containing α . Explicitly,

$$(5) \quad Z_{\text{for}}(T) = \sum_G Y_G \left((L-1)^{n-\dim G} \sum_{u \in \sigma_G^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)} \right),$$

where G varies over all nonempty compact faces of $\text{Newt}(f)$, σ_G denotes the dual cone to G , C° denotes the relative interior of a polyhedral cone C , N is a certain piecewise linear function, and Y_G and L are formal variables satisfying the following relations:

- (1) $Y_V = 1$ if V is a primitive vertex of $\text{Newt}(f)$, and
- (2) $Y_G + Y_F = \frac{(L-1)^{\dim F}}{1-L^{-1}T}$ if F is a compact B_1 -face with nonempty base G .

See Definition 5.3.1 for details. The key relation above is (2), which specializes to a natural relation in the $\hat{\mu}$ -equivariant Grothendieck ring of varieties. See Lemma 5.2.2.

Given a subset $C \subset \mathbb{R}_{\geq 0}^n$, we can define the *contribution* $Z_{\text{for}}(T)|_C$ of C to $Z_{\text{for}}(T)$ to be the same expression as the right hand side of (5), except that the second summation runs over $u \in \sigma_G^\circ \cap C \cap \mathbb{N}^n$. See (23). When F is a compact B_1 -face with nonempty base G and apex A in the direction e_ℓ^* , and $C' \subset \sigma_F^\circ$ is a nonzero rational polyhedral cone, then we deduce the following relation:

$$(6) \quad Z_{\text{for}}(T)|_{C^\circ} + Z_{\text{for}}(T)|_{(C')^\circ} = (L-1)^n \left(\sum_{u \in (C^\circ \cup (C')^\circ) \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A \rangle} \right),$$

where $C \subset \sigma_G$ is the cone spanned by C' and e_ℓ^* . See Lemma 5.4.1. The above equation (6) is a key technical tool underlying our strategy to show fakeness of poles, and it is analogous to a formula involving the local topological zeta function in [ELT22, Lemma 3.3]. For example, if $\text{Contrib}(\alpha)$ consists of a single B_1 (and hence UB_1) facet F with apex A , then one can deduce an expression for $Z_{\text{for}}(T)$ with no candidate pole at α by applying (6) with $C' = \sigma_G^\circ$, as G varies over all faces of G not containing A . This is analogous to approaches to showing fakeness of poles under certain assumptions for the local topological zeta function in [ELT22, LVP11] and the local naive motivic zeta function in [Que, Theorem A].

When we only assume that every face in $\text{Contrib}(\alpha)$ is B_1 , it may not be possible to extend the above approach. The key difficulty is that it may not be possible to choose a single base direction e_ℓ^* for all faces of $\text{Contrib}(\alpha)$. In our case, we assume that all elements of $\text{Contrib}(\alpha)$ are UB_1 . This assumption implies that we may choose base directions for elements of $\text{Contrib}(\alpha)$ satisfying a natural compatibility condition: to every face G in $\text{Contrib}(\alpha)$, we may assign a pair (A_G, e_G^*) such that G is B_1 with apex A_G and base direction e_G^* , and, if $G \subset G'$ and $A_G = A_{G'}$, then $e_G^* = e_{G'}^*$. See Definition 6.1.3 and Lemma 6.2.1.

We now sketch the remainder of the proof, and refer the reader to Section 6.1 for a more detailed overview. We first fix a minimal element M of $\text{Contrib}(\alpha)$ and reduce to considering only elements of $\text{Contrib}(\alpha)$ that contain M . See Section 6.3. We then use the above compatibility condition to construct a fan with support \mathbb{R}^n satisfying certain properties. See Section 6.5 and Section 6.6. In particular, we assign to every cone τ in the fan a coordinate vector e_τ^* such that, if $M \subset G$ and $\sigma_G \cap \tau \neq (0)$, then G is a B_1 -face with base direction e_τ^* . In this sense, we locally choose a single base direction. Then $Z_{\text{for}}(T)$ is the sum of all contributions $Z_{\text{for}}(T)|_{\tau^\circ}$ as τ varies over all cones of the fan. Analogously to the case when $\text{Contrib}(\alpha)$ is a single B_1 -facet, we then intersect each cone τ with the dual fan to $\text{Newt}(f)$ and repeatedly apply (6) to obtain an expression for $Z_{\text{for}}(T)|_{\tau^\circ}$ with no candidate pole α , allowing us to complete the proof of Theorem 1.4.7. See Section 6.4.

1.4.6. *Beyond the simplicial case.* In this introduction, we have stated our main results under the assumption that $\text{Newt}(f)$ is simplicial. However, both our arguments about eigenvalues and about poles are carried out in somewhat greater generality. In Section 7, we state our most general result on the local motivic monodromy conjecture for nondegenerate singularities (Theorem 7.1.1), which is sufficient to prove the local motivic monodromy conjecture in all cases when f is nondegenerate and $n = 3$ (Theorem 7.2.1).

Finally, observe that $\exp(2\pi i\alpha)$ appearing as a zero or pole of the monodromy zeta function implies that $\exp(2\pi i\alpha)$ is an eigenvalue of monodromy, but the converse is not true. When $\text{Newt}(f)$ is simplicial and X_f is nondegenerate, we prove that there is a set of candidate poles \mathcal{P} such that, for all $\alpha \in \mathcal{P} \setminus \mathbb{Z}$, $\exp(2\pi i\alpha)$ is a zero or pole of the monodromy zeta function at the generic point of some coordinate subspace $\mathbb{A}^I \subset X_f$. This stronger statement about when certain monodromy zeta functions are sufficient to detect eigenvalues is not true when $\text{Newt}(f)$ is not simplicial and $n \geq 4$. In such cases, there may be poles of local topological zeta functions such that $\exp(2\pi i\alpha)$ appears as a zero or pole of the monodromy zeta function only at points along strata that are properly contained in coordinate subspaces. See [ELT22, Example 7.5]. For one combinatorial approach to detecting such eigenvalues, see [Est21].

1.5. **Notation.** We now set up some additional notation which we will use for the remainder. With the exception of Section 7, the notation of the various sections are otherwise largely independent.

Let $F \subset \mathbb{R}_{\geq 0}^n$ be a rational polyhedron whose affine span does not contain the origin, and let $\text{span}(F)$ denote the linear span of F . Let ψ_F be the unique \mathbb{Q} -linear function on $\text{span}(F)$ with value 1 on F . The *lattice distance* ρ_F of F from the origin is the smallest positive integer ρ_F such that $\rho_F \psi_F$ is a \mathbb{Z} -linear function. If $F \subset G$ is an inclusion of such rational polyhedra, then ρ_F divides ρ_G .

A face of $\text{Newt}(f)$ is *interior* if it meets $\mathbb{R}_{> 0}^n$. The functions ψ_F , for F a face of a proper interior face of $\text{Newt}(f)$, assemble into a function ψ on $\mathbb{R}_{\geq 0}^n$ that is piecewise linear with respect to Δ .

For a polyhedral cone C , let ∂C denote its boundary, defined to be the union of all faces of C of dimension strictly less than $\dim C$. Let $C^\circ = C \setminus \partial C$ denote the relative interior of a polyhedral cone. A nonzero vector v in \mathbb{Z}^n is *primitive* if it generates the group $\mathbb{R}v \cap \mathbb{Z}^n$. Recall that C is *simplicial* if it is a pointed cone generated by $\dim C$ rays. For a set of vectors S in \mathbb{R}^n , let $\langle S \rangle$ denote the cone that they span.

A *geometric triangulation* of a simplex is a subdivision of a geometric simplex into a union of geometric simplices that meet along shared faces.

For a positive integer ℓ , we write $[\ell] = \{1, \dots, \ell\}$.

Acknowledgments. We thank M. Mustaș and J. Nicaise for helpful conversations related to the monodromy conjectures, and M. H. Quek for insightful comments on an earlier draft of this paper and telling us about Example 2.2.2. The work of ML is supported by an NDSEG fellowship, and the work of SP is supported in part by NSF grants DMS-2001502 and DMS-2053261.

2. EXAMPLES

2.1. Basic examples. Our first two examples are intended to serve as a guide to the main constructions in the paper. In these examples, $\text{Newt}(f)$ is simplicial and f is supported at the vertices of $\text{Newt}(f)$, so f is nondegenerate [BO16].

Below, $\tilde{E}(\mathcal{F}_x) \in \mathbb{Z}[\mathbb{Q}/\mathbb{Z}]$ is an alternative encoding of $\frac{\zeta_x(t)}{1-t}$ (see (7) in Section 3). The local formal zeta function $Z_{\text{for}}(T)$ lies in a quotient ring of $\mathbb{Z}[L, L^{-1}][Y_K : K \text{ nonempty compact face of } \text{Newt}(f)][[T]]$, where L, T, Y_K are formal variables. See Definition 5.3.1.

Example 2.1.1. Let $f(x_1, x_2) = x_2^2 - x_1^3$. Then f has an isolated cusp at 0, and $\text{Newt}(f)$ is convenient and has a unique compact facet F with vertices $v = (3, 0)$ and $w = (0, 2)$. Note that $\alpha_F = -5/6$, and F is not B_1 . Theorem 1.4.6 says that $\exp(2\pi i \alpha_F)$ is an eigenvalue of monodromy for f at 0.

Then Δ is the trivial fan, and $\ell(\Delta, C; t)$ equals 1 if $C = C_F$, and equals 0 otherwise. We have monodromy zeta function $\zeta_0(t) = \frac{1-t}{1-t+t^2}$, and $\tilde{E}(\mathcal{F}_0) = [1/6] + [5/6]$. The local formal zeta function is

$$Z_{\text{for}}(T) = \frac{(L-1)(Y_F L^{-5} T^6 + Y_v L^{-2} T^3(1+L^{-3} T^3) + Y_w L^{-1} T^2(1+L^{-2} T^2 + L^{-4} T^{-4}))}{1-L^{-5} T^6}.$$

The local motivic zeta function $Z_{\text{mot}}(T)$ is

$$\frac{(\mathbb{L}-1)\left(\left(\frac{(\mathbb{L}-1)\mathbb{L}^{-1}T}{1-\mathbb{L}^{-1}T} + [Y_F(1)]\right)\mathbb{L}^{-5}T^6 + [\mu_3]\mathbb{L}^{-2}T^3(1+\mathbb{L}^{-3}T^3) + [\mu_2]\mathbb{L}^{-1}T^2(1+\mathbb{L}^{-2}T^2 + \mathbb{L}^{-4}T^{-4})\right)}{1-\mathbb{L}^{-5}T^6},$$

where $Y_F(1)$ is an elliptic curve minus 6 points, with a free μ_6 -action, and $Y_F(1)/\mu_6$ is isomorphic to \mathbb{P}^1 minus 3 points. After simplification, the local naive motivic zeta function is

$$\frac{(\mathbb{L}-1)(\mathbb{L}^{-1}T^2 - \mathbb{L}^{-4}T^5 + \mathbb{L}^{-4}T^6 - \mathbb{L}^{-6}T^7)}{(1-\mathbb{L}^{-1}T)(1-\mathbb{L}^{-5}T^6)}.$$

For $p \notin \{2, 3\}$, f has good reduction mod p , and the local p -adic zeta function is

$$Z_{(p)}(s) = \frac{(p-1)(p^{5s+5} - p^{2s+2} + p^{s+2} - 1)}{(p^{s+1} - 1)(p^{6s+5} - 1)}.$$

The local topological zeta function is $Z_{\text{top}}(s) = \frac{(4s+5)}{(s+1)(6s+5)}$.

Example 2.1.2. Let $f(x_1, x_2, x_3) = x_1^2 - x_2^2 x_3$. Then X_f is the Whitney umbrella. There are three coordinate subspaces contained in X_f : $\mathbb{A}^\emptyset = \{0\}$, $\mathbb{A}^{\{2\}} = \{x_1 = x_3 = 0\}$ and $\mathbb{A}^{\{3\}} = \{x_1 = x_2 = 0\}$. The singular locus of X_f is $\mathbb{A}^{\{3\}}$. In particular, f does not have an isolated singularity at the origin and $\text{Newt}(f)$ is not convenient. The only maximal compact face of $\text{Newt}(f)$ is a 1-dimensional face F with vertices

$v = (2, 0, 0)$ and $w = (0, 2, 1)$. Note that F is UB_1 and $(1, 1, 1) \notin \text{span}(F)$. There are two unbounded interior facets: $F_1 = \{e_1^* + e_2^* = 2\} = F + \mathbb{R}_{\geq 0}e_3$ and $F_2 = \{e_1^* + 2e_3^* = 2\} = F + \mathbb{R}_{\geq 0}e_2$, with $F_1 \cap F_2 = F$.

Then $\alpha_{F_1} = -1$, and F_1 is not B_1 . Theorem 1.4.6 predicts that $\exp(2\pi i \alpha_{F_1}) = 1$ is a nearby eigenvalue of monodromy for reduced cohomology. Also, $\alpha_{F_2} = -3/2$, and F_2 is UB_1 . Then Theorem 1.4.7 predicts that there is a set of candidate poles for $Z_{\text{mot}}(T)$ not containing $-3/2$. In this particular case, the corresponding candidate eigenvalue $\exp(2\pi i \alpha_{F_2}) = -1$ is a nearby eigenvalue of monodromy.

The fan Δ has two maximal cones C_{F_1} and C_{F_2} intersecting in a unique interior 2-dimensional face C_F . We have

$$\ell(\Delta, C; t) = \begin{cases} 1 & \text{if } C = C_{F_1} \text{ or } C = C_{F_2}, \\ 1+t & \text{if } C = C_F, \\ 0 & \text{otherwise.} \end{cases}$$

The monodromy zeta function at a general point x_I of each coordinate subspace \mathbb{A}^I is given by: $\zeta_0(t) = (1-t)(1+t)$, $\zeta_{x_{\{2\}}} = 1-t$, $\zeta_{x_{\{3\}}} = 1$. Then $\prod_{\mathbb{A}^I \subset X_f} \left(\frac{\zeta_{x_I}(t)}{1-t} \right)^{(-1)^{n-1-|I|}} = 1-t^2$. Equivalently, $\tilde{E}(\mathcal{F}_0) = [1/2]$, $\tilde{E}(\mathcal{F}_{x_{\{2\}}}) = 0$, $\tilde{E}(\mathcal{F}_{x_{\{3\}}}) = -1$, and $\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) = 1 + [1/2]$. The local formal zeta function is

$$Z_{\text{for}}(T) = \frac{L^{-3}T^2(L-1)^3(Y_v(1-L^{-1}T) + L^{-1}T(1-L^{-1}))}{(1-L^{-1})^2(1-L^{-1}T)(1-L^{-2}T^2)}.$$

The local motivic zeta function is

$$Z_{\text{mot}}(T) = \frac{\mathbb{L}^{-3}T^2(\mathbb{L}-1)^3([\mu_2](1-\mathbb{L}^{-1}T) + \mathbb{L}^{-1}T(1-\mathbb{L}^{-1}))}{(1-\mathbb{L}^{-1})^2(1-\mathbb{L}^{-1}T)(1-\mathbb{L}^{-2}T^2)}.$$

After simplifying, the local naive motivic zeta function is

$$\frac{\mathbb{L}^{-1}T^2(\mathbb{L}-1)(1-\mathbb{L}^{-2}T)}{(1-\mathbb{L}^{-1}T)(1-\mathbb{L}^{-2}T^2)}.$$

For $p \neq 2$, f has good reduction mod p , and the local p -adic zeta function is

$$Z_{(p)}(s) = \frac{(p-1)(p^{s+2}-1)}{(p^{s+1}-1)^2(p^{s+1}+1)}.$$

The local topological zeta function is $Z_{\text{top}}(s) = \frac{(s+2)}{2(s+1)^2}$.

2.2. Counterexamples. We now present counterexamples to [ELT22, Conjecture 1.3], [Que, Question 5.1.8], [ELT22, Proposition 3.7], [ELT22, Lemma 5.18]. The polyhedral computations in these examples were done using polymake [GJ00], and the computation of the zeta functions can be verified using the Sage code of [VS12].

Example 2.2.1. Let

$$f(x_1, x_2, x_3, x_4, x_5, x_6) = x_1^8 + x_2^5 + x_3^{24} + x_4^{13} + x_5^{17} + x_6^{14} + x_3x_5x_6 + x_2^3x_4 + x_4x_5x_6 + x_1x_3^2x_4x_6.$$

Then f is a nondegenerate polynomial with an isolated singularity at 0 whose Newton polyhedron is simplicial and convenient, with sixteen compact facets and ten vertices. There are five compact facets containing the face with vertices $\{(8, 0, 0, 0, 0, 0), (0, 5, 0, 0, 0, 0), (0, 0, 1, 0, 1, 1), (0, 3, 0, 1, 0, 0)\}$, each of which contributes the candidate pole $-69/40$. All of these facets are B_1 , and no other facets contribute $-69/40$. Two of these facets are obtained by adding either $\{(0, 0, 0, 0, 0, 14), (0, 0, 0, 1, 1, 1)\}$ or $\{(0, 0, 0, 0, 0, 14), (1, 0, 2, 1, 0, 1)\}$ to the above face. The existence of these two facets implies that the condition in [ELT22, Conjecture 1.3] is not satisfied, so the conjecture predicts that $\exp(2\pi i(-69/40))$ is an eigenvalue of monodromy. But this is not one of the 1912 eigenvalues of monodromy at the origin.

The local topological zeta function of f is $Z_{\text{top}}(s) = -\frac{6142656s^3 - 2948088s^2 - 93769198s - 115234075}{17(s+1)(104s+157)^2(168s+275)}$, which does not have $-69/40$ as a pole. One can deduce from Theorem 6.1.2 that there is a set of candidate poles for $Z_{\text{mot}}(T)$ which does not contain $-69/40$.

Example 2.2.2. Let

$$f(x_1, x_2, x_3, x_4) = x_1^{10} + x_2^5 + x_3^6 + x_4^{10} + x_2x_4^2 + x_2x_3 + x_1x_2 + x_1x_3.$$

Then f is a nondegenerate polynomial with an isolated singularity at the origin whose Newton polyhedron is simplicial and convenient, with six compact facets and eight vertices. There are two compact facets containing the face with vertices $\{(0, 0, 0, 10), (0, 1, 0, 2), (1, 0, 1, 0)\}$, each of which contributes the candidate pole $-19/10$. Both of these facets are B_1 , and they are obtained by adding either $(0, 1, 1, 0)$ or $(1, 1, 0, 0)$. The condition in [ELT22, Conjecture 1.3] is not satisfied, so their conjecture predicts that $\exp(2\pi i(-19/10))$ is an eigenvalue of monodromy. However, the only eigenvalues of monodromy are $\{1, \pm i\}$. This contradicts the claim in [ELT22] that they have proven [ELT22, Conjecture 1.3] for polynomials in four variables.

The local topological zeta function of f is $Z_{\text{top}}(s) = \frac{s+7}{(s+1)(4s+7)}$, which does not have $-19/10$ as a pole. One can deduce from Theorem 1.4.7 that there is a set of candidate poles for $Z_{\text{mot}}(T)$ which does not contain $-19/10$. We thank M. H. Quek for informing us that there are counterexamples to [ELT22, Conjecture 1.3] in four variables.

Example 2.2.3. Let

$$f(x_1, x_2, x_3, x_4, x_5) = x_1^{21} + x_2^{22} + x_3^{24} + x_4^6 + x_5^{12} + x_1x_2 + x_2x_3^6x_5^5 + x_3x_4x_5 + x_3^2x_5^9.$$

Then f is a nondegenerate polynomial with an isolated singularity at the origin whose Newton polyhedron is simplicial and convenient, with ten compact facets and nine vertices. Every compact facet contains the face with vertices $\{(1, 1, 0, 0, 0), (0, 0, 1, 1, 1)\}$, and the candidate pole of every facet is -2 . These ten facets have a choice of compatible apices in the sense of [Que, Definition 5.1.5], so [Que, Question 5.1.8] predicts that $\{-1\}$ is a set of candidate poles for the local naive motivic zeta function.

The local topological zeta function of f is $Z_{\text{top}}(s) = \frac{7s^2 + 45s + 96}{24(s+1)(s+2)^2}$, so any set of candidate poles for the local naive motivic zeta function contains -2 . When $n = 6$, there are counterexamples to [Que, Question 5.1.8] whose candidate pole is not an integer.

Example 2.2.4. In [ELT22, Definition 3.1], the authors give a different definition of a B_1 -facet when the facet is non-compact. They say that a facet F with $\text{Unb}(C_F) \neq \emptyset$ is a B_1 -facet of non-compact type if the image \bar{F} of F under the projection $\mathbb{R}^n \rightarrow \mathbb{R}^n / \langle \text{Unb}(C_F) \rangle$ is a B_1 -facet. Then [ELT22, Proposition 3.7] claims that if a pole $\alpha \neq -1$ is contributed only by a single B_1 -facet, then α is not a pole of $Z_{\text{top}}(s)$. Consider the nondegenerate polynomial

$$f(x_1, x_2, x_3, x_4) = x_1x_3 + x_2x_3 + x_2x_4^5 + x_4^6.$$

Then $\text{Newt}(f)$ has four vertices and eight facets, one of which is compact. There is B_1 -facet F of non-compact type with vertices $\{(1, 0, 1, 0), (0, 1, 1, 0), (0, 1, 0, 5)\}$ and $\text{Unb}(C_F) = \{e_1, e_2\}$ whose candidate pole is $-6/5$. It is the only facet whose candidate pole is $-6/5$, so [ELT22, Proposition 3.7] claims that $-6/5$ is not a pole of $Z_{\text{top}}(s)$. In fact, $Z_{\text{top}}(s) = \frac{6}{(s+1)(5s+6)}$.

At the origin, the monodromy zeta function is 1. The singular locus of X_f is the set of points of the form $\{(c, -c, 0, 0)\}$. At any $c \neq 0$, the monodromy zeta function is equal to $-(t-1)(t^4 + t^3 + t^2 + t + 1)$, so $\exp(2\pi i(-6/5))$ is an eigenvalue of monodromy.

Example 2.2.5. Let

$$f(x_1, x_2, x_3, x_4) = x_1^{45} + x_2^{45} + x_3^{23} + x_4^{18} + x_1x_4 + 2x_2x_4.$$

Then f is a nondegenerate polynomial with an isolated singularity at the origin whose Newton polyhedron is convenient, with two compact facets and six vertices. The vertices

$$\{(45, 0, 0, 0), (0, 45, 0, 0), (0, 0, 23, 0), (1, 0, 0, 1), (0, 1, 0, 1)\}$$

form a facet F . Then F is not a “ B -facet” in the sense of [ELT22, Definition 3.10], but any four affinely independent vertices form a B_1 facet. This contradicts [ELT22, Lemma 5.18].

The candidate pole associated to F is $-1103/1035$. The topological zeta function is $Z_{\text{top}}(s) = \frac{67s+1103}{(s+1)(1035s+1103)}$, so $-1103/1035$ is a pole of the topological zeta function. The lattice point $(1, 44, 0, 0) = 1/45(45, 0, 0, 0) + 44/45(0, 45, 0, 0)$ is contained in F , and the lattice simplex with vertices

$$\{(1, 44, 0, 0), (0, 0, 23, 0), (1, 0, 0, 1), (0, 1, 0, 1)\}$$

is inscribed in F and is not B_1 . This implies that F is not pseudo- UB_1 , and so $\exp(2\pi i(-1103/1035))$ is an eigenvalue of monodromy by Theorem 7.1.1(1).

3. A NONNEGATIVE FORMULA FOR NEARBY EIGENVALUES

Here and throughout, $f \in \mathbb{k}[x_1, \dots, x_n]$ is a nondegenerate polynomial that vanishes at 0. In this section, we do not assume that $\text{Newt}(f)$ is simplicial. For a geometric point x in the hypersurface X_f , we write $\tilde{m}_x(\alpha)$ to denote the multiplicity of $\exp(2\pi i\alpha)$ in the virtual representation $\tilde{\chi}(\mathcal{F}_x) := \sum_i (-1)^i \tilde{H}^i(\mathcal{F}_x, \mathbb{C})$, where \tilde{H} denotes reduced cohomology. We define

$$(7) \quad \tilde{E}(\mathcal{F}_x) := \sum_{[\alpha] \in \mathbb{Q}/\mathbb{Z}} \tilde{m}_x(\alpha) [\alpha] \in \mathbb{Z}[\mathbb{Q}/\mathbb{Z}],$$

where $\mathbb{Z}[\mathbb{Q}/\mathbb{Z}]$ is the group algebra of \mathbb{Q}/\mathbb{Z} . For example, when $\text{Newt}(f)$ is convenient, then X_f has an isolated singularity at the origin [Kou76, 1.13(ii)]. In this case, the Milnor fiber \mathcal{F}_0 has the homotopy type of a wedge sum of $(n-1)$ -dimensional spheres [Mil68], and $\exp(2\pi i\alpha)$ is a nearby eigenvalue for reduced cohomology if and only if the coefficient $\tilde{m}_0(\alpha)$ is nonzero.

Note that $\tilde{E}(\mathcal{F}_x)$ encodes information equivalent to that in the monodromy zeta function $\zeta_x(t)$. This alternative notation is useful when studying how the monodromy action interacts with finer invariants of the cohomology of the Milnor fiber, such as its mixed Hodge structure [Sta17, Section 6.3]. The additive structure of $\tilde{E}(\mathcal{F}_x)$ and the restriction to reduced cohomology is also more natural from a combinatorial perspective: it aligns with standard formulas for local h -polynomials and thereby leads to the nonnegative formula for nearby eigenvalues that we prove here.

Given $\beta \in \mathbb{Q}$, let $D(\beta) \in \mathbb{Z}_{>0}$ be the denominator of β , written as a reduced fraction. Fix $M \in \mathbb{Z}_{>0}$, and consider the following \mathbb{Z} -module homomorphism:

$$\Psi_M: \mathbb{Z}[\mathbb{Q}/\mathbb{Z}] \rightarrow \mathbb{Z}[\mathbb{Q}/\mathbb{Z}], \text{ given by } \Psi_M([\beta]) = \begin{cases} [\beta] & \text{if } M \text{ divides } D(\beta), \\ 0 & \text{otherwise.} \end{cases}$$

For example, when $M = 1$, then Ψ_M is the identity map.

Let x_I denote a general point in the coordinate subspace $\mathbb{A}^I \subset \mathbb{A}^n$ for $I \subset [n]$. Under appropriate conditions, the main result of this section is a formula with nonnegative integer coefficients for

$$\Psi_M \left(\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) \right).$$

For example, when $\text{Newt}(f)$ is simplicial or convenient, we will see that we obtain a non-negative formula when $M = 1$ (see Remark 3.2.2). The set of coordinate subspaces contained in X_f depends only on $\text{Newt}(f)$. Explicitly, $\mathbb{A}^I \subset X_f$ if and only if $\mathbb{R}_{\geq 0}^I \cap \text{Newt}(f) = \emptyset$.

3.1. The local h -polynomial. Let Δ' be a simplicial fan with support $\mathbb{R}_{\geq 0}^n$. If we forget the lattice structure of the fan Δ' , we may view Δ' as encoding a triangulation of a simplex, e.g., by slicing with a transverse hyperplane. For C' a cone in Δ' , let $\sigma(C')$ be the smallest face of the cone $\mathbb{R}_{\geq 0}^n$ containing C' . Set

$$e(C') := \dim \sigma(C') - \dim C'.$$

We use the following definition of the local h -polynomial (see, for example, [KS16, Lemma 4.12]).

Definition 3.1.1. *Let Δ' be a simplicial fan with support $\mathbb{R}_{\geq 0}^n$, and let C' be a cone in Δ' . Then the local h -polynomial $\ell(\Delta', C'; t)$ is defined as*

$$\ell(\Delta', C'; t) := \sum_{C' \subset C \in \Delta'} (-1)^{\text{codim}(C)} t^{\text{codim}(C') - e(C)} (t-1)^{e(C)}.$$

The local h -polynomial has several important properties. Most important for our purposes is that its coefficients are nonnegative integers [Ath12a]. We refer the reader to [KS16] for more details and a more general setting. Observe that

$$(8) \quad \ell(\Delta', C'; 1) := \sum_{\substack{C' \subset C \in \Delta' \\ e(C)=0}} (-1)^{\text{codim}(C)}.$$

Recall that Δ denotes the fan over the faces of $\text{Newt}(f)$. The following definition extends the notions of $\text{Gen}(C)$, $\text{Vert}(C)$ and $\text{Unb}(C)$, for C a cone in Δ .

Definition 3.1.2. *Let C' be a polyhedral cone contained in a cone of Δ , such that every ray of C' not in Δ intersects $\partial \text{Newt}(f)$ at a lattice point. Let $\text{Gen}(C') = \text{Vert}(C') \cup \text{Unb}(C')$ be the set of distinguished lattice point generators of the rays of C' defined as follows:*

$$\begin{aligned} \text{Vert}(C') &= \{w \in \mathbb{Z}^n : \{w\} = r \cap \partial \text{Newt}(f) \text{ for some ray } r \text{ of } C'\}, \text{ and} \\ \text{Unb}(C') &= \{e_i : e_i \in C', \mathbb{R}_{\geq 0} e_i \cap \text{Newt}(f) = \emptyset\}. \end{aligned}$$

When C' is simplicial, we need the following definition.

Definition 3.1.3. *Let C' be a simplicial cone contained in a cone of Δ , such that every ray of C' not in Δ intersects $\partial \text{Newt}(f)$ at a lattice point. Let $\text{Gen}(C') = \{w_1, \dots, w_r\}$, and define finite sets $\text{Box}_{C'}^\circ$ and $\text{Box}_{C'}$ as follows:*

$$\text{Box}_{C'}^\circ := \left\{ w \in \mathbb{Z}^n : w = \sum_{i=1}^r \lambda_i w_i, 0 < \lambda_i < 1 \right\} \quad \text{and} \quad \text{Box}_{C'} := \left\{ w \in \mathbb{Z}^n : w = \sum_{i=1}^r \lambda_i w_i, 0 \leq \lambda_i < 1 \right\}.$$

When $C' = \{0\}$, then $\text{Box}_{C'}^\circ = \text{Box}_{C'} = \{0\}$.

Note that $\text{Box}_{C'} = \cup_{C \subset C'} \text{Box}_C^\circ$, and $\text{Box}_{C'} = \{0\}$ if $\text{Gen}(C') \subset \{e_1, \dots, e_n\}$.

Recall that if $F \subset \mathbb{R}_{\geq 0}^n$ is a rational polyhedron whose affine span $\text{aff}(F)$ does not contain the origin, we write ρ_F for the lattice distance of F to the origin. If F is a lattice polytope, then we may consider the *normalized volume* $\text{Vol}(F) \in \mathbb{Z}_{>0}$ of F , i.e., the Euclidean volume on $\text{aff}(F)$ scaled such that the volume of a unimodular lattice simplex is 1. When $F = \emptyset$, $\rho_F = \text{Vol}(F) = 1$. We will need the following basic lemma.

Lemma 3.1.4. *Let C' be a simplicial cone contained in a cone of Δ such that every ray of C' not in Δ intersects $\partial \text{Newt}(f)$ at a lattice point. Let F be the convex hull of the elements of $\text{Gen}(C')$, and let ϕ be the linear function on C' with value 1 on F . Then*

$$\sum_{w \in \text{Box}_{C'}} [-\phi(w)] = \text{Vol}(F) \sum_{i=0}^{\rho_F-1} [i/\rho_F].$$

Proof. The result follows from the fact that ϕ induces a group homomorphism:

$$\tilde{\phi}: (\text{span}(C') \cap \mathbb{Z}^n) / (\mathbb{Z}w_1 + \cdots + \mathbb{Z}w_r) \rightarrow \mathbb{Q}/\mathbb{Z},$$

where the domain is a finite set in bijection with $\text{Box}_{C'}$, $\text{im}(\tilde{\phi}) = \frac{1}{\rho_F} \mathbb{Z}/\mathbb{Z}$ and $|\ker(\tilde{\phi})| = \text{Vol}(F)$. See also [Sta17, Examples 4.12-4.13]. \square

3.2. Nearby eigenvalues along coordinate subspaces. We now state our nonnegative formula for the multiplicity of nearby eigenvalues along coordinate subspaces. We first introduce some notation.

Recall that ψ is the unique piecewise linear function on $\mathbb{R}_{\geq 0}^n$ with value 1 on all interior faces of $\partial \text{Newt}(f)$. Let C be a cone in Δ . If $C = C_F$ for some face F of $\text{Newt}(f)$, then we set $\rho_C := \rho_F$ to be the lattice distance of F to the origin. Otherwise, we set $\rho_C := 1$. Equivalently, ρ_C is the smallest positive integer such that $\rho_C \psi|_C$ is the restriction of a \mathbb{Z} -linear function on $\text{span}(C)$. Observe that if $C \subset \tilde{C} \in \Delta$, then ρ_C divides $\rho_{\tilde{C}}$.

Fix $M \in \mathbb{Z}_{>0}$. Let Δ_M be the (possibly empty) subfan of Δ consisting of all maximal cones C in Δ such that M divides ρ_C , together with all the faces of C . Observe that if C is a cone in Δ and M divides ρ_C , then all cones in Δ containing C lie in Δ_M .

If Δ' is a fan refining Δ , let Δ'_M denote the restriction of Δ' to Δ_M . Given a cone C' contained in a cone of Δ , let $\tau(C')$ denote the smallest cone in Δ containing C' .

Theorem 3.2.1. *Assume that f is nondegenerate. Let $M \in \mathbb{Z}_{>0}$ and let Δ' be a simplicial fan refining Δ . Assume that every ray of $\Delta' \setminus \Delta$ intersects the boundary of $\text{Newt}(f)$ at a lattice point, and $\text{Unb}(C') = \text{Unb}(\tau(C'))$ for all C' in Δ'_M . Then*

$$(9) \quad \Psi_M \left(\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) \right) = \Psi_M \left(\sum_{C' \in \Delta'} \ell(\Delta', C'; 1) \sum_{w \in \text{Box}_{C'}^\circ} [-\psi(w)] \right).$$

Remark 3.2.2. We consider two important special cases when such a Δ' exists when $M = 1$. Firstly, if $\text{Newt}(f)$ is simplicial, then the hypotheses of the theorem hold with $\Delta' = \Delta$. Secondly, if $\text{Newt}(f)$ is convenient, then there exists a simplicial fan Δ' that refines Δ and has the same rays as Δ . In this case, as $\text{Unb}(C') = \text{Unb}(\tau(C')) = \emptyset$ for all C' in Δ' , the hypotheses of the theorem hold.

As a simple example where the hypotheses fail, suppose there exists $C \in \Delta$ with $\dim C = 3$ and $|\text{Vert}(C)| = |\text{Unb}(C)| = 2$. Then there is no simplicial refinement of C in which all maximal cones contain $\text{Unb}(C)$.

Remark 3.2.3. When $M = 1$, (9) can be restated in terms of monodromy zeta functions at x_I , as follows:

$$\prod_{\mathbb{A}^I \subset X_f} \left(\frac{\zeta_{x_I}(t)}{1-t} \right)^{(-1)^{n-1-|I|}} = \prod_{C' \in \Delta'} \prod_{w \in \text{Box}_{C'}^\circ} (1 - \exp(-2\pi i \psi(w)) t)^{\ell(\Delta', C'; 1)}.$$

For the remainder, we work in terms of $\tilde{E}(\mathcal{F}_{x_I})$ rather than the corresponding monodromy zeta function.

We now give three examples which show that the sum $\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I})$ appearing in Theorem 3.2.1 can fail to detect nearby eigenvalues and can have strictly negative coefficients.

Example 3.2.4. In Example 2.2.4, consider the facet F with candidate pole $\alpha = -6/5$. It was shown that there exists $x \in X_f$ arbitrarily close to the origin such that $\tilde{m}_x(\alpha)$ is nonzero. On the other hand, X_f is smooth at x_I for $I \neq \emptyset$, so $\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) = -\tilde{E}(\mathcal{F}_0) = 1$. In particular, $[-\alpha]$ does not appear. See also Example 3.4.9 below.

Example 3.2.5. The following example appeared in [ELT22, Example 7.4]. Consider the nondegenerate polynomial $f(x_1, x_2, x_3, x_4) = x_3^6 + x_2^4 x_3^5 + (x_1^2 + x_4^2) x_2^{13} x_3^2$. Let $F \subset \mathbb{R}^4$ be the 3-dimensional lattice simplex with vertices $w_1 = (0, 0, 6, 0)$, $w_2 = (0, 4, 5, 0)$, $w_3 = (2, 13, 2, 0)$, $w_4 = (0, 13, 2, 2)$. Then F is the unique compact facet of $\text{Newt}(f)$, and it has candidate pole $\alpha = -1/3$. The authors showed that $\tilde{E}(\mathcal{F}_0) = -(\sum_{0 \leq i < 24} [i/24] - \sum_{1 \leq i < 6} [i/6])$, and $[\alpha]$ does not appear in $\tilde{E}(\mathcal{F}_{x_I})$ for a general point x_I in any $\mathbb{A}^I \subset X_f$, but that there exists $x \in X_f$ arbitrarily close to the origin such that $\tilde{E}(\mathcal{F}_x)$ contains $[\alpha]$. We have

$$(-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) = \begin{cases} \sum_{0 \leq i < 24} [i/24] - \sum_{1 \leq i < 6} [i/6] & \text{if } I = \emptyset \\ \sum_{1 \leq i < 5} [i/5] & \text{if } I = \{2\} \\ -\sum_{0 \leq i < 78} [i/78] + \sum_{1 \leq i < 6} [i/6] & \text{if } I = \{1\} \text{ or } \{4\} \\ \sum_{0 \leq i < 78} [i/78] - \sum_{1 \leq i < 6} [i/6] & \text{if } I = \{1, 4\} \\ -[1/2] & \text{if } I = \{1, 2\} \text{ or } \{2, 4\} \\ [1/2] & \text{if } I = \{1, 2, 4\}. \end{cases}$$

In particular, $\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I})$ has strictly negative integer coefficients. See also Example 3.4.10 below.

Example 3.2.6. Consider f as a polynomial in $n+1$ variables, i.e., $f \in \mathbb{k}[x_1, \dots, x_n] \subset \mathbb{k}[x_1, \dots, x_n, x_{n+1}]$. Then one can verify that both sides of (9) are identically zero. Geometrically, if $I \subset \{1, \dots, n\}$ and $I' = I \cup \{n+1\}$, then $(-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) + (-1)^{n-1-|I'|} \tilde{E}(\mathcal{F}_{x_{I'}}) = 0$.

We will deduce Theorem 3.2.1 from Varchenko's formula for the monodromy zeta function of a nondegenerate singularity [Var76, Theorem 4.1]. First, recall that ρ_C is the lattice distance from the origin to F if $C = C_F$, and is 1 otherwise. Recall that Δ_M is the (possibly empty) subfan of Δ consisting of all maximal cones C in Δ such that M divides ρ_C , together with all the faces of C .

Lemma 3.2.7. *Let $M \in \mathbb{Z}_{>0}$. If C is a cone in $\Delta \setminus \Delta_M$, then $\Psi_M([i/\rho_C]) = 0$ for any $i \in \mathbb{Z}$. In particular, $\Psi_M([- \psi(w)]) = 0$ for all $w \in C \cap \mathbb{Z}^n$.*

Proof. If $\Psi_M([i/\rho_C]) \neq 0$, then M divides $D(i/\rho_C)$. Therefore for \tilde{C} a maximal cone in Δ containing C , M divides $\rho_{\tilde{C}}$ and hence $C \in \Delta_M$, a contradiction. The second statement follows since the restriction of $\rho_C \psi$ to C is the restriction of a \mathbb{Z} -linear function on $\text{span}(C)$ for all cones C in Δ . \square

Observe that when Δ_M is empty, the proposition below states that $\Psi_M(\tilde{E}(\mathcal{F}_0)) = 0$.

Proposition 3.2.8. *Let $M \in \mathbb{Z}_{>0}$ and let Δ'_M be a simplicial fan refining Δ_M . Assume that every ray of $\Delta'_M \setminus \Delta_M$ intersects the boundary of $\text{Newt}(f)$ at a lattice point, and $\text{Unb}(C') = \text{Unb}(\tau(C'))$ for all C' in Δ'_M . Then*

$$\Psi_M(\tilde{E}(\mathcal{F}_0)) = \sum_{\substack{C' \in \Delta'_M \\ \text{Unb}(C') = \emptyset \\ e(C') = 0}} (-1)^{\dim C' + 1} \Psi_M \left(\sum_{w \in \text{Box}_{C'}} [-\psi(w)] \right).$$

Proof. Suppose that C in Δ satisfies $\text{Unb}(C) = \emptyset$. Then $C = C_F$ for some bounded face F of $\text{Newt}(f)$, and we may set $\text{Vol}(C) := \text{Vol}(F)$. For example, when $C = \{0\}$, then $F = \emptyset$ and $\text{Vol}(C) = 1$. With this notation, Varchenko's formula for the monodromy zeta function [Var76, Theorem 4.1] states that

$$(10) \quad \tilde{E}(\mathcal{F}_0) = \sum_{\substack{C \in \Delta \\ \text{Unb}(C) = \emptyset \\ e(C) = 0}} (-1)^{\dim C + 1} \text{Vol}(C) \sum_{i=0}^{\rho_C - 1} [i/\rho_C].$$

Let $S = \{C \in \Delta_M : \text{Unb}(C) = \emptyset, e(C) = 0\}$. Applying Ψ_M to both sides of the above equation, and using Lemma 3.2.7, we obtain the equation

$$\Psi_M(\tilde{E}(\mathcal{F}_0)) = \sum_{C \in S} (-1)^{\dim C + 1} \text{Vol}(C) \Psi_M \left(\sum_{i=0}^{\rho_C - 1} [i/\rho_C] \right).$$

Let $C \in S$ and set $S'_C = \{C' \in \Delta'_M : \tau(C') = C, \dim C' = \dim C\}$. Let $C' \in S'_C$. By assumption, $\text{Unb}(C') = \text{Unb}(C) = \emptyset$. Then $C = C_F$ for some lattice polytope F , and C' is the cone over a lattice polytope $G \subset F$. Define $\text{Vol}(C') := \text{Vol}(G)$ and $\rho_{C'} := \rho_G$. Note that $\dim C' = \dim C$ implies that $\rho_{C'} = \rho_C$. By the additivity of normalized volume, we have

$$\Psi_M(\tilde{E}(\mathcal{F}_0)) = \sum_{C \in S} (-1)^{\dim C + 1} \Psi_M \left(\sum_{i=0}^{\rho_C - 1} [i/\rho_C] \right) \sum_{C' \in S'_C} \text{Vol}(C').$$

Since $\dim C' = \dim C$, the condition $e(C') = 0$ is equivalent to the condition $e(C) = 0$. Let $S' = \{C' \in \Delta'_M : \text{Unb}(C') = \emptyset, e(C') = 0\}$. Then rearranging the above equation gives

$$\Psi_M(\tilde{E}(\mathcal{F}_0)) = \sum_{C' \in S'} (-1)^{\dim C' + 1} \Psi_M \left(\text{Vol}(C') \sum_{i=0}^{\rho_{C'} - 1} [i/\rho_{C'}] \right)$$

By Lemma 3.1.4, we obtain our desired result. \square

We also need the following remark. Given $c = (c_1, \dots, c_n) \in \mathbb{k}^n$, let $f_c(x_1, \dots, x_n) := f(x_1 + c_1, \dots, x_n + c_n)$. Consider a coordinate subspace $\mathbb{A}^I \subset X_f$, and a general point x_I in \mathbb{A}^I . Let $J = [n] \setminus I$, and consider the projection map $\text{pr}_J: \mathbb{R}^n \rightarrow \mathbb{R}^J$ and the polyhedron $\text{Newt}(f)_J := \text{pr}_J(\text{Newt}(f)) \subset \mathbb{R}^J$. Let g be a nondegenerate polynomial with Newton polyhedron P and Milnor fiber $\hat{\mathcal{F}}_0$ at the origin. By [Var76, Theorem 4.1], $\tilde{E}(\hat{\mathcal{F}}_0)$ depends only on P , and not on the choice of g . We set

$$\tilde{E}(P) := \tilde{E}(\hat{\mathcal{F}}_0).$$

Remark 3.2.9. With the notation above, \mathcal{F}_{x_I} is the Milnor fiber of f_{x_I} at the origin. It follows from [ELT22, Proposition 7.2] and its proof that f_{x_I} is nondegenerate with Newton polyhedron equal to $\text{Newt}(f)_J \times \mathbb{R}_{\geq 0}^I$. Then

$$(11) \quad \tilde{E}(\mathcal{F}_{x_I}) = \tilde{E}(\text{Newt}(f)_J \times \mathbb{R}_{\geq 0}^I) = \tilde{E}(\text{Newt}(f)_J).$$

We deduce that if the coefficient of $[\alpha]$ in $\tilde{E}(\text{Newt}(f)_J)$ is nonzero for some such choice of I , then $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy (for reduced cohomology).

Recall that for a set of vectors S , $\langle S \rangle$ denotes the cone that they span. We now prove Theorem 3.2.1. Our strategy is to apply Proposition 3.2.8 to each coordinate projection of $\text{Newt}(f)$.

Proof of Theorem 3.2.1. Consider a coordinate subspace \mathbb{A}^I in X_f . Let $J = [n] \setminus I$, and consider the projection map $\text{pr}_J: \mathbb{R}^n \rightarrow \mathbb{R}^J$ and the polyhedron $\text{Newt}(f)_J = \text{pr}_J(\text{Newt}(f)) \subset \mathbb{R}^J$. By (11), $\tilde{E}(\mathcal{F}_{x_I}) = \tilde{E}(\text{Newt}(f)_J)$, where $\tilde{E}(\text{Newt}(f)_J)$ is the invariant \tilde{E} applied to the Milnor fiber at the origin of any nondegenerate polynomial with Newton polyhedron $\text{Newt}(f)_J$. Our first goal is to apply Proposition 3.2.8 to compute $\Psi_M(\tilde{E}(\text{Newt}(f)_J))$.

If $C \subset \mathbb{R}_{\geq 0}^n$ is a cone, then we use the notation $C_J := \text{pr}_J(C)$. Let Δ_J be the fan over the faces of $\text{Newt}(f)_J$. Then $\Delta_J = \{C_J : C \in \Delta, \mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C) \rangle\}$. Let $\Delta_{J,M}$ be the subfan of Δ_J consisting of all maximal cones C_J in Δ_J such that M divides ρ_{C_J} , together with all the faces of C_J . Observe that if $C \in \Delta$ and $\mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C) \rangle$, then $\rho_C = \rho_{C_J}$. It follows that $\Delta_{J,M} = \{C_J : C \in \Delta_M, \mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C) \rangle\}$.

If $\Delta_{J,M}$ is empty, then Proposition 3.2.8 implies that $\Psi_M(\tilde{E}(\text{Newt}(f)_J)) = 0$. Assume that $\Delta_{J,M}$ is nonempty. Then $\mathbb{R}_{\geq 0}^I \in \Delta_M$. In order to apply Proposition 3.2.8, we want to construct a simplicial refinement of $\Delta_{J,M}$. Since each ray in $\Delta' \setminus \Delta$ intersects $\text{Newt}(f)$, no such ray is contained in $\mathbb{R}_{\geq 0}^I$, and hence $\mathbb{R}_{\geq 0}^I \in \Delta'_M$. Consider the simplicial fan $\Delta'_{J,M} = \{C'_J : C' \in \Delta'_M, \mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C') \rangle\}$. This is the star of $\mathbb{R}_{\geq 0}^I$ in Δ'_M , and it follows from [CLS11, Exercise 3.4.8] that $\Delta'_{J,M}$ is a refinement of $\Delta_{J,M}$.

We next verify that $\Delta'_{J,M}$ satisfies the hypotheses of Proposition 3.2.8. Firstly, the intersection of a ray of $\Delta'_{J,M} \setminus \Delta_{J,M}$ with the boundary of $\text{Newt}(f)_J$ is the image of the intersection of a ray in $\Delta'_M \setminus \Delta_M$ with the boundary of $\text{Newt}(f)$, and hence is a lattice point. Secondly, for a cone $C \subset \mathbb{R}^n$ containing $\mathbb{R}_{\geq 0}^I$, let $\text{Unb}_J(C_J) := \{e_i \in \mathbb{R}^J : e_i \in C_J, \mathbb{R}_{\geq 0} e_i \cap \text{Newt}(f)_J = \emptyset\}$. Fix C' in Δ'_M such that $\mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C') \rangle$. Since $\text{Unb}(C') = \text{Unb}(\tau(C'))$ by assumption, we compute:

$$\text{Unb}_J(C'_J) = \{\text{pr}_J(e_i) : e_i \in \text{Unb}(C'), i \notin I\} = \{\text{pr}_J(e_i) : e_i \in \text{Unb}(\tau(C')), i \notin I\} = \text{Unb}_J(\tau(C'))_J.$$

Moreover, $\tau(C')_J$ is the smallest cone in Δ_J containing C'_J . We conclude that the hypotheses of Proposition 3.2.8 hold.

Let ψ_J be the unique piecewise linear function on \mathbb{R}^J with value 1 on all interior faces of $\partial \text{Newt}(f)_J$. Let $\sigma_J(C'_J)$ be the smallest face of $\mathbb{R}_{\geq 0}^I$ containing C'_J , and let $e_J(C'_J) = \dim \sigma_J(C'_J) - \dim C'_J$. Then applying Proposition 3.2.8 gives

$$\Psi_M(\tilde{E}(\text{Newt}(f)_J)) = \sum_{\substack{C'_J \in \Delta'_{J,M} \\ \text{Unb}_J(C'_J) = \emptyset \\ e_J(C'_J) = 0}} (-1)^{\dim C'_J + 1} \Psi_M \left(\sum_{w \in \text{Box}_{C'_J}} [-\psi_J(w)] \right).$$

We compute that $e_J(C'_J) = \dim \sigma_J(C'_J) - \dim C'_J = (\dim \sigma(C') - |I|) - (\dim C' - |I|) = e(C')$, so

$$\Psi_M(\tilde{E}(\text{Newt}(f)_J)) = \sum_{\substack{C' \in \Delta'_M \\ \langle \text{Unb}(C') \rangle = \mathbb{R}_{\geq 0}^I \\ e(C') = 0}} (-1)^{\dim C' + |I| + 1} \Psi_M \left(\sum_{w \in \text{Box}_{C'_J}} [-\psi_J(w)] \right).$$

Consider $C' \in \Delta'_M$ such that $\mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C') \rangle$. We define a bijection $\phi: \text{Box}_{C'} \rightarrow \text{Box}_{C'_J}$ as follows. Write $\text{Gen}(C') = \{w_1, \dots, w_r\} \cup \{e_i : i \in I\}$. If $w = \sum_{i=1}^r \lambda_i w_i + \sum_{i \in I} \mu_i e_i \in \text{Box}_{C'}$, then $\phi(w) = \sum_{i=1}^r \lambda_i \text{pr}_J(w_i)$. Observe that $\psi(w) = \psi_J(\phi(w))$. We deduce that

$$(12) \quad \sum_{w \in \text{Box}_{C'}} [-\psi(w)] = \sum_{w \in \text{Box}_{(C')_J}} [-\psi_J(w)].$$

Substituting (12) into the above equation gives

$$\Psi_M(\tilde{E}(\text{Newt}(f)_J)) = \sum_{\substack{C' \in \Delta'_M \\ \langle \text{Unb}(C') \rangle = \mathbb{R}_{\geq 0}^I \\ e(C') = 0}} (-1)^{\dim C' + |I| + 1} \Psi_M \left(\sum_{w \in \text{Box}_{C'}} [-\psi(w)] \right).$$

When $\Delta_{J,M}$ is empty, we have seen that $\Psi_M(\tilde{E}(\text{Newt}(f)_J)) = 0$, so the above formula holds then as well.

Let $C' \in \Delta'$ and assume that $\Psi_M(-\psi(w)) \neq 0$ for some $w \in \text{Box}_{C'}$. By Lemma 3.2.7, every cone in Δ containing C' lies in Δ_M , and so every cone in Δ' containing C' lies in Δ'_M .

Putting this all together, we compute

$$\begin{aligned}
\Psi_M \left(\sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \tilde{E}(\mathcal{F}_{x_I}) \right) &= \sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \Psi_M(\tilde{E}(\text{Newt}(f)_J)) \\
&= \sum_{\mathbb{A}^I \subset X_f} (-1)^{n-1-|I|} \sum_{\substack{C' \in \Delta'_M \\ \langle \text{Unb}(C') \rangle = \mathbb{R}_{\geq 0}^I \\ e(C')=0}} (-1)^{\dim C' + |I| + 1} \Psi_M \left(\sum_{w \in \text{Box}_{C'}} [-\psi(w)] \right) \\
&= \sum_{\substack{C' \in \Delta'_M \\ e(C')=0}} (-1)^{\text{codim } C'} \Psi_M \left(\sum_{w \in \text{Box}_{C'}} [-\psi(w)] \right) \\
&= \sum_{C \in \Delta'_M} \Psi_M \left(\sum_{w \in \text{Box}_C^\circ} [-\psi(w)] \right) \sum_{\substack{C \subset C' \in \Delta'_M \\ e(C')=0}} (-1)^{\text{codim } C'} \\
&= \Psi_M \left(\sum_{C \in \Delta'} \sum_{w \in \text{Box}_C^\circ} [-\psi(w)] \sum_{\substack{C \subset C' \in \Delta' \\ e(C')=0}} (-1)^{\text{codim } C'} \right) \\
&= \Psi_M \left(\sum_{C \in \Delta'} \sum_{w \in \text{Box}_C^\circ} [-\psi(w)] \ell(\Delta, C; 1) \right).
\end{aligned}$$

Here the final equality follows from (8). \square

3.3. An existence result for nearby eigenvalues of monodromy. We use Theorem 1.4.5, a vanishing result for the local h -polynomial, and Theorem 3.2.1 to study nearby eigenvalues of monodromy.

Let Δ' be a simplicial fan refining Δ . Assume that every ray of $\Delta' \setminus \Delta$ intersects the boundary of $\text{Newt}(f)$ at a lattice point. Let C' be a cone in Δ' with $\text{Gen}(C') = \{w_1, \dots, w_r\}$. We consider the function

$$\text{box}_{C'} : \text{span}(C') \cap \mathbb{Z}^n \rightarrow \text{Box}_{C'}, \text{ defined by } \text{box}_{C'} \left(\sum_{i=1}^r \lambda_i w_i \right) = \sum_{i=1}^r \{\lambda_i\} w_i,$$

where $\lambda_i \in \mathbb{Q}$ and $\{\lambda_i\}$ denotes the fractional part of λ_i .

Throughout this section, let G be a lattice simplex contained in $\partial \text{Newt}(f)$ such that $\mathbf{1} \in \text{span}(G)$ and $C_G \in \Delta'$. Let ψ_G be the unique linear function on $\text{span}(G)$ with value 1 on G . Equivalently, ψ_G is determined by the condition that $\psi_G|_{C_G} = \psi|_{C_G}$. Let $\alpha = -\psi_G(\mathbf{1})$. Then α is the candidate pole associated to any proper face F of $\text{Newt}(f)$ containing G .

Definition 3.3.1. *The essential face of G is the unique face $E \subset G$ such that $\text{box}_{C_G}(\mathbf{1})$ is in $\text{Box}_{C_E}^\circ$.*

Equivalently, one may verify that if $\{w_1, \dots, w_r\}$ are the vertices of G and we write $\mathbf{1} = \sum_{i=1}^r \lambda_i w_i$ for some $\lambda_i \in \mathbb{Q}$, then E is the unique face of G with $\text{Gen}(C_E) = \{w_i : \lambda_i \notin \mathbb{Z}\}$. Note that, for any lattice point $w \in \text{span}(C_G) \cap \mathbb{Z}_{\geq 0}^n$, $[\psi_G(w)] = [\psi(\text{box}_{C_G}(w))]$ in \mathbb{Q}/\mathbb{Z} . We deduce that

$$(13) \quad [\alpha] = [-\psi(\text{box}_{C_G}(\mathbf{1}))] \text{ in } \mathbb{Q}/\mathbb{Z}.$$

We deduce the following corollary of Theorem 3.2.1. When $\text{Newt}(f)$ is simplicial and we set $\Delta' = \Delta$, this is equivalent to Corollary 1.4.4.

Corollary 3.3.2. *Let $\alpha \in \mathbb{Q}$, and let $M = D(\alpha)$. Let Δ' be a simplicial fan refining Δ . Assume that every ray of $\Delta' \setminus \Delta$ intersects the boundary of $\text{Newt}(f)$ at a lattice point, and $\text{Unb}(C') = \text{Unb}(\tau(C'))$ for all C' in Δ'_M . Let G be a lattice simplex contained in $\partial \text{Newt}(f)$ such that $\mathbf{1} \in \text{span}(G)$ and $C_G \in \Delta'$, and let E be the essential face of G . Assume that $\alpha = -\psi_G(\mathbf{1})$. If $\ell(\Delta', C_E; t)$ is nonzero, then the coefficient of $[\alpha]$ in $\tilde{E}(\mathcal{F}_{x_I})$ is nonzero at a general point x_I of some coordinate subspace $\mathbb{A}^I \subset X_f$.*

Proof. By definition, $\Psi_{D(\alpha)}([\alpha]) = [\alpha]$. By (13), $\ell(\Delta', C_E; 1)[\alpha]$ is a term in the right-hand side of (9). The result now follows from the nonnegativity of the local h -polynomial. \square

Extending Definition 1.4.3, we may define the notion of G being UB_1 exactly as for (compact) faces of $\text{Newt}(f)$. Explicitly, a vertex A of G is an *apex* with *base direction* e_ℓ^* if $\langle e_\ell^*, A \rangle > 0$, and $\langle e_\ell^*, V \rangle = 0$ for all $V \in \text{Gen}(C_G)$ with $V \neq A$, i.e., for all vertices of G not equal to A . Then G is UB_1 if there exists an apex A in G with a unique choice of base direction e_ℓ^* , and $\langle e_\ell^*, A \rangle = 1$.

The following definition is a special case of Definition 4.1.1. We say that $C_G \setminus C_E$ in $\text{lk}_{\Delta'}(C_E)$ is a *U-pyramid* if there exists an apex A in G with a unique choice of base direction e_ℓ^* , and $A \notin \text{Gen}(C_E)$.

Lemma 3.3.3. *With the notation above, G is UB_1 if and only if $C_G \setminus C_E$ in $\text{lk}_{\Delta'}(C_E)$ is a U-pyramid.*

Proof. Let $\text{Gen}(C_G) = \{w_1, \dots, w_r\}$, and uniquely write $\mathbf{1} = \sum_{i=1}^r \lambda_i w_i$ for some $\lambda_i \in \mathbb{Q}$. Let w_i be an apex with a base direction e_ℓ^* . Let $h = \langle e_\ell^*, w_i \rangle \in \mathbb{Z}_{>0}$. Then $\lambda_i = 1/h \in \mathbb{Z}_{>0}$. The result then follows since

$$w_i \notin C_E \iff \lambda_i \in \mathbb{Z} \iff h = 1. \quad \square$$

Let \mathcal{A}_G be the set of apices of G which are not in E . For a face G' of G , let $\sigma(G')$ be the smallest face of $\mathbb{R}_{\geq 0}^n$ containing G' . Let $B_G = \{\ell \in [n] : \text{there exists } A \in \mathcal{A}_G \text{ with base direction } e_\ell^*\}$. Below, we identify faces of $\mathbb{R}_{\geq 0}^n$ with their corresponding subsets of $[n]$ and identify simplices with their set of vertices.

The following definition is a special case of Definition 4.1.2. Below, by associating faces of G with their corresponding cones in Δ' , we may view faces of $G \setminus E$ as faces in $\text{lk}_{\Delta'}(C_E)$. A *full partition* of $G \setminus E$ is a decomposition

$$G \setminus E = G_1 \sqcup G_2 \sqcup \mathcal{A}_G$$

such that

- (1) $\sigma(G_1 \sqcup \mathcal{A}_G \sqcup E) = [n]$,
- (2) $\sigma(G_2 \sqcup E) = [n] \setminus B_G$.

Lemma 3.3.4. *With the notation above, $G \setminus E$ admits a full partition.*

Proof. Let $\text{Gen}(C_G) = \{w_1, \dots, w_r\}$, and uniquely write $\mathbf{1} = \sum_{i=1}^r \lambda_i w_i$ for some $\lambda_i \in \mathbb{Q}$. We have

$$(14) \quad G \setminus E = G_1 \sqcup G_2 \sqcup \mathcal{A}_G,$$

where $G_1 = \{w_i : w_i \notin \mathcal{A}_G, \lambda_i \in \mathbb{Z}_{>0}\}$, and $G_2 = \{w_i : \lambda_i \in \mathbb{Z}_{\leq 0}\}$. Note that $w_i \in \mathcal{A}_G$ implies that $\lambda_i = 1$. We claim that (14) is a full partition.

For each w_i in $\text{Gen}(C_G)$, write $(w_i)_\ell \in \mathbb{Z}_{\geq 0}$ for the ℓ th coordinate of w_i . For each coordinate $\ell \in [n]$,

$$(15) \quad 1 = \sum_{i=1}^n \lambda_i (w_i)_\ell.$$

If $\ell \notin \sigma(\mathcal{A}_G \sqcup G_1 \sqcup E)$, then the right-hand side of (15) is a sum of nonpositive terms, a contradiction. We conclude that $\sigma(\mathcal{A}_G \sqcup G_1 \sqcup E) = [n]$.

It remains to show that $\sigma(G_2 \sqcup E) = [n] \setminus B_G$. It follows from the definitions that $\sigma(G_2 \sqcup E) \subset [n] \setminus B_G$. It remains to prove that $[n] \setminus \sigma(G_2 \sqcup E) \subset B_G$. Suppose that $\ell \in [n] \setminus \sigma(G_2 \sqcup E)$. Then all terms on the right-hand side of (15) are nonnegative integers, and we deduce that there is a unique index k such that $\lambda_k = (w_k)_\ell = 1$ and $\lambda_i(w_i)_\ell = 0$ for $i \neq k$. If $(w_i)_\ell \neq 0$ for some $i \neq k$, then $\lambda_i = 0$ and hence $\ell \in \sigma(G_2)$, a contradiction. We deduce that $w_i \in \mathcal{A}_G$ has base direction e_ℓ^* . That is, $\ell \in B_G$. \square

The following corollary is immediate from Theorem 1.4.5, together with Lemma 3.3.3 and Lemma 3.3.4. Here Theorem 1.4.5 is a consequence of Theorem 4.1.3, whose proof is the subject of Section 4.

Corollary 3.3.5. *With the notation above, if $\ell(\Delta', C_E; t) = 0$, then G is UB_1 .*

3.4. Existence of simplicial refinements. We now give a criterion for the existence of a simplicial refinement of Δ that satisfies the hypotheses of Corollary 3.3.2 and allows us to prove our strongest result on the existence of eigenvalues. We will obtain Theorem 1.4.6 as a consequence.

We first introduce a combinatorial condition on the unbounded faces of $\text{Newt}(f)$.

Definition 3.4.1. *Say that a cone C in Δ has good projection if for any face C' of C such that $C' \cap \text{Unb}(C) = \emptyset$, $\dim(C' + \langle \text{Unb}(C) \rangle) = \dim C' + |\text{Unb}(C)|$. Equivalently, for any face C' of C disjoint from $\text{Unb}(C)$, the images of the elements of $\text{Unb}(C)$ are linearly independent in $\mathbb{R}^n / \text{span}(C')$.*

We say that $\text{Newt}(f)$ has good projection if all cones in Δ have good projection. Let $M \in \mathbb{Z}_{>0}$. Then $\text{Newt}(f)$ has M -good projection if every maximal cone C in Δ such that M divides ρ_C has good projection.

Observe that Δ has M -good projection if and only if all cones in Δ_M have good projection. Clearly, $\text{Newt}(f)$ has good projection if and only if $\text{Newt}(f)$ has M -good projection for $M = 1$ if and only if $\text{Newt}(f)$ has M -good projection for all $M \in \mathbb{Z}_{>0}$. If all cones $C \in \Delta$ with $\text{Unb}(C) \neq \emptyset$ are simplicial, then $\text{Newt}(f)$ has good projection. For example, when $\text{Newt}(f)$ is simplicial or convenient, then $\text{Newt}(f)$ has good projection. Also, if $|\text{Unb}(C)| \leq 1$ for all maximal cones C in Δ , then $\text{Newt}(f)$ has good projection.

Lemma 3.4.2. *Let $\alpha \in \mathbb{Q}$. If $\text{Newt}(f)$ has $D(\alpha)$ -good projection, then C_F has good projection for all $F \in \text{Contrib}(\alpha)$.*

Proof. By definition, $\alpha = i/\rho_{C_F}$ for some $i \in \mathbb{Z}$, and hence $D(\alpha)$ divides ρ_{C_F} . Since $\text{Newt}(f)$ has $D(\alpha)$ -good projection, it follows that C_F has good projection. \square

If $\text{Newt}(f)$ has $D(\alpha)$ -good projection, then we will be able to apply Theorem 3.2.1 to deduce that $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy if a certain local h -polynomial does not vanish. The condition that $\text{Newt}(f)$ has $D(\alpha)$ -good projection is inspired in part by a stricter condition in Saito [Sai19, Definition 3.12], that itself follows ideas from [TT16]. In the language of our paper, they consider the condition on $\text{Newt}(f)$ that every maximal cone C in Δ such that $D(\alpha)$ divides ρ_C satisfies $\text{Unb}(C) = \emptyset$.

We now formulate the condition on $\text{Newt}(f)$ that will allow us to apply Corollary 3.3.5. Let F be a face of $\text{Newt}(f)$ such that $C_F \in \Delta$. Let \bar{F} denote the image of F under the projection $\mathbb{R}^n \rightarrow \mathbb{R}^n / \langle \text{Unb}(C_F) \rangle$.

Definition 3.4.3. *Let F be a face of $\text{Newt}(f)$ such that $C_F \in \Delta$. Assume that C_F has good projection. Then F is pseudo- UB_1 if every $(\dim \bar{F})$ -dimensional lattice simplex inscribed in \bar{F} is UB_1 .*

If $\text{Newt}(f)$ is simplicial, then all pseudo- UB_1 faces are UB_1 . The B_2 -facets of [ELT22, Definition 3.9] are examples of pseudo- UB_1 faces which are not UB_1 .

Remark 3.4.4. Suppose that C in Δ has good projection. Then $\Sigma = \{C_1 + C_2 : C_1 \subset C, \text{Unb}(C_1) = \emptyset, C_2 \subset \langle \text{Unb}(C) \rangle\}$ is a fan refining C . Consider $I \subset [n]$ such that $\mathbb{R}_{\geq 0}^I = \langle \text{Unb}(C) \rangle$. Let $J = [n] \setminus I$ and consider the projection $\text{pr}_J : \mathbb{R}^n \rightarrow \mathbb{R}^J$. Then

$$\text{Star}_\Sigma(\langle \text{Unb}(C) \rangle) := \{\text{pr}_J(C') : \langle \text{Unb}(C) \rangle \subset C' \in \Sigma\} = \{\text{pr}_J(C_1) : C_1 \subset C, \text{Unb}(C_1) = \emptyset\}$$

is a refinement of $\text{pr}_J(C)$ [CLS11, Exercise 3.4.8]. Let F be a face of $\text{Newt}(f)$ such that $C = C_F$ has good projection. Then it follows that F is UB_1 if and only if $\overline{F} = \text{pr}_J(F)$ is UB_1 .

Lemma 3.4.5. *Let F be a face of $\text{Newt}(f)$ such that $C_F \in \Delta$ and C_F has good projection. If F is UB_1 , then F is pseudo- UB_1 .*

Proof. Using Remark 3.4.4, we reduce to the case when F is compact. In that case, suppose F has an apex A with unique base direction e_ℓ^* , and $\langle e_\ell^*, A \rangle = 1$. Let G be a lattice simplex inscribed in F with $\dim G = \dim F$. Then A is an apex of G with base direction e_ℓ^* . Suppose e_j^* is a base direction of A in G . If $V \neq A$ is a vertex of F , then $V \in \text{span}(F \cap \{e_\ell^* = 0\}) = \text{span}(G \cap \{e_\ell^* = 0\}) = \text{span}(G \cap \{e_j^* = 0\})$. Hence e_j^* is a base direction of A in F , and $j = \ell$. We conclude that G is UB_1 , as desired. \square

We now state our strongest result on the existence of eigenvalues of monodromy.

Theorem 3.4.6. *Suppose f is nondegenerate. Let $\alpha \in \mathbb{Q}$. Assume that $\text{Newt}(f)$ has $D(\alpha)$ -good projection. Then either every face in $\text{Contrib}(\alpha)$ is pseudo- UB_1 , or $\exp(2\pi i\alpha)$ is a nearby eigenvalue of monodromy (for reduced cohomology).*

Before giving the proof, we present some applications and examples.

Proof of Theorem 1.4.6. Assume that $\text{Newt}(f)$ is simplicial. Then $\text{Newt}(f)$ has good projection, and a face F of $\text{Newt}(f)$ is pseudo- UB_1 if and only if it is UB_1 . The result now follows from Theorem 3.4.6. \square

Theorem 3.4.7. *Let f be a nondegenerate polynomial with $\text{Newt}(f) = kP$ for some $k \geq 2$ and some Newton polyhedron P . If $\text{Newt}(f)$ has good projection, then every candidate eigenvalue is a nearby eigenvalue of monodromy.*

Proof. Note that none of the vertices of kP have any coordinate equal to one, so no face of kP is pseudo- UB_1 . The result follows from Theorem 3.4.6. \square

Example 3.4.8. Suppose that $\text{Newt}(f)$ is convenient. Then $\text{Newt}(f)$ has good projection. In this case, Theorem 3.4.6 states that either every face in $\text{Contrib}(\alpha)$ is pseudo- UB_1 , or $\exp(2\pi i\alpha)$ is a nearby eigenvalue of monodromy (for reduced cohomology).

Example 3.4.9. In Example 2.2.4 and Example 3.2.4, consider the facet F with candidate pole $\alpha = -6/5$. We have $\text{Contrib}(\alpha) = \{F\}$ and F is not UB_1 , although \overline{F} is UB_1 . By Remark 3.4.4, C_F does not have good projection. In particular, F is not pseudo- UB_1 . By Lemma 3.4.2, $\text{Newt}(f)$ does not have $D(\alpha)$ -good projection, so Theorem 3.4.6 does not apply.

Example 3.4.10. Consider the set up of Example 3.2.5 with F the bounded facet with candidate pole $\alpha = -1/3$. We have $\text{Contrib}(\alpha) = \{F\}$, and F is not UB_1 . Consider the unbounded facet G of $\text{Newt}(f)$ defined by $\psi_G = \frac{1}{78}(4e_2^* + 13e_3^*) = 1$. Then $\text{Vert}(C_G) = \{w_1, w_3, w_4\}$ and $\text{Unb}(C_G) = \{e_1, e_4\}$. In particular, C_G does not have good projection. Since $D(\alpha) = 3$ divides $\rho_{C_G} = 78$, we conclude that $\text{Newt}(f)$ does not have $D(\alpha)$ -good projection, so Theorem 3.4.6 does not apply.

Proof of Theorem 3.4.6. Let F be a face in $\text{Contrib}(\alpha)$. Suppose there exists a $(\dim \overline{F})$ -dimensional lattice simplex G inscribed in \overline{F} that is not UB_1 . We need to show that $\exp(2\pi i\alpha)$ is a nearby eigenvalue of monodromy (for reduced cohomology).

First assume that F is compact. Our first goal is to construct an appropriate simplicial fan Δ' that refines Δ and contains C_G as a cone. Let $\{i_1, \dots, i_s\} = \{i \in [n] : \mathbb{A}^{\{i\}} \subset X_f\}$. Consider positive integers $0 \ll m_{i_1} \ll \dots \ll m_{i_s}$, and let $\hat{f} := f + \sum_{j=1}^s x_{i_j}^{m_{i_j}}$ with corresponding Newton polyhedron $\text{Newt}(\hat{f})$ and fan over the faces $\hat{\Delta}$. Then $\hat{\Delta}$ refines Δ and has the same rays as Δ . If $C \in \Delta$ has good projection, then every

cone in $\widehat{\Delta}|_C$ is a sum of a cone C_1 with $\text{Unb}(C_1) = \emptyset$ and a cone spanned by a subset of $\text{Unb}(C)$. Using, for example, a pulling triangulation [DLRS10, Section 4.3.2], one can construct a simplicial refinement Δ' of $\widehat{\Delta}$ such that C_G is a cone in Δ' , and the rays of Δ' are the union of the rays of Δ and the rays of C_G .

Given $C' \in \Delta'$, recall that $\tau(C')$ denotes the smallest face of Δ containing C' , and $\text{Unb}(C') \subset \text{Unb}(\tau(C'))$. If $C \in \Delta$ has good projection, then every cone in $\Delta'|_C$ is a sum of a cone $C_1 \in \Delta'|_C$ with $\text{Unb}(C_1) = \emptyset$ and a cone spanned by a subset of $\text{Unb}(C)$. In particular, if $C' \in \Delta'|_C$, then $\text{Unb}(C') = \text{Unb}(\tau(C'))$.

By Corollary 3.3.2 and Corollary 3.3.5, the coefficient of $[\alpha]$ in $\widetilde{E}(\mathcal{F}_{x_I})$ is nonzero at a general point x_I of some coordinate subspace $\mathbb{A}^I \subset X_f$.

Now consider the case when F is not necessarily compact. Consider $I \subset [n]$ such that $\mathbb{R}_{\geq 0}^I = \langle \text{Unb}(C_F) \rangle$. Let $J = [n] \setminus I$ and consider the projection $\text{pr}_J : \mathbb{R}^n \rightarrow \mathbb{R}^J$, and $\text{Newt}(f)_J := \text{pr}_J(\text{Newt}(f)) \subset \mathbb{R}^J$. Let Δ_J be the fan over the faces of $\text{Newt}(f)_J$. The maximal cones of $\text{Newt}(f)_J$ are precisely the cones of the form $\text{pr}_J(C)$, where C is a maximal cone of Δ such that $\mathbb{R}_{\geq 0}^I \subset \langle \text{Unb}(C) \rangle$. For any such cone C , $\rho_C = \rho_{\text{pr}_J(C)}$ and if C has good projection, then $\text{pr}_J(C)$ has good projection. We deduce that $\text{Newt}(f)_J$ has $D(\alpha)$ -good projection. Also, $\overline{F} = \text{pr}_J(F)$ has candidate pole α .

Let g be a nondegenerate polynomial with Newton polyhedron $\text{Newt}(f)_J$ and Milnor fiber $\widehat{\mathcal{F}}_0$ at the origin. By the compact case above, we deduce that the coefficient of $[\alpha]$ in $\widetilde{E}(\mathcal{F}_{y_{\hat{I}}})$ is nonzero at a general point $y_{\hat{I}}$ of some coordinate subspace $\mathbb{A}^{\hat{I}} \subset X_g$ with $\hat{I} \subset J$. Let $I' = I \cup \hat{I}$ and $J' = [n] \setminus I'$. Applying (11) to both $\widetilde{E}(\mathcal{F}_{x_{I'}})$ and $\widetilde{E}(\mathcal{F}_{y_{\hat{I}}})$ yields the equality $\widetilde{E}(\mathcal{F}_{x_{I'}}) = \widetilde{E}(\text{Newt}(f)_{J'}) = \widetilde{E}(\mathcal{F}_{y_{\hat{I}}})$. We conclude that the coefficient of $[\alpha]$ in $\widetilde{E}(\mathcal{F}_{x_{I'}})$ is nonzero. \square

Finally, as a corollary of the proof above, we may extend the result of Budur and van der Veer [BvdV22, Theorem 1.10] on dilates of Newton polyhedron by removing the convenient hypothesis.

Proposition 3.4.11. *Fix a Newton polyhedron P . Let f be a nondegenerate polynomial with $\text{Newt}(f) = kP$ for some $k \in \mathbb{Z}_{>0}$ chosen sufficiently large. Then every candidate eigenvalue is a nearby eigenvalue of monodromy.*

Proof. Let F be a facet of P and let α be the corresponding candidate pole. Then α/k is the corresponding candidate pole associated to the facet kF of kP . After possibly replacing F by \overline{F} , we reduce to the case when F is compact. Then the proof of [BvdV22, Theorem 1.10] applies. Explicitly, assume that F is compact and let $c_k \in \mathbb{Z}$ denote the coefficient of $[\alpha/k]$ in $\widetilde{E}(kP)$. Then Varchenko's Theorem (see (10) above) implies that

$$(16) \quad \lim_{k \rightarrow \infty} c_k/k^{n-1} = (-1)^n \sum_{F'} \text{Vol}(F'),$$

where F' varies over all facets of P such that $D(\alpha)$ divides $\rho_{F'}$. Since $F' = F$ appears in the sum on the right hand side of (16), we deduce that the left hand side of (16) is nonzero, and the result follows. \square

4. A NECESSARY CONDITION FOR THE VANISHING OF THE LOCAL h -POLYNOMIAL

4.1. Overview. In this section, we prove a necessary condition for the vanishing of the local h -polynomial of a geometric triangulation of a simplex. The section is self-contained and combinatorial in nature. As such, the notation used is independent from the rest of the paper. In Section 4.2, we recall the combinatorial commutative algebra interpretation of the local h -polynomial. In Section 4.3, we reduce our result to proving a positivity result, Proposition 4.3.5, which we then prove in Section 4.4.

Let $\sigma : \mathcal{S} \rightarrow 2^{[n]}$ be a geometric triangulation of a simplex. A face G of \mathcal{S} is *interior* if $\sigma(G) = [n]$. Let E be a face of \mathcal{S} , and let $F \in \text{lk}_{\mathcal{S}}(E)$ be a face. Then F is a *pyramid* with apex $A \in F$ if $F \sqcup E$ is interior and

$(F \sqcup E) \setminus A$ is not interior. Let

$$\mathcal{A}_F := \{A \in F : F \text{ is a pyramid with apex } A\}, \quad \text{and } V_A = V_A(F) := [n] \setminus \sigma((F \sqcup E) \setminus A)$$

for $A \in \mathcal{A}_F$. In what follows, we identify simplices with their sets of vertices.

Definition 4.1.1. *We say that F is a U -pyramid if $|V_A| = 1$ for some $A \in \mathcal{A}_F$.*

Definition 4.1.2. *A full partition of F is a decomposition*

$$F = F_1 \sqcup F_2 \sqcup \mathcal{A}_F$$

such that $F_1 \sqcup \mathcal{A}_F \sqcup E$ is interior and $\sigma(F_2 \sqcup E) = [n] \setminus \bigcup_{A \in \mathcal{A}_F} V_A$.

Recall that $\ell(\mathcal{S}, E; t)$ denotes the corresponding local h -polynomial. See Definition 3.1.1. Our goal is to prove the following theorem.

Theorem 4.1.3. *Let $\sigma : \mathcal{S} \rightarrow 2^{[n]}$ be a geometric triangulation of a simplex, and fix a face $E \in \mathcal{S}$. Let $F \in \text{lk}_{\mathcal{S}}(E)$ be a face that admits a full partition $F = F_1 \sqcup F_2 \sqcup \mathcal{A}_F$. If the coefficient of $t^{|F_1|+|\mathcal{A}_F|}$ in $\ell(\mathcal{S}, E; t)$ is zero, then F is a U -pyramid.*

Our strategy is as follows: assume that F admits a full partition and is not a U -pyramid. We argue that the nonvanishing of the local h -polynomial is implied by the nonvanishing of a specific element in the highest degree cohomology of an associated complete toric variety, expressed as a non-squarefree monomial in the torus-invariant divisors. We then compute an explicit nonzero formula for this element, completing the proof.

4.2. The commutative algebra of local h -polynomials. Let Δ be a rational simplicial fan in \mathbb{R}^n with support $\mathbb{R}_{\geq 0}^n$. For each ray of Δ , choose a rational, nonzero point v . Consider the unique piecewise \mathbb{Q} -linear function $\psi : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}$ defined by $\psi(v) = 1$ for all such v , and let $\mathcal{S} = \{x \in \mathbb{R}_{\geq 0}^n : \psi(x) = 1\}$. Then \mathcal{S} is a simplicial complex with vertices $\{v\}$, and \mathcal{S} induces a geometric triangulation $\sigma : \mathcal{S} \rightarrow 2^{[n]}$ of a simplex by projecting onto a transverse hyperplane. The combinatorial type of this triangulation is independent of both the choice of $\{v\}$ and the choice of transverse hyperplane. Explicitly, if F is a face of \mathcal{S} , then $\mathbb{R}^{\sigma(F)}$ is the smallest coordinate hyperplane containing F . Conversely, given a geometric triangulation of a simplex, we may deform the vertices without changing the combinatorial type to assume that the triangulation is rational, and then the triangulation is realized by some such \mathcal{S} .

If F is a face of \mathcal{S} , let C_F denote the cone over F . For example, when $F = \emptyset$, then $C_F = \{0\}$. Then $\Delta = \{C_F : F \in \mathcal{S}\}$. Fix a face E of \mathcal{S} . Then the collection of cones Δ_E given by the images of $\{C_F : F \in \text{lk}_{\mathcal{S}}(E)\}$ in $\mathbb{R}^n / \text{span}(E)$, forms a fan. For example, $\Delta_{\emptyset} = \Delta$, and Δ_E is complete if and only if E is an interior face of \mathcal{S} . Consider the standard lattice $\mathbb{Z}^n \subset \mathbb{R}^n$, and let X_E denote the toric variety associated to Δ_E . The torus orbits in X_E are in inclusion reversing bijection with the faces in $\text{lk}_{\mathcal{S}}(E)$. If $E \subset E'$, then $X_{E'}$ is the closure in X_E of the torus orbit corresponding to the face $E' \setminus E$ of $\text{lk}_{\mathcal{S}}(E)$.

Given a finite simplicial complex \mathcal{T} , let $\mathbb{Q}[\mathcal{T}]$ denote the *face ring* of \mathcal{T} over \mathbb{Q} , i.e., the quotient of the polynomial ring over \mathbb{Q} with variables corresponding to the vertices of \mathcal{T} by the ideal generated by monomials corresponding to non-faces. For a face $F \in \mathcal{T}$, let $x^F \in \mathbb{Q}[\mathcal{T}]$ denote the product of the variables corresponding to the vertices of F . Note that $\mathbb{Q}[\mathcal{T}]$ is graded by degree. We write $|G|$ for the number of vertices in a face G . In particular, x^F is a squarefree monomial of degree $|F|$.

A *linear system of parameters* (l.s.o.p.) for a finitely generated graded \mathbb{Q} -algebra R of Krull dimension d is a sequence of elements $\theta_1, \dots, \theta_d$ in R_1 such that $R/(\theta_1, \dots, \theta_d)$ is a finite-dimensional \mathbb{Q} -vector space. If \mathcal{T} has dimension $d - 1$, then $\mathbb{Q}[\mathcal{T}]$ has Krull dimension d .

Let $c = n - |E|$. Note that c is the Krull dimension of $\mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]$. The *support* of an element $\theta = \sum a_v x^v \in \mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]_1$ is $\text{supp}(\theta) := \{v : a_v \neq 0\}$. A linear system of parameters $\theta_1, \dots, \theta_c$ for $\mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]$ is *special*, as

defined in [Sta92, Ath12a] if, for each vertex $v \in [n] \setminus \sigma(E)$, there is an element θ_v of the l.s.o.p. such that $\text{supp}(\theta_v)$ consists of vertices w in $\text{lk}_{\mathcal{S}}(E)$ such that $v \in \sigma(w)$, and such that $\theta_v \neq \theta_{v'}$ for $v \neq v'$.

Proposition 4.2.1. [Ath12a, Ath12b], *see also* [LPS22, Proof of Theorem 1.2] *Let I be the ideal in $\mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]$ generated by $\{x^F : F \sqcup E \text{ is interior}\}$. Let $L(\mathcal{S}, E)$ be the image of I in $\mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]/(\theta_1, \dots, \theta_c)$, where $\theta_1, \dots, \theta_c$ is a special l.s.o.p. Then the Hilbert series of $L(\mathcal{S}, E)$ is $\ell(\mathcal{S}, E; t)$.*

We call $L(\mathcal{S}, E)$ the *local face module*. Note that the local face module depends on the choice of a special l.s.o.p. In this paper, we will consider a particular special l.s.o.p. that is defined in terms of Δ .

Below we view elements of $(\mathbb{Q}^n / \text{span}(E))^* \hookrightarrow (\mathbb{Q}^n)^*$ as \mathbb{Q} -linear functions vanishing on $\text{span}(E)$. For $u \in (\mathbb{Q}^n / \text{span}(E))^*$, let $\theta_u = \sum_{v \in \text{lk}_{\mathcal{S}}(E)} \langle u, v \rangle x^v \in \mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]$. Consider the ideal $J_E = (\theta_u : u \in (\mathbb{Q}^n / \text{span}(E))^*)$ in $\mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]$. Note that J_E is generated by a special l.s.o.p., obtained by extending $\{e_i^* : i \in [n] \setminus \sigma(E)\}$ to a basis for $(\mathbb{Q}^n / \text{span}(E))^*$.

Let $H^*(E) = \mathbb{Q}[\text{lk}_{\mathcal{S}}(E)]/J_E$. Then $H^*(E)$ is isomorphic to the rational cohomology ring $H^*(X_E, \mathbb{Q})$ of X_E . The ideal in $H^*(E)$ generated by $\{x^F : F \in \text{lk}_{\mathcal{S}}(E), F \sqcup E \text{ interior}\}$ is $L(\mathcal{S}, E)$. We will show the nonvanishing of $\ell(\mathcal{S}, E; t)$ by showing that a certain element of $L(\mathcal{S}, E)$ is nonzero. To achieve this, we require the three constructions.

First, if $E \subset E'$ is an inclusion of faces in \mathcal{S} , then there is a graded \mathbb{Q} -algebra homomorphism $\iota^* = \iota_{E, E'}^* : H^*(E) \rightarrow H^*(E')$ corresponding to the pullback map on cohomology. The *closed star* $\text{Star}(E' \setminus E)$ of $E' \setminus E$ is the subcomplex of $\text{lk}_{\mathcal{S}}(E)$ that consists of faces H such that $H \cup (E' \setminus E)$ is a face of $\text{lk}_{\mathcal{S}}(E)$. Then ι^* may be characterized as follows: let $v \in \text{lk}_{\mathcal{S}}(E)$. Then

- (1) $\iota^*(x^v) = 0$ if $v \notin \text{Star}(E' \setminus E)$,
- (2) $\iota^*(x^v) = x^v$ if $v \in \text{lk}_{\mathcal{S}}(E')$.

Note that $\text{Star}(E' \setminus E)$ is the join of $\text{lk}_{\mathcal{S}}(E')$ with $E' \setminus E$. If $v \in E' \setminus E$, then there exists a linear form u_v in $(\mathbb{Q}^n / \text{span}(E))^*$ that takes value 1 on v and vanishes on all other $v' \in E'$, and the above properties imply that $\iota^*(x^v) = -\sum_{v' \in \text{lk}_{\mathcal{S}}(E')} \langle u_v, v' \rangle x^{v'}$.

Second, let $j_* = j_{E', E, *}: H^*(E') \rightarrow H^*(E)$ be defined by $j_*(x^G) = x^G x^{E' \setminus E}$ for all $G \in \text{lk}_{\mathcal{S}}(E')$, corresponding to the Gysin pushforward map on cohomology. It then follows from the characterization of ι^* via (1) and (2), that $j_* \circ \iota^* : H^*(E) \rightarrow H^*(E)$ is multiplication by $x^{E' \setminus E}$.

Finally, assume that E is interior. Then X_E is a complete toric variety, and the degree map on top cohomology gives rise to a \mathbb{Q} -vector space isomorphism $\text{deg}_E : H^c(E) \rightarrow \mathbb{Q}$. We have the following explicit description. For a facet H of \mathcal{S} , let $m(H)$ be the absolute value of the determinant of the matrix whose columns are the coordinates of the vertices of H in \mathbb{R}^n . If G is a facet of $\text{lk}_{\mathcal{S}}(E)$, then $\text{deg}_E(x^G) = 1/m(G \sqcup E)$. If $E \subset E'$, then it follows from the description of the degree map that $\text{deg}_{E'} = \text{deg}_E \circ j_*$. Let $c' = n - |E'|$. Then for any element $z \in H^{c'}(E)$, we compute

$$(17) \quad \text{deg}_E(z x^{E' \setminus E}) = \text{deg}_E(j_* \circ \iota^*(z)) = \text{deg}_{E'}(\iota^*(z)).$$

Remark 4.2.2. If we replace $\{e_i^*\}$ by $\{\lambda_i e_i^*\}$ for some $\lambda_i \in \mathbb{Q}_{>0}$, then the definitions of $H^*(E)$, ι^* and j_* are unaffected, while deg_E is composed with multiplication by $\prod_{i=1}^n \lambda_i$.

4.3. Reduction steps. In this subsection, we reduce Theorem 4.1.3 to an explicit calculation in the top cohomology group of a complete toric variety. We continue with the notation above.

Lemma 4.3.1. *Let $F \in \text{lk}_{\mathcal{S}}(E)$ be a face with a full partition $F = F_1 \sqcup F_2 \sqcup \mathcal{A}_F$. Let $G \in \text{lk}_{\mathcal{S}}(E \sqcup F)$. Consider $F' = F \sqcup G$ in $\text{lk}_{\mathcal{S}}(E)$. Then*

$$F' = F'_1 \sqcup F'_2 \sqcup \mathcal{A}_{F'}$$

is a full partition, where $\widehat{\mathcal{A}}_F = \{A \in \mathcal{A}_F : V_A \subset \sigma(G)\}$, $F'_1 = F_1 \sqcup \widehat{\mathcal{A}}_F$, $F'_2 = F_2 \sqcup G$ and $\mathcal{A}_{F'} = \mathcal{A}_F \setminus \widehat{\mathcal{A}}_F$.

Proof. As $F \sqcup E$ is interior and $F \subset F'$, if F' is a pyramid with apex A , then $A \in \mathcal{A}_F$. For any $A \in \mathcal{A}_F$, F' is a pyramid with apex A if and only if $A \notin \widehat{\mathcal{A}}_F$, so $\mathcal{A}_{F'} = \mathcal{A}_F \setminus \widehat{\mathcal{A}}_F$. We have that $(F_1 \sqcup \widehat{\mathcal{A}}_F) \sqcup (\mathcal{A}_F \setminus \widehat{\mathcal{A}}_F) \sqcup E = F_1 \sqcup \mathcal{A}_F \sqcup E$ is interior by the full partition condition on F . The definition of $\widehat{\mathcal{A}}_F$ and the full partition condition on F imply that $\sigma(F_2 \sqcup G \sqcup E) = [n] \setminus \cup_{A \in \mathcal{A}_{F'}} V_A$. \square

Remark 4.3.2. Let F be a face of $\text{lk}_{\mathcal{S}}(E)$ such that $F \sqcup E$ is interior. Assume that $F \in \text{lk}_E(\mathcal{S})$ is not a U -pyramid. Then $\text{codim}(F \sqcup E) \geq |\mathcal{A}_F|$, and equality implies that $|V_A| = 2$ for all $A \in \mathcal{A}_F$.

Definition 4.3.3. Let F be a face of $\text{lk}_{\mathcal{S}}(E)$. We say that F is a maximal non- U -pyramid if $F \sqcup E$ is interior, F is not a U -pyramid, and for any $F \subset F' \in \text{lk}_{\mathcal{S}}(E)$ with $F \neq F'$, F' is a U -pyramid.

Proposition 4.3.4. Let F be a face of $\text{lk}_{\mathcal{S}}(E)$ that is a maximal non- U -pyramid. Then $\text{codim}(F \sqcup E) = |\mathcal{A}_F|$.

Proof. By Remark 4.3.2, $\text{codim}(F \sqcup E) \geq |\mathcal{A}_F|$. Assume that $\text{codim}(F \sqcup E) > |\mathcal{A}_F|$. We need to show that there exists $F \subset F'$ with $F \neq F'$, and F' is not a U -pyramid.

We first describe a basis for $(\mathbb{Q}^n / \text{span}(F \sqcup E))^*$. For each $A \in \mathcal{A}_F$, choose an ordering on $V_A = \{i_{A,1}, \dots, i_{A,|V_A|}\}$. For each $1 \leq j \leq |V_A| - 1$, let $u_{A,j} = \langle e_{i_{A,j+1}}^*, A \rangle e_{i_{A,j}}^* - \langle e_{i_{A,j}}^*, A \rangle e_{i_{A,j+1}}^*$. Then we can find $\{t_k\}$ such that $\{u_{A,j}\} \cup \{t_k\}$ is a basis for $(\mathbb{Q}^n / \text{span}(F \sqcup E))^*$, and $\langle t_k, e_{i_{A,j}} \rangle = 0$ for all choices of A, j, k . We consider the corresponding isomorphism $\phi: \mathbb{R}^n / \text{span}(F \sqcup E) \rightarrow (\mathbb{R}^n / \text{span}(F \sqcup E))^*$, given by $\phi(V) = \sum_{A,j} \langle u_{A,j}, V \rangle u_{A,j} + \sum_k \langle t_k, V \rangle t_k$.

Consider the complete fan $\Delta_{F \sqcup E}$ given by the images of $\{C_G : G \in \text{lk}_{\mathcal{S}}(F \sqcup E)\}$ in $\mathbb{R}^n / \text{span}(F \sqcup E)$. Given a ray R in $(\mathbb{R}^n / \text{span}(F \sqcup E))^*$, there is a unique element $V \in \mathbb{R}^n$ in the subcomplex $\text{lk}_{\mathcal{S}}(F \sqcup E)$ of \mathcal{S} such that the image of V in $\mathbb{R}^n / \text{span}(F \sqcup E)$ maps via ϕ to an element of R . Let G be the unique nonempty face of $\text{lk}_{\mathcal{S}}(F \sqcup E)$ containing V in its relative interior. Then for each $1 \leq i \leq n$, $\langle e_i^*, V \rangle > 0$ if and only if there exists a vertex W of G such that $\langle e_i^*, W \rangle > 0$. For any choice of R , we may consider the face $F' = F \sqcup G$ of $\text{lk}_{\mathcal{S}}(E)$. It remains to choose R such that F' is not a U -pyramid. There are two cases.

First, suppose that $|V_A| > 2$ for some A . Consider the ray $R := \{\lambda u_{A,|V_A|-1} : \lambda \leq 0\}$. For each $A' \neq A$, $\langle u_{A',j}, V \rangle = 0$ for $1 \leq j \leq |V_{A'}| - 1$, and hence $\{\langle e_{i_{A',j}}^*, V \rangle\}_{1 \leq j \leq |V_{A'}|-1}$ are either all nonzero or all zero. In the latter case, $A' \notin \mathcal{A}_{F'}$. In the former case, $A' \in \mathcal{A}_{F'}$, and $|V_{A'}(F')| = |V_{A'}(F)| \geq 2$. Also, $\langle u_{A,|V_A|-1}, V \rangle < 0$ implies that $\langle e_{i_{A,|V_A|}}^*, V \rangle > 0$, and $\langle u_{A,j}, V \rangle = 0$, for $1 \leq j \leq |V_A| - 2$, implies that $\{\langle e_{i_{A,j}}^*, V \rangle\}_{1 \leq j \leq |V_A|-2}$ are either all nonzero or all zero. In the latter case, $A \notin \mathcal{A}_{F'}$. In the former case, $A \in \mathcal{A}_{F'}$, and $|V_A(F')| = |V_A(F)| - 1 \geq 2$. We conclude that F' is not a U -pyramid.

Second, suppose that $|V_A| = 2$ for all A . Then we may consider the ray $R := \{\lambda t_1 : \lambda > 0\}$. For each A , $\langle u_{A,j}, V \rangle = 0$ for $1 \leq j \leq |V_A| - 1$, and hence $\{\langle e_{i_{A,j}}^*, V \rangle\}_{1 \leq j \leq |V_A|-1}$ are either all nonzero or all zero. In the latter case, $A \notin \mathcal{A}_{F'}$. In the former case, $A \in \mathcal{A}_{F'}$, and $|V_A(F')| = |V_A(F)| \geq 2$. We again conclude that F' is not a U -pyramid. \square

The key computation used to prove Theorem 4.1.3 is the following proposition. We explain how it implies Theorem 4.1.3, and then we prove it in the following section.

Proposition 4.3.5. Let F be a face of $\text{lk}_{\mathcal{S}}(E)$ that is a maximal non- U -pyramid. Then

$$(-1)^{|\mathcal{A}_F|} \deg_{F \sqcup E}(\iota_{E, F \sqcup E}^*(x^{A_F})) > 0.$$

Proof of Theorem 4.1.3. Suppose $F \in \text{lk}_{\mathcal{S}}(E)$ admits a full partition $F = F_1 \sqcup F_2 \sqcup \mathcal{A}_F$ and is not a U -pyramid. There exists $G \in \text{lk}_{\mathcal{S}}(F \sqcup E)$ such that $F' = F \sqcup G$ is a maximal non- U -pyramid. By Lemma 4.3.1, F' admits a full partition $F'_1 \sqcup F'_2 \sqcup \mathcal{A}_{F'}$ with $F'_1 = F_1 \sqcup \widehat{\mathcal{A}}_F$, $F'_2 = F_2 \sqcup G$, and $\mathcal{A}_{F'} = \mathcal{A}_F \setminus \widehat{\mathcal{A}}_F$ for some subface $\widehat{\mathcal{A}}_F$ of \mathcal{A}_F . By the definition of a full partition, $F_1 \sqcup \mathcal{A}_F \sqcup E$ is interior, so $x^{F_1 \sqcup \mathcal{A}_F} = x^{F'_1 \sqcup \mathcal{A}_{F'}} \in H^*(E)$ is contained in $L(\mathcal{S}, E)$. If we can show that $x^{F'_1 \sqcup \mathcal{A}_{F'}} \in H^*(E)$ is nonzero, then $L(\mathcal{S}, E)$ is nonzero in degree $|F_1| + |\mathcal{A}_F|$, as desired.

We apply (17) with E replaced by $F'_2 \sqcup \mathcal{A}'_F \sqcup E$, E' replaced by $F' \sqcup E$, and $z = \iota_{E, F'_2 \sqcup \mathcal{A}'_F \sqcup E}^*(x^{\mathcal{A}_{F'}})$, to obtain

$$\deg_{F'_2 \sqcup \mathcal{A}'_F \sqcup E}(\iota_{E, F'_2 \sqcup \mathcal{A}'_F \sqcup E}^*(x^{\mathcal{A}_{F'}})x^{F'_1}) = \deg_{F'_2 \sqcup \mathcal{A}'_F \sqcup E}(\iota_{E, F'_2 \sqcup \mathcal{A}'_F \sqcup E}^*(x^{F'_1 \sqcup \mathcal{A}_{F'}})) = \deg_{F' \sqcup E}(\iota_{E, F' \sqcup E}^*(x^{\mathcal{A}_{F'}})).$$

By Proposition 4.3.5, the right-hand side of the above equation is nonzero, and hence $x^{F'_1 \sqcup \mathcal{A}_{F'}}$ is nonzero. \square

Remark 4.3.6. The statement of Proposition 4.3.5 has the following geometric interpretation. The refined self-intersection of the compact T -invariant subvariety $X_{F \sqcup E}$ of half-dimension in the toric variety $X_{(F \sqcup E) \setminus \mathcal{A}_F}$ is not zero, and its sign is determined by the number of apices $|\mathcal{A}_F|$.

4.4. Positivity for maximal non- U -pyramids. In this section, we prove Proposition 4.3.5. We first fix our setup. Let F be a face of $\text{lk}_S(E)$ that is a maximal non- U -pyramid. Let $r = \text{codim}(F \sqcup E)$. By Remark 4.3.2 and Proposition 4.3.4, we may let $\mathcal{A}_F = \{A_1, \dots, A_r\}$ and assume that $V_{A_i}(F) = \{2i-1, 2i\}$. For a vector $V \in \mathbb{R}^n$, let $V_j = \langle e_j^*, V \rangle$. By Remark 4.2.2, we may rescale the first $2r$ coordinates, and assume that $(A_i)_{2i-1} = (A_i)_{2i} = 1$. Recall that for $u \in (\mathbb{Q}^n / \text{span}(E))^*$, we have the equality in $H^*(E)$

$$(18) \quad \theta_u := \sum_{v \in \text{lk}_S(E)} \langle u, v \rangle x^v = 0.$$

We introduce the following notation: given $F \in \text{lk}_S(E)$, we write $y^F := \iota_{E, F \sqcup E}^*(x^F) \in H^*(F \sqcup E)$. Then our goal is to prove that $(-1)^r \deg_{F \sqcup E}(y^{A_1} \cdots y^{A_r}) > 0$. For G a face in $\text{lk}_S(F \sqcup E)$, we define $\text{Supp}_G := \{1 \leq j \leq 2r : V_j \neq 0 \text{ for some vertex } V \in G\}$. Let $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^r$ be defined by $\pi(x_1, \dots, x_n) = (x_1 - x_2, x_3 - x_4, \dots, x_{2r-1} - x_{2r})$ (cf. the proof of Proposition 4.3.4). Then π induces an isomorphism $\mathbb{R}^n / \text{span}(F \sqcup E) \cong \mathbb{R}^r$, mapping the complete fan $\Delta_{F \sqcup E}$ to the complete fan with cones $\{\pi(C_G) : G \in \text{lk}_S(F \sqcup E)\}$.

4.4.1. Motivation for the proof. In order to help motivate and guide the reader, we give a quick explanation of the proof when $r = 2$, and then explain some of the key ideas for the general case.

Assume that $r = 2$. To compute the degree of $y^{A_1} y^{A_2} \in H^2(F \sqcup E)$, we want to show it is equivalent to a sum $\sum_G \lambda_G y^G$, where G varies over the facets of $\text{lk}_S(F \sqcup E)$ for some $\lambda_G \in \mathbb{Q}$. Then $\deg_{F \sqcup E}(y^{A_1} y^{A_2}) = \sum_G \lambda_G \deg_{F \sqcup E}(y^G)$. This is always possible, but, in general, it will lead to a large sum with positive and negative contributions. Instead, our goal is to arrange that the sum consists of a single term.

Let $\mu = e_1^*$. Applying (18) with $u = \mu$, implies that

$$-y^{A_1} y^{A_2} = \sum_{v \in \text{lk}_S(F \sqcup E)} v_1 y^v y^{A_2}.$$

Consider $v \in \text{lk}_S(F \sqcup E)$ such that $v_1 > 0$. Suppose that $v_2 > 0$. We claim that $y^v y^{A_2} = 0$. Indeed, maximality implies that $F \sqcup \{v\}$ is a U -pyramid. Hence, without loss of generality, we may assume that $v_3 = 0$ and $v_4 > 0$. Applying (18) with $u = e_3^*$, implies that:

$$-y^v y^{A_2} = \sum_{w \in \text{lk}_S(F \sqcup v \sqcup E)} w_3 y^v y^w.$$

If $w_3 > 0$ in any term above, then $F \sqcup v \sqcup w$ is not a U -pyramid, contradicting maximality. Therefore $v_2 = 0$, so $\pi_1(v) = v_1 > 0$. We deduce that

$$(19) \quad -y^{A_1} y^{A_2} = \sum_{\substack{v \in \text{lk}_S(F \sqcup E) \\ \pi_1(v) > 0}} \pi_1(v) y^v y^{A_2}.$$




FIGURE 1. Examples of fans and their unique positive maximal cones.

For $x, y \in \mathbb{R}^n$, let $A(x, y) = \begin{bmatrix} \pi_1(x) & \pi_1(y) \\ x_3 & y_3 \end{bmatrix}$, and $A'(x, y) = \begin{bmatrix} \pi_1(x) & \pi_1(y) \\ \pi_2(x) & \pi_2(y) \end{bmatrix}$. Fix $v \in \text{lk}_{\mathcal{S}}(F \sqcup E)$, and consider the linear function $\mu = \mu_v: \mathbb{R}^n \rightarrow \mathbb{R}$, defined by $\mu(x) = \det(A(v, x))$. Applying (18) with $u = \mu$ to each term in the right-hand side of (19) implies that

$$y^{A_1} y^{A_2} = \sum_{\substack{v \sqcup w \in \text{lk}_{\mathcal{S}}(F \sqcup E) \\ \pi_1(v) > 0}} \det A(v, w) y^v y^w,$$

where the sum is over ordered pairs (v, w) of distinct vertices such that $v \sqcup w$ is a face in $\text{lk}_{\mathcal{S}}(F \sqcup E)$. Switching the roles of v and w cancels the contributions where $\pi_1(w) > 0$, leaving:

$$y^{A_1} y^{A_2} = \sum_{\substack{v \sqcup w \in \text{lk}_{\mathcal{S}}(F \sqcup E) \\ \pi_1(v) > 0, \pi_1(w) \leq 0}} \det A(v, w) y^v y^w.$$

Consider a nonzero term above. Then $\det A(v, w) > 0$. We claim that $\det A(v, w) = \det A'(v, w)$. Indeed, if $\pi_1(w) < 0$, then since $F \sqcup v \sqcup w$ is not a U -pyramid, $v_4 = w_4 = 0$ because v_3 and w_3 cannot both vanish. If $\pi_1(w) = 0$, then $\det A(v, w) = \pi_1(v)w_3$. Since $F \sqcup w$ is not a U -pyramid, $w_4 = 0$. In both cases, the claim follows. We then deduce:

$$y^{A_1} y^{A_2} = \sum_{\substack{v \sqcup w \in \text{lk}_{\mathcal{S}}(F \sqcup E) \\ \pi_1(v) > 0, \pi_1(w) \leq 0 \\ \det A'(v, w) > 0}} \det A'(v, w) y^v y^w.$$

There is a unique facet G in $\text{lk}(F \sqcup E)$ such that $G = v \sqcup w$ appears in the summation above. Explicitly, G is the unique facet such that the cone $\pi(C_G)$ contains $(0, 1) \in \mathbb{R}^2$, and the projection of $\pi(C_G)$ onto the first coordinate contains $1 \in \mathbb{R}$. Equivalently, G is the unique facet such that $\pi(C_G)$ contains $(\epsilon, 1)$ for $0 < \epsilon \ll 1$. We call a cone *positive* if it satisfies these equivalent conditions. See Figure 1. We deduce that $y^{A_1} y^{A_2} = \text{mult}(G) y^G$, where $\text{mult}(G) = \det A'(v, w) = |\det A'(v, w)| > 0$ is the multiplicity of C_G . Since $\deg_{F \sqcup E}(y^G) > 0$ by definition, this completes the proof of Proposition 4.3.5.

We next discuss the general case. Below, we will generalize and extend both the notion of positive cones, and the expansion techniques above. We aim to show that $(-1)^r \deg_{F \sqcup E}(y^{A_1} \cdots y^{A_r}) = \text{mult}(G) y^G$, where G is the unique facet in $\text{lk}_{\mathcal{S}}(F \sqcup E)$ such that $\pi(C_G)$ is a positive cone. As above, this will complete the proof of Proposition 4.3.5.

The final obstruction is not visible in the $r = 2$ case. Namely, if one follows the techniques described above when $r > 2$, error terms naturally appear that must be shown to vanish. One can apply similar techniques to expand the error terms, and secondary error terms naturally appear that also must be shown to vanish. Continuing in this way, we obtain an infinite series of error terms that we must show vanish. This leads one to set up a more general problem involving all the terms we wish to analyze. More specifically, we introduce the elements of interest in Definition 4.4.25, and our main result is to compute them in Proposition 4.4.31. As a special case, $(-1)^r \deg_{F \sqcup E}(y^{A_1} \cdots y^{A_r}) = \text{mult}(G)y^G$, as desired.

Additional notation. We introduce the following notation for use in the proof. Let S, T be (possibly empty) ordered sets that are subsets of $[r]$. Suppose we have an inclusion of sets $S \subset T$. Given an ordered set $U = \{u_{i_j}\}_{i_j \in T}$ indexed by T , we write $U|_S := \{v_{i_j}\}_{i_j \in S}$. If f is a function defined on the elements of U , we write $f(U) = \{f(u_{i_j})\}_{i_j \in T}$. Let $\mathbb{R}^S, \mathbb{R}^T$ denote real vector spaces with coordinates indexed by S, T respectively, and let $\text{pr}_{T,S}: \mathbb{R}^T \rightarrow \mathbb{R}^S$ denote the associated projection map. If $S = \emptyset$, then $\mathbb{R}^S = \{0\}$. If U is an ordered set of elements of \mathbb{R}^S , indexed by T , then let A_U to be the $|S| \times |T|$ matrix with columns given by the elements of U , ordered by T , and rows indexed by S .

4.4.2. *Positive cones.* We now study the combinatorics of certain cones which will play an important role in the sequel. Throughout this section, $S = \{i_1 < \cdots < i_s\}$ will be a (possibly empty) ordered set of $s = |S|$ positive integers, and $i_0 = 0$.

Definition 4.4.1. We say that C is positive if $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_k]}(C) \subset \mathbb{R}^{S \cap [i_k]}$ for $1 \leq k \leq s$.

For example, when $S = \emptyset$, $C = \mathbb{R}^S = \{0\}$ is positive. For $\epsilon > 0$, let $\vec{\epsilon}_S := (\epsilon^{-i_1}, \epsilon^{-i_2}, \dots, \epsilon^{-i_s}) \in \mathbb{R}^S$. If $S = \emptyset$, let $\vec{\epsilon}_S = 0 \in \mathbb{R}^S = \{0\}$.

Remark 4.4.2. Let $W \subset \mathbb{R}^S$ be a finite union of proper affine subspaces. Since $\{\epsilon_S : \epsilon > 0\} \cap W$ is finite, it follows that $\epsilon_S \cap W = \emptyset$ for $\epsilon > 0$ sufficiently small.

Lemma 4.4.3. A cone C is positive if and only if $\vec{\epsilon}_S \in C$ for all $\epsilon > 0$ sufficiently small.

Proof. Suppose $\vec{\epsilon}_S \in C$ for all $\epsilon > 0$ sufficiently small. For $1 \leq k \leq s$, we compute in $\mathbb{R}^{S \cap [i_k]}$:

$$(0, \dots, 0, 1) = \lim_{\epsilon \rightarrow 0} \epsilon^{i_k} \vec{\epsilon}_{S \cap [i_k]} = \lim_{\epsilon \rightarrow 0} \text{pr}_{S, S \cap [i_k]}(\epsilon^{i_k} \vec{\epsilon}_S) \in \text{pr}_{S, S \cap [i_k]}(C).$$

Hence, C is positive. Conversely, assume that C is positive. By hypothesis, there exists an ordered set $U = \{u_{i_1}, \dots, u_{i_s}\}$ of elements of C such that $\text{pr}_{S, S \cap [i_k]}(u_{i_k}) = (0, \dots, 0, 1)$ for $1 \leq k \leq s$. The associated matrix A_U is lower triangular with 1's on the diagonal. Since A_U^{-1} is also lower triangular with 1's on the diagonal, $\lim_{\epsilon \rightarrow 0} \epsilon^{i_k} (A_U^{-1} \vec{\epsilon}_S)_{i_k} = 1$, and hence, for ϵ sufficiently small, $A_U^{-1} \vec{\epsilon}_S$ has positive entries, and $\vec{\epsilon}_S = A_U(A_U^{-1} \vec{\epsilon}_S) \in C$. \square

In particular, positive cones are full-dimensional, and every complete polyhedral fan contains a unique positive cone. The goal for the remainder of this section is to develop an alternative inductive criterion for C to be positive, Proposition 4.4.14. We will use the following linear algebra lemma.

Lemma 4.4.4. Let $m, n \in \mathbb{Z}_{\geq 0}$ with $p: \mathbb{R}^m \rightarrow \mathbb{R}^n$ a linear map. Let $C \subset \mathbb{R}^m$ be a simplicial cone, and let \widehat{C} be the smallest face of C containing $C \cap \ker(p)$. Then $p(\widehat{C})$ is the largest linear subspace contained in $p(C)$.

Moreover, if $\text{span}(C \cap \ker(p)) = \text{span}(C) \cap \ker(p)$, then $p(C)/p(\widehat{C}) \subset \mathbb{R}^n/p(\widehat{C})$ is a simplicial cone of dimension $\text{codim}(\widehat{C}, \text{span}(C))$. The faces of $p(C)$ are $\{p(C') : \widehat{C} \subset C' \subset C\}$, and $\text{codim}(p(C'), \text{span}(p(C))) = \text{codim}(C', \text{span}(C))$ for $\widehat{C} \subset C' \subset C$.

Proof. Replacing \mathbb{R}^m with $\text{span}(C)$ and \mathbb{R}^n with $\text{span}(p(C))$, we reduce to the case when C and $p(C)$ are full-dimensional. The result holds when $m = 0$. Assume that $m > 0$.

Let u_1, \dots, u_m be generators of the rays of C . Let L be the largest linear subspace contained in $p(C)$. Suppose $x \in L$. Then $x = \sum_{i=1}^m \alpha_i p(u_i)$ for some $\alpha_i \geq 0$, and $-x = \sum_{i=1}^m \beta_i p(u_i)$ for some $\beta_i \geq 0$. Hence $\sum_{i=1}^m (\alpha_i + \beta_i) u_i \in C \cap \ker(p)$. If $\alpha_i > 0$, then $u_i \in \widehat{C}$, and hence $x \in p(\widehat{C})$.

Conversely, suppose $u_j \in \widehat{C}$. Then there exists $y = \sum_{i=1}^m \alpha_i u_i \in C \cap \ker(p)$ for some $\alpha_i \geq 0$ such that $\alpha_j > 0$. Then $-p(u_j) = (1/\alpha_j) \sum_{i \neq j} \alpha_i p(u_i) \in p(C)$, and hence $p(u_j) \in L$. We conclude that $p(\widehat{C}) \subset L$.

Assume that $\text{span}(C \cap \ker(p)) = \ker(p)$. Then the restriction $p|_{\text{span}(\widehat{C})} : \text{span}(\widehat{C}) \rightarrow \text{span}(p(\widehat{C}))$ is surjective with kernel $\ker(p)$, and hence $\dim p(\widehat{C}) = \dim \widehat{C} - \dim \ker(p)$. Then $\text{codim}(\widehat{C}, \mathbb{R}^m) = \text{codim}(p(\widehat{C}), \mathbb{R}^n)$. Since $p(C)/p(\widehat{C})$ is spanned by the images of the $\text{codim}(\widehat{C}, \mathbb{R}^m)$ rays in $C \setminus \widehat{C}$, we deduce that $p(C)/p(\widehat{C})$ is simplicial. The final statement about the faces of $p(C)$ follows. \square

Definition 4.4.5. Assume that C is simplicial. For $0 \leq k \leq s$, let $C^{(i_k)}$ be the smallest face of C containing $C \cap \ker(\text{pr}_{S, S \setminus [i_k]})$.

For example, $\ker(\text{pr}_{S, S \setminus [i_0]}) = \mathbb{R}^S$ and $C^{(i_0)} = C$. Also, $C^{(i_s)} = \{0\}$.

Remark 4.4.6. It follows from Lemma 4.4.4 that if C is simplicial, then $\text{pr}_{S, S \setminus [i_k]}(C^{(i_k)})$ is the largest linear subspace contained in $\text{pr}_{S, S \setminus [i_k]}(C)$.

Corollary 4.4.7. Assume C is simplicial. For $1 \leq j \leq k \leq s$, $\text{pr}_{S, \{i_j\}}(C^{(i_k)}) \subset \mathbb{R}^{\{i_j\}}$ equals $\{0\}$ or $\mathbb{R}^{\{i_j\}}$.

Proof. Note that $\text{pr}_{S, \{i_j\}}$ factors through $\text{pr}_{S, S \setminus [i_k]}$, and, by Remark 4.4.6, $\text{pr}_{S, S \setminus [i_k]}(C^{(i_k)})$ is a linear space. This implies that $\text{pr}_{S, \{i_j\}}(C^{(i_k)})$ is a linear space. \square

Lemma 4.4.8. Assume that C is full-dimensional and simplicial. Fix $1 \leq k \leq s$. Assume that $(0, \dots, 0, 1) \in \text{pr}_{S, S \setminus [i_j]}(C) \subset \mathbb{R}^{S \setminus [i_j]}$ for $j > k$. Then $\text{pr}_{S, S \setminus [i_k]}(C)/\text{pr}_{S, S \setminus [i_k]}(C^{(i_k)})$ is a simplicial cone of dimension $\text{codim}(C^{(i_k)}, \mathbb{R}^S)$. The faces of $\text{pr}_{S, S \setminus [i_k]}(C)$ are

$$\{\text{pr}_{S, S \setminus [i_k]}(C') : C^{(i_k)} \subset C' \subset C\},$$

and $\text{codim}(\text{pr}_{S, S \setminus [i_k]}(C'), \mathbb{R}^{S \setminus [i_k]}) = \text{codim}(C', \mathbb{R}^S)$, for $C^{(i_k)} \subset C' \subset C$.

Proof. Consider an ordered set $U = \{u_{i_{k+1}}, \dots, u_{i_s}\}$ of elements of C such that $\text{pr}_{S, S \setminus [i_j]}(u_{i_j}) = (0, \dots, 0, 1)$ for $k < j \leq s$. Then U is a basis of $\ker(\text{pr}_{S, S \setminus [i_k]})$ contained in C , so $\text{span}(C \cap \ker(\text{pr}_{S, S \setminus [i_k]})) = \text{span}(C) \cap \ker(\text{pr}_{S, S \setminus [i_k]})$. The result then follows from Lemma 4.4.4. \square

Lemma 4.4.9. Assume that C is full-dimensional and simplicial. Fix $1 < k \leq s$. Assume that $(0, \dots, 0, 1) \in \text{pr}_{S, S \setminus [i_j]}(C) \subset \mathbb{R}^{S \setminus [i_j]}$ for $j > k$. Consider the projection map $p_k := \text{pr}_{S \setminus [i_k], S \setminus [i_{k-1}]} : \mathbb{R}^{S \setminus [i_k]} \rightarrow \mathbb{R}^{S \setminus [i_{k-1}]}$. Then, for $\epsilon > 0$ sufficiently small, there is a bijection

$\psi_\epsilon : \partial(\text{pr}_{S, S \setminus [i_k]}(C)) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \setminus [i_k]})) \rightarrow \{C^{(i_k)} \subset C' \subset C : \text{codim}(C', \mathbb{R}^S) = 1, \text{pr}_{S, S \setminus \{i_k\}}(C') \text{ is positive}\},$
such that $x \in \partial(\text{pr}_{S, S \setminus [i_k]}(C)) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \setminus [i_k]}))$ lies in the relative interior of the face $\text{pr}_{S, S \setminus [i_k]}(\psi_\epsilon(x)) \subset \text{pr}_{S, S \setminus [i_k]}(C)$, and $\{\psi_\epsilon^{-1}(C')\} = \text{pr}_{S, S \setminus [i_k]}(C') \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \setminus [i_k]}))$.

Proof. Applying Lemma 4.4.8, we have that the faces of $\text{pr}_{S, S \setminus [i_k]}(C)$ are $\{\text{pr}_{S, S \setminus [i_k]}(C') : C^{(i_k)} \subset C' \subset C\}$, and $\text{codim}(\text{pr}_{S, S \setminus [i_k]}(C'), \mathbb{R}^{S \setminus [i_k]}) = \text{codim}(C', \mathbb{R}^S)$ for $C^{(i_k)} \subset C' \subset C$. It follows from the assumption that $(0, \dots, 0, 1) \in \text{pr}_{S, S \setminus [i_j]}(C^{(i_k)}) \subset \mathbb{R}^{S \setminus [i_j]}$ for $j > k$. If $C^{(i_k)} \subset C'$, then it follows from Definition 4.4.1 that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive if and only if $\text{pr}_{S, S \setminus [i_{k-1}]}(C') = p_k(\text{pr}_{S, S \setminus [i_k]}(C'))$ is positive.

Suppose $C^{(i_k)} \subset C' \not\subset C$, and $x \in \text{pr}_{S, S \cap [i_k]}(C')$. Then $x \in p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ if and only if $p_k(x) = p_k(\vec{\epsilon}_{S \cap [i_k]}) = \vec{\epsilon}_{S \cap [i_{k-1}]}$. By Remark 4.4.2, for $\epsilon > 0$ sufficiently small, $\vec{\epsilon}_{S \cap [i_{k-1}]} \in p_k(\text{pr}_{S, S \cap [i_k]}(C'))$ implies that $p_k(\text{pr}_{S, S \cap [i_k]}(C')) \subset \mathbb{R}^{S \cap [i_{k-1}]}$ is full-dimensional, and hence $\text{codim}(\text{pr}_{S, S \cap [i_k]}(C'), \mathbb{R}^{S \cap [i_k]}) = \text{codim}(C', \mathbb{R}^S) = 1$, and $\text{pr}_{S, S \cap [i_k]}(C') \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ is a single point. By Lemma 4.4.3, we deduce that, for $\epsilon > 0$ sufficiently small, $\partial(\text{pr}_{S, S \cap [i_k]}(C')) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ is a union of points $\{x_{\epsilon, C'}\}$ indexed by the codimension 1 faces $C' \subset C$ such that $p_k(\text{pr}_{S, S \cap [i_k]}(C'))$ is positive, where $x_{\epsilon, C'}$ lies in the relative interior of $\text{pr}_{S, S \cap [i_k]}(C')$. \square

We continue with the assumptions and notation of Lemma 4.4.9. More specifically, $S = \{i_1 < i_2 < \dots < i_s\}$ is an ordered set, $C \subset \mathbb{R}^S$ is a full dimensional, simplicial cone, $1 < k \leq s$, and $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$. Then for $\epsilon > 0$ sufficiently small, there are 5 possibilities for $\text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$:

- (1) \emptyset ,
- (2) $x_\epsilon + \mathbb{R}_{\geq 0}(0, \dots, 0, 1)$,
- (3) $x_\epsilon + \mathbb{R}_{\geq 0}(0, \dots, 0, -1)$,
- (4) $[x_\epsilon, y_\epsilon]$,
- (5) $\text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]})) \cong \mathbb{R}$,

for some $x_\epsilon, y_\epsilon \in \mathbb{R}^{S \cap [i_k]}$. The corresponding form of $\partial(\text{pr}_{S, S \cap [i_k]}(C)) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ is $\emptyset, \{x_\epsilon\}, \{x_\epsilon\}, \{x_\epsilon, y_\epsilon\}$, and \emptyset respectively. Using the bijection ψ_ϵ , the number of codimension 1 faces $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive equals 0, 1, 1, 2, and 0 respectively.

In particular, given a codimension 1 face $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive, we can define an ‘orientation’ $\sigma(C') \in \{-1, 1\}$ by the condition that $(\psi_\epsilon^{-1}(C') + \mathbb{R}_{> 0}(0, \dots, 0, \sigma(C'))) \cap \text{pr}_{S, S \cap [i_k]}(C) \neq \emptyset$. For any other proper face $C^{(i_k)} \subset C' \not\subset C$, let $\sigma(C') = 0$. Let $\zeta_{C, k} := \sum_{C^{(i_k)} \subset C' \not\subset C} \sigma(C')$.

Lemma 4.4.10. *Assume that C is full-dimensional and simplicial. Fix $1 < k \leq s$. Assume that $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$. Then $\zeta_{C, k} \in \{-1, 0, 1\}$, and $\zeta_{C, k} \neq 0$ if and only if there is a unique $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive.*

Moreover, the following are equivalent:

- (1) $\zeta_{C, k} = 1$.
- (2) $\zeta_{C, k} \neq 0$ and $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_k]}(C)$.
- (3) $\zeta_{C, k} \neq 0$ and $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$.
- (4) C is positive and $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$.

Proof. From the above discussion and the definition of σ , $\zeta_{C, k} \neq 0$ if and only if $\text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ has the form either $x_\epsilon + \mathbb{R}_{\geq 0}(0, \dots, 0, 1)$ or $x_\epsilon + \mathbb{R}_{\geq 0}(0, \dots, 0, -1)$ for some x_ϵ , which holds if and only if the number of codimension 1 faces $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive equals 1. Note also that, since $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$, C is positive if and only if $\text{pr}_{S, S \cap [i_k]}(C)$ is positive.

In particular, $\zeta_{C, k} = 1$ if and only if $\text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ has the form $x_\epsilon + \mathbb{R}_{\geq 0}(0, \dots, 0, 1)$ for some x_ϵ . If $\zeta_{C, k} \neq 0$, then the latter condition holds if and only if $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_k]}(C)$ if and only if $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$. That is, conditions (1)-(3) are equivalent.

Assume that (4) holds. Then $\text{pr}_{S, S \cap [i_k]}(C)$ is positive, so, by Lemma 4.4.3, $\epsilon_{S \cap [i_k]} \in \text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ for $\epsilon > 0$ sufficiently small. By the definition of positive, $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_k]}(C)$. Hence $\text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ contains $\epsilon_{S \cap [i_k]} + \mathbb{R}_{\geq 0}(0, \dots, 0, 1)$. Since $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$, $\text{pr}_{S, S \cap [i_k]}(C) \cap p_k^{-1}(p_k(\vec{\epsilon}_{S \cap [i_k]}))$ has the form $x_\epsilon + \mathbb{R}_{\geq 0}(0, \dots, 0, 1)$ for some x_ϵ , and (1)-(3) hold.

Conversely, suppose (1)-(3) hold. By assumption, $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_k]}(C)$. There is a unique codimension 1 face $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive. Since $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive,

$$(0, \dots, 0, 1) \in \text{pr}_{S \setminus \{i_k\}, S \cap [i_j]}(\text{pr}_{S, S \setminus \{i_k\}}(C')) \subset \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$$

for $1 \leq j \leq k-1$. We conclude that $\text{pr}_{S, S \cap [i_k]}(C)$ is positive and (4) holds. \square

Let $U = \{u_{i_1}, \dots, u_{i_s}\}$ be an ordered set of elements of \mathbb{R}^S . Then U is *positively ordered* if, for all $1 \leq k \leq s$, $\{\text{pr}_{S, S \cap [i_k]}(u_j) : 1 \leq j \leq k\}$ spans a positive cone in $\mathbb{R}^{S \cap [i_k]}$. Given a permutation $w \in \text{Sym}_s$ and an ordered set $U = \{u_{i_1}, \dots, u_{i_s}\}$, we write $w(U) = \{u_{i_{w(1)}}, \dots, u_{i_{w(s)}}\}$.

Lemma 4.4.11. *Assume that C is simplicial and positive. Let $U = \{u_{i_1}, \dots, u_{i_s}\}$ be an ordered set of generators of the rays of C . Then there exists a unique permutation $w \in \text{Sym}_s$ such that $w(U)$ is positively ordered.*

Proof. We may assume that $s > 1$. By Lemma 4.4.10, there exists a unique codimension 1 face $C' \subset C$ such that $\text{pr}_{S, S \cap [i_{s-1}]}(C')$ is positive. Let $u_{i_{w(s)}}$ be the unique element of U in $C \setminus C'$. Then C' is also simplicial, so the result follows by induction. \square

Lemma 4.4.12. *Assume that C is simplicial and positive. Fix a positively ordered set of generators $U = \{u_{i_1}, \dots, u_{i_s}\}$ of the rays of C . Then $\det(A_U) > 0$.*

Proof. The assumptions imply that there are $|S| \times |S|$ matrices \mathcal{U}_U and \mathcal{L}_U such that $A_U \mathcal{U}_U^{-1} = \mathcal{L}_U$, \mathcal{U}_U^{-1} has nonnegative entries and is upper triangular, and \mathcal{L}_U is lower triangular with all 1s on the diagonal. \square

Assume that C is full-dimensional and simplicial. Let $U = \{u_{i_1}, \dots, u_{i_s}\}$ be an ordered set of generators of the rays of C . Fix $1 \leq k \leq s$. Then U is *i_k -weakly positively ordered* or *i_k -WPO* if the ordered set $\text{pr}_{S, S \setminus \{i_k\}}(U|_{S \setminus \{i_k\}})$ of elements in $\mathbb{R}^{S \setminus \{i_k\}}$ is positively ordered and $u_{i_k} \notin C^{(i_k)}$. By Lemma 4.4.11, there is a bijection between permutations $w \in \text{Sym}_s$ such that $w(U)$ is i_k -WPO, and codimension 1 faces $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive, where $w(U)|_{S \setminus \{i_k\}}$ is a positively ordered set of generators of the rays of C' , and $u_{i_{w(k)}}$ generates the unique ray in $C \setminus C'$.

We say that a nonzero element $x \in \mathbb{R}$ has $\text{sign } x/|x| \in \{-1, 1\}$.

Lemma 4.4.13. *Assume that C is full-dimensional and simplicial. Fix $1 < k \leq s$. Assume that $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$. Let $U = \{u_{i_1}, \dots, u_{i_s}\}$ be an ordered set of generators of the rays of C . Assume that U is i_k -WPO and $U|_{S \setminus \{i_k\}}$ is a set of generators of the rays of $C^{(i_k)} \subset C' \not\subset C$. Then $\det(A_U)$ has sign $\sigma(C') \in \{-1, 1\}$.*

Proof. Let $U' = \text{pr}_{S, S \setminus \{i_k\}}(U|_{S \setminus \{i_k\}})$. Since U' is positively ordered, $\det(A_{U'}) > 0$ by Remark 4.4.12. Since $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$, we can replace u_{i_k} by adding an element of $\ker(\text{pr}_{S, S \cap [i_k]}(C))$ without changing $\det(A_U)$. By Lemma 4.4.8, $\text{pr}_{S, S \cap [i_k]}(C')$ has codimension 1 in $\mathbb{R}^{S \cap [i_k]}$. It follows that we can write

$$\text{pr}_{S, S \cap [i_k]}(u_{i_k}) = \alpha(0, \dots, 0, 1) \in \mathbb{R}^{S \cap [i_k]} / \text{span}(\text{pr}_{S, S \cap [i_k]}(C')),$$

for some $\alpha \neq 0$ with $\text{sign } \sigma(C')$. The result follows since $\det(A_U) = \alpha \det(A_{U'})$. \square

The following result is the main conclusion of our study of positive cones.

Proposition 4.4.14. *Assume C is full-dimensional and simplicial. Let $U = \{u_{i_1}, \dots, u_{i_s}\}$ be an ordered set of generators of the rays of C . Fix $1 < k \leq s$, and assume $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C)$ for $j > k$. Let*

$$\lambda(U) := \sum_{\substack{w \in \text{Sym}_s \\ w(U) \text{ is } i_k\text{-WPO}}} \det(A_{w(U)})$$

Then $\lambda(U) \neq 0$ if and only if there is a unique permutation $w \in \text{Sym}_s$ such that $w(U)$ is i_k -WPO. Moreover, the following are equivalent:

- (1) $\lambda(U) > 0$,
- (2) $\lambda(U) \neq 0$ and $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_k]}(C)$,
- (3) $\lambda(U) \neq 0$ and $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$, and
- (4) C is positive and $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$.

Proof. By the above discussion, there is a bijection between permutations $w \in \text{Sym}_s$ such that $w(U)$ is i_k -WPO, and codimension 1 faces $C^{(i_k)} \subset C' \not\subset C$ such that $\text{pr}_{S, S \setminus \{i_k\}}(C')$ is positive. By Lemma 4.4.13, if w corresponds to C' under this bijection, then $\det(A_{w(U)}) = \sigma(C') |\det(A_U)|$. Hence $\lambda(U) = |\det(A_U)| \zeta_{C,k}$. The result now follows from Lemma 4.4.10. \square

4.4.3. *Expansion.* We now develop some tools that will help us expand $y^{A_1} y^{A_2} \dots y^{A_r} \in H^*(F \sqcup E)$ as a sum of squarefree products of vertices of $\text{lk}_S(F \sqcup E)$.

Lemma 4.4.15. *Let $S \subset [r]$ be a nonempty subset, and let G be a face of $\text{lk}_S(F \sqcup E)$. Assume that $\{2j-1, 2j\} \subset \text{Supp}_G$ for all $j \in S$. Then $y^G(\prod_{j \notin S} y^{A_j}) = 0$ in $H^*(F \sqcup E)$.*

Proof. The proof proceeds by induction on $r - |S|$. We first show that the assumption that F is a maximal non- U -pyramid means that $r - |S| = 0$ is impossible. Indeed, there exists $1 \leq i \leq r$ such that precisely one of $\{2i-1, 2i\}$ lies in Supp_G . In particular, $i \notin S$, and hence $S \neq [r]$ and $r - |S| > 0$. Without loss of generality, assume that $2i \in \text{Supp}_G$ and $2i-1 \notin \text{Supp}_G$. We apply (18) with the global linear function e_{2i-1} to get the relation in $H^*(E)$:

$$x^{A_i} + \sum_{\substack{v \in \text{lk}_S(E) \\ v \notin \{A_1, \dots, A_r\}}} v_{2i-1} x^v = 0.$$

Let $v \in \text{lk}_S(E)$ be a vertex with $v_{2i-1} \neq 0$ appearing in the above sum. Since A_i is an apex of $F \sqcup G \sqcup E$ with base direction e_{2i-1}^* , it follows that $v \notin F \sqcup G \sqcup E$, and $y^v y^G = 0$ in $H^*(F \sqcup E)$ unless $G \sqcup v$ is a face in $\text{link}_S(F \sqcup E)$. In the latter case, $\{2i-1, 2i\} \in \text{Supp}_{G \sqcup v}$. We now compute in $H^*(F \sqcup E)$:

$$y^G(\prod_{j \notin S} y^{A_j}) = - \sum_{v \in \text{lk}_S(F \sqcup E)} v_{2i-1} y^v y^G (\prod_{j \notin S \cup \{i\}} y^{A_j}).$$

Finally, the induction hypothesis implies that if $v_{2i-1} \neq 0$, then $y^{G \sqcup v}(\prod_{j \notin S \cup \{i\}} y^{A_j}) = 0$. \square

Definition 4.4.16. *Let $R \subset [r]$. Let $\pi_R: \mathbb{R}^n \rightarrow \mathbb{R}^r$ be the linear function defined by*

$$\pi_R(x)_i = \begin{cases} \pi(x)_i = x_{2i-1} - x_{2i} & \text{if } i \notin R, \\ x_{2i-1} & \text{if } i \in R. \end{cases}$$

Definition 4.4.17. *Let $R, S, T \subset [r]$. Consider an ordered set V of elements of \mathbb{R}^n indexed by T . If S and T are nonempty, we may consider the ordered set $\text{pr}_{[r], S}(\pi_R(V))$ of elements of \mathbb{R}^S indexed by T , and we define the $|S| \times |T|$ matrix $A_{V, S, R} := A_{\text{pr}_{[r], S}(\pi_R(V))}$.*

Let G be a nonempty face in $\text{lk}_S(F \sqcup E)$, and assume that $S \neq \emptyset$. Let V_G be an ordered set consisting of the vertices of G , indexed by T . Let $C_{G, S, R} \subset \mathbb{R}^S$ be the cone spanned by the columns of $A_{V_G, S, R}$. If $G = \emptyset$ or $S = \emptyset$, then $C_{G, S, R} = \{0\}$. If $|G| = |S|$, we define the multiplicity of (G, S, R) to be $\text{mult}(G, S, R) := |\det(A_{V_G, S, R})|$. If $G = S = \emptyset$, then $\text{mult}(G, S, R) = 1$.

For example, if G is a face in $\text{lk}_S(F \sqcup E)$, $S = [r]$, and $R = \emptyset$, then $C_{G, S, R} = \pi(C_G)$, where $C_G \in \Delta$ is the cone over G in \mathbb{R}^n . Note that the definitions of $C_{G, S, R}$ and $\text{mult}(G, S, R)$ above are independent of the choice of an ordering V_G of the vertices of G .

Proposition 4.4.18. *Let H be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \dots < i_s\} \subset [r]$ for some $s > 1$. Fix $1 \leq k \leq s$, and assume that $k < j$ for all $i_j \in R$. Assume that $|H| = s - 1$ and $(0, \dots, 0, 1) \in C_{H, S \cap [i_j], R \cup \{i_k\}} \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$. Consider an ordering V_H of the vertices of H , indexed by $S \setminus \{i_k\}$. If $H \subset G$ and $|G| = s$, consider the unique ordering $V_{H,G}$ of the vertices of G indexed by S such that $V_{H,G}|_{S \setminus \{i_k\}} = V_H$. Then we have the following equality in $H^*(F \sqcup E)$:*

$$\det(A_{V_H, S \setminus \{i_k\}, R}) y^H \left(\prod_{i \notin S \setminus \{i_k\}} y^{A_i} \right) = - \sum_{\substack{H \subset G \\ |G|=s}} \det(A_{V_{H,G}, S, R \cup \{i_k\}}) y^G \left(\prod_{i \in S} y^{A_i} \right).$$

Proof. Our goal is to expand $y^{A_{i_k}}$ on the left hand side by constructing a linear function and applying (18). We define a linear function $\mu = \mu_{H, S, R, i_k} : \mathbb{R}^n \rightarrow \mathbb{R}$ as follows: if $x \in \mathbb{R}^n$, then

$$\mu(x) = \det(A_{V_x, S, R \cup \{i_k\}}),$$

where V_x is the ordered set indexed by S defined by $V_x|_{S \setminus \{i_k\}} = V_H$ and $V_x|_{\{i_k\}} = \{x\}$. By definition, $\mu(x) = 0$ if x is a vertex of H . If x is a vertex of $F \sqcup E$ and $j \leq k$, then

$$(A_{V_x, S, R \cup \{i_k\}})_{i_j, i_k} = \pi_R(x)_{i_j} = \begin{cases} 1 & \text{if } j = k, x = A_{i_k}, \\ 0 & \text{otherwise.} \end{cases}$$

Since $(0, \dots, 0, 1) \in C_{H, S \cap [i_j], R \cup \{i_k\}} \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$, it follows that we can apply column operations to ensure that $(A_{V_x, S, R \cup \{i_k\}})_{i_j, i_k} = 0$ for $j > k$, without affecting $\mu(x)$ or the values of any other elements of $A_{V_x, S, R \cup \{i_k\}}$. We deduce that if x is a vertex of $F \sqcup E$, then $\mu(x) = \det(A_{V_H, S \setminus \{i_k\}, R})$ if $x = A_{i_k}$, and $\mu(x) = 0$ otherwise.

We apply (18) with the global linear function μ to get the relation in $H^*(E)$:

$$\mu(A_{i_k}) x^{A_{i_k}} + \sum_{\substack{v \in \text{lk}_S(E) \\ v \notin F \sqcup H}} \mu(v) x^v = 0.$$

Note that if $v \notin F \sqcup H$, then $y^v y^H = 0$ in $H^*(F \sqcup E)$ unless $H \sqcup v$ is a face in $\text{lk}_S(F \sqcup E)$. Considering the image of the above equality in $H^*(F \sqcup E)$ and multiplying both sides by $y^H (\prod_{i \notin S} y^{A_i})$ gives the result. \square

We will use the formula in Proposition 4.4.18 to repeatedly to expand $y^{A_1} \dots y^{A_r}$. The terms that appear at some point in this expansion depend on a tuple (\widehat{H}, S, R, i_k) , where \widehat{H} is a face of $\text{lk}_S(F \sqcup E)$, $R \subset S$, and $0 \leq k \leq s$. We introduce two concepts that will be used to control terms in this expansion.

Definition 4.4.19. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \dots < i_s\} \subset [r]$. Fix $0 \leq k \leq s$. Then (\widehat{H}, S, R, i_k) is material if $C_{\widehat{H}, S, R}$ is simplicial, $C_{\widehat{H}, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$, and $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C_{\widehat{H}, S, R}) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$.*

Remark 4.4.20. When $k = 0$, the condition $C_{\widehat{H}, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$ holds immediately. When $k = s$, the condition $C_{\widehat{H}, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$ holds if and only if $C_{\widehat{H}, S, R} = \{0\}$. It follows that if $\widehat{H} = \emptyset$ and $k = s$, then (\widehat{H}, S, R, i_k) is material.

Remark 4.4.21. Note that if (\widehat{H}, S, R, i_k) is material, then $\ker(\text{pr}_{S, S \cap [i_k]}) \subset \text{span}(C_{\widehat{H}, S, R})$. Indeed, by assumption there exists $u_1, \dots, u_{s-k} \in C_{\widehat{H}, S, R}$ such that $\text{pr}_{S, S \cap [i_{k+j}]}(u_j) = (0, \dots, 0, 1)$ for $1 \leq j \leq s - k$. Since $\dim(\ker(\text{pr}_{S, S \cap [i_k]})) = s - k$, it follows that $\ker(\text{pr}_{S, S \cap [i_k]}) = \text{span}(u_1, \dots, u_{s-k}) \subset \text{span}(C_{\widehat{H}, S, R})$.

Lemma 4.4.22. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \dots < i_s\} \subset [r]$. Fix $1 \leq k \leq s$. Assume that $(v_{2i_k-1}, v_{2i_k}) = (0, 0)$ for all vertices $v \in \widehat{H}$. Then (\widehat{H}, S, R, i_k) is material if and only if $(\widehat{H}, S, R \cup \{i_k\}, i_k)$ is material. Moreover, if (\widehat{H}, S, R, i_k) is material, then $(\widehat{H}, S \setminus \{i_k\}, R, i_{k-1})$ is material.*

Proof. For all vertices $v \in \widehat{H}$, $(v_{2i_k-1}, v_{2i_k}) = (0, 0)$, and hence $\pi_R(v)_{i_k} = \pi_{R \cup \{i_k\}}(v)_{i_k} = 0$. It follows that $C_{\widehat{H}, S, R} = C_{\widehat{H}, S, R \cup \{i_k\}}$. This establishes the first statement, so we now prove the second statement. When $k = 1$, the claim is immediate, so we may assume that $1 < k \leq s$. Let $\psi: C_{\widehat{H}, S, R} \rightarrow C_{\widehat{H}, S \setminus \{i_k\}, R}$ denote the restriction of $\text{pr}_{S, S \setminus \{i_k\}}$. Then ψ is surjective by definition. Since $\pi_R(v)_{i_k} = 0$ for all vertices $v \in \widehat{H}$, $\ker(\text{pr}_{S, S \setminus \{i_k\}}) \cap C_{\widehat{H}, S, R} = \{0\}$, and hence ψ is an isomorphism of cones. Also, $\text{pr}_{S, S \setminus \{i_k\}}$ induces an isomorphism from $\ker(\text{pr}_{S, S \cap [i_k]})$ to $\ker(\text{pr}_{S \setminus \{i_k\}, S \cap [i_k-1]})$. It follows that $\psi(C_{\widehat{H}, S, R}^{(i_k)}) = C_{\widehat{H}, S \setminus \{i_k\}, R}^{(i_{k-1})}$, which implies the result. \square

Lemma 4.4.23. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \dots < i_s\} \subset [r]$. Fix $1 \leq k \leq s$. Assume that $i_k \notin R$ and (\widehat{H}, S, R, i_k) is material. Then either $\{2i_k - 1, 2i_k\} \subset \text{Supp}_{\widehat{H}}$ or $(v_{2i_k-1}, v_{2i_k}) = (0, 0)$ for all vertices $v \in \widehat{H}$.*

Proof. By Corollary 4.4.7, $\text{pr}_{S, \{i_k\}}(C_{\widehat{H}, S, R})$ is $\{0\}$ or $\mathbb{R}^{\{i_k\}}$. Assume there is a vertex $v \in \widehat{H}$ such that $(v_{2i_k-1}, v_{2i_k}) \neq (0, 0)$. If $\pi(v)_{i_k} = 0$, then $v_{2i_k-1} = v_{2i_k} \neq 0$. If $\pi(v)_{i_k} \neq 0$, then $\text{pr}_{S, \{i_k\}}(C_{\widehat{H}, S, R}) = \mathbb{R}^{\{i_k\}}$, and there is $v' \in \widehat{H}$ such that $\pi(v')_{i_k} \neq 0$ has the opposite sign to $\pi(v)_{i_k}$. In either case, $2i_k - 1, 2i_k \in \text{Supp}_{\widehat{H}}$. \square

Let G be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \dots < i_s\} \subset [r]$. Assume that $|G| = s$. Then we say that (G, S, R) is *positive* if $C_{G, S, R} \subset \mathbb{R}^S$ is positive. An ordering V of the vertices of G indexed by S is *positively ordered* if $\text{pr}_{[r], S}(\pi_R(V))$ is positively ordered. For example, if $G = S = R = \emptyset$, then $C_{G, S, R} = \mathbb{R}^S = \{0\}$ and (G, S, R) is positive.

Remark 4.4.24. If (G, S, R) is positive then $C_{G, S, R} \subset \mathbb{R}^S$ is a full-dimensional simplicial cone, with faces $\{C_{H, S, R} : H \subset G\}$. Also, in this case, Lemma 4.4.11 implies that there is a unique positive ordering of the vertices of G .

4.4.4. *Computing the degree.* We now define an element of $H^*(F \sqcup E)$ that depends on a tuple (\widehat{H}, S, R, i_k) , which we call $\theta(\widehat{H}, S, R, i_k)$. For some tuples, the formula for $\theta(\widehat{H}, S, R, i_k)$ is very simple, and it behaves well with respect to the formula in Proposition 4.4.18. This allows us to compute the degree of $y^{A_1} \dots y^{A_r}$.

Definition 4.4.25. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \dots < i_s\} \subset [r]$. Fix $0 \leq k \leq s$. We define an element $\theta(\widehat{H}, S, R, i_k) \in H^*(F \sqcup E)$ by*

$$\theta(\widehat{H}, S, R, i_k) := \sum_{\substack{\widehat{H} \subset G, |G|=s \\ C_{G, S, R}^{(i_k)} = C_{\widehat{H}, S, R} \\ (G, S, R) \text{ positive}}} \text{mult}(G, S, R) y^G \left(\prod_{i \notin S} y^{A_i} \right).$$

Remark 4.4.26. We highlight a special case of the above definition. We have that

$$\theta(\widehat{H}, S, R, i_0) = \begin{cases} \text{mult}(\widehat{H}, S, R) y^{\widehat{H}} (\prod_{i \notin S} y^{A_i}) & \text{if } |\widehat{H}| = s \text{ and } (\widehat{H}, S, R) \text{ is positive,} \\ 0 & \text{otherwise.} \end{cases}$$

In particular, if $S = \emptyset$ (and therefore $R = \emptyset$ and $k = 0$),

$$\theta(\widehat{H}, S, R, i_k) = \begin{cases} y^{A_1} \dots y^{A_r} & \text{if } \widehat{H} = \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

We now state four properties of θ . They will allow us to compute $\theta(\emptyset, \emptyset, \emptyset, i_0) = y^{A_1} \cdots y^{A_r}$ via a recursion, which we use to complete the proof of Proposition 4.3.5. Afterwards, we will prove the four properties.

Lemma 4.4.27. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \cdots < i_s\} \subset [r]$. Fix $0 \leq k \leq s$. Then $\theta(\widehat{H}, S, R, i_k) \neq 0$ implies that (\widehat{H}, S, R, i_k) is material. If (\widehat{H}, S, R, i_k) is material and $\widehat{H} \subset G$ such that $|G| = s$ and (G, S, R) is positive, then $C_{G, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$. In particular, if (\widehat{H}, S, R, i_k) is material, then*

$$\theta(\widehat{H}, S, R, i_k) = \sum_{\substack{\widehat{H} \subset G, |G|=s \\ (G, S, R) \text{ positive}}} \text{mult}(G, S, R) y^G \left(\prod_{i \notin S} y^{A_i} \right).$$

Lemma 4.4.28. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \cdots < i_s\} \subset [r]$. Fix $1 \leq k \leq s$. If (\widehat{H}, S, R, i_k) is material, then*

$$\theta(\widehat{H}, S, R, i_k) = \sum_{\widehat{H} \subset \widehat{H}'} \theta(\widehat{H}', S, R, i_{k-1}).$$

Lemma 4.4.29. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \cdots < i_s\} \subset [r]$. Fix $1 \leq k \leq s$. Assume that $\{2i_k - 1, 2i_k\} \not\subset \text{Supp}_{\widehat{H}}$. Then*

$$\theta(\widehat{H}, S, R, i_{k-1}) = \theta(\widehat{H}, S, R \cup \{i_k\}, i_{k-1}).$$

Lemma 4.4.30. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \cdots < i_s\} \subset [r]$. Fix $1 \leq k \leq s$. Assume that $k < j \leq s$ for all $i_j \in R$. Assume that $(\widehat{H}, S, R \cup \{i_k\}, i_k)$ and $(\widehat{H}, S \setminus \{i_k\}, R, i_{k-1})$ are material. Then*

$$\theta(\widehat{H}, S, R \cup \{i_k\}, i_k) = -\theta(\widehat{H}, S \setminus \{i_k\}, R, i_{k-1}).$$

Proposition 4.4.31. *Let \widehat{H} be a face in $\text{lk}_S(F \sqcup E)$. Let $R \subset S = \{i_1 < \cdots < i_s\} \subset [r]$. Fix $0 \leq k \leq s$. Assume that $k < j \leq s$ for all $i_j \in R$. Assume that $2i_j - 1, 2i_j \in \text{Supp}_{\widehat{H}}$ for $k < j \leq s$. Then*

$$\theta(\widehat{H}, S, R, i_k) = \begin{cases} (-1)^s y^{A_1} \cdots y^{A_r} & \text{if } \widehat{H} = R = \emptyset, k = s \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We proceed by induction on k . Assume first that $k = 0$. If $s = 0$, then the result follows from Remark 4.4.26. Assume that $s > 0$. By Remark 4.4.26,

$$\theta(\widehat{H}, S, R, i_0) = \begin{cases} \text{mult}(\widehat{H}, S, R) y^{\widehat{H}} \left(\prod_{i \notin S} y^{A_i} \right) & \text{if } |\widehat{H}| = s, (\widehat{H}, S, R) \text{ is positive} \\ 0 & \text{otherwise.} \end{cases}$$

By assumption, $2i_j - 1, 2i_j \in \text{Supp}_{\widehat{H}}$ for all $1 \leq j \leq s$. Then Lemma 4.4.15 implies that $\theta(\widehat{H}, S, R, i_0) = 0$.

Assume that $k > 0$. In particular, $s > 0$. By Lemma 4.4.27 and Remark 4.4.20, we may assume that (\widehat{H}, S, R, i_k) is material. By Lemma 4.4.28,

$$\theta(\widehat{H}, S, R, i_k) = \sum_{\widehat{H} \subset \widehat{H}'} \theta(\widehat{H}', S, R, i_{k-1}).$$

By induction, this simplifies to

$$(20) \quad \theta(\widehat{H}, S, R, i_k) = \sum_{\substack{\widehat{H} \subset \widehat{H}' \\ \{2i_k - 1, 2i_k\} \not\subset \text{Supp}_{\widehat{H}'}}} \theta(\widehat{H}', S, R, i_{k-1}).$$

In particular, we may assume that $\{2i_k - 1, 2i_k\} \not\subset \text{Supp}_{\widehat{H}}$, else the right-hand side of (20) is zero, and the result holds. Since $i_k \notin R$ by assumption, Lemma 4.4.23 implies that $(v_{2i_k - 1}, v_{2i_k}) = (0, 0)$ for all

vertices $v \in \widehat{H}$. By Lemma 4.4.22, $(\widehat{H}, S, R \cup \{i_k\}, i_k)$ and $(\widehat{H}, S \setminus \{i_k\}, R, i_{k-1})$ are material. As above, by Lemma 4.4.28 and the induction hypothesis,

$$\theta(\widehat{H}, S, R \cup \{i_k\}, i_k) = \sum_{\substack{\widehat{H} \subset \widehat{H}' \\ \{2i_k-1, 2i_k\} \not\subset \text{Supp}_{\widehat{H}'}}} \theta(\widehat{H}', S, R \cup \{i_k\}, i_{k-1}).$$

Comparing with (20), Lemma 4.4.29 implies that

$$\theta(\widehat{H}, S, R, i_k) = \theta(\widehat{H}, S, R \cup \{i_k\}, i_k).$$

By Lemma 4.4.30,

$$\theta(\widehat{H}, S, R \cup \{i_k\}, i_k) = -\theta(\widehat{H}, S \setminus \{i_k\}, R, i_{k-1}).$$

The result now follows by induction. \square

Proof of Proposition 4.3.5. Recall that for each face $G \in \text{lk}_S(F \sqcup E)$, $C_{G, [r], R} = \pi(C_G)$, where $C_G \in \Delta$ is the cone over G in \mathbb{R}^n . Moreover, the cones $\{\pi(C_G) : G \in \text{lk}_S(F \sqcup E)\}$ form a complete fan. Therefore, there is a unique facet $\widehat{G} \in \text{lk}_S(F \sqcup E)$ such that $\pi(C_{\widehat{G}})$ is positive. By Definition 4.4.25, $\theta(\emptyset, [r], \emptyset, i_r) = \text{mult}(\widehat{G}, [r], \emptyset) y^{\widehat{G}}$. By Proposition 4.4.31, we have the following equality in $H^*(F \sqcup E)$:

$$(-1)^r y^{A_1} \cdots y^{A_r} = \text{mult}(\widehat{G}, [r], \emptyset) y^{\widehat{G}}.$$

Since $\text{mult}(\widehat{G}, [r], \emptyset)$ and $\deg_{F \sqcup E}(y^{\widehat{G}})$ are both strictly positive, we deduce that $(-1)^r \deg_{F \sqcup E}(y^{A_1} \cdots y^{A_r}) > 0$. \square

Proof of Lemma 4.4.27. If $\theta(\widehat{H}, S, R, i_k) \neq 0$, then there is a face $G \supset \widehat{H}$ with $|G| = s$, $C_{G, S, R}^{(i_k)} = C_{\widehat{H}, S, E}$, and (G, S, R) positive. As $C_{G, S, R}^{(i_k)}$ is a face of the simplicial cone $C_{G, S, R}$, $C_{\widehat{H}, S, E} = C_{G, S, R}^{(i_k)}$ is simplicial. We also see that $C_{\widehat{H}, S, E}^{(i_k)} = C_{\widehat{H}, S, E}$. The positivity of (G, S, R) implies that $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C_{\widehat{H}, S, E})$ for $j > k$, so (\widehat{H}, S, R, i_k) is material.

Suppose (\widehat{H}, S, R, i_k) is material and $G \supset \widehat{H}$ with $|G| = s$ and (G, S, R) positive. By Remark 4.4.24, $C_{G, S, R} \subset \mathbb{R}^S$ is a full-dimensional simplicial cone with faces $\{C_{H, S, R} : H \subset G\}$. Then $C_{\widehat{H}, S, E}$ is a face of $C_{G, S, R}$, and it follows from Remark 4.4.21 that $C_{G, S, R}^{(i_k)} = C_{\widehat{H}, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$. Then the formula for $\theta(\widehat{H}, S, R, i_k)$ is immediate. \square

Proof of Lemma 4.4.28. Let G be a face of $\text{lk}_S(F \sqcup E)$ such that $|G| = s$ and (G, S, R) is positive. Let $\widehat{H}' \subset G$ be the unique face such that $C_{G, S, R}^{(i_{k-1})} = C_{\widehat{H}', S, R}$. By Lemma 4.4.27, if $\widehat{H} \subset G$, then $C_{G, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$. It is enough to show that $\widehat{H} \subset G$ if and only if $\widehat{H} \subset \widehat{H}'$. Clearly, if $\widehat{H} \subset \widehat{H}'$ then $\widehat{H} \subset G$. If $\widehat{H} \subset G$, then $C_{G, S, R}^{(i_k)} = C_{\widehat{H}, S, R}$ which is a face of $C_{G, S, R}^{(i_{k-1})} = C_{\widehat{H}', S, R}$, and the result follows from Remark 4.4.24. \square

Proof of Lemma 4.4.29. Let R_1, R_2 be sets such that $\{R_1, R_2\} = \{R, R \cup \{i_k\}\}$. Consider a face $G \supset \widehat{H}$ such that $|G| = s$. Assume that $C_{G, S, R_1}^{(i_{k-1})} \subset C_{\widehat{H}, S, R_1}$ and (G, S, R_1) is positive. By definition, $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C_{G, S, R_1}^{(i_{k-1})})$ for $k \leq j \leq s$. In particular, $\text{pr}_{S, \{i_k\}}(C_{G, S, R_1}^{(i_{k-1})}) \not\subset \mathbb{R}_{\leq 0}$. Since $C_{G, S, R_1}^{(i_{k-1})} \subset C_{\widehat{H}, S, R_1}$ and $\pi_{R_1}(v)_{i_k} \in \{\pi(v)_{i_k}, v_{2i_k-1}\}$ for $v \in \widehat{H}$, we deduce that $2i_k-1 \in \text{Supp}_{\widehat{H}}$. Since $\{2i_k-1, 2i_k\} \not\subset \text{Supp}_{\widehat{H}}$, we have $2i_k \notin \text{Supp}_{\widehat{H}}$, and hence $\pi_{R_1}(v)_{i_k} = \pi_{R_2}(v)_{i_k}$ for $v \in \widehat{H}$. We deduce that $C_{G, S, R_1}^{(i_{k-1})} \subset C_{\widehat{H}, S, R_1} = C_{\widehat{H}, S, R_2} \subset C_{G, S, R_2}$. In particular, $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C_{G, S, R_2})$ for $k \leq j \leq s$, and $C_{G, S, R_2}^{(i_{k-1})} \subset C_{G, S, R_1}^{(i_{k-1})} \subset C_{\widehat{H}, S, R_2}$. For $1 \leq j < k$, $(0, \dots, 0, 1) \in \text{pr}_{S, S \cap [i_j]}(C_{G, S, R_1}) = \text{pr}_{S, S \cap [i_j]}(C_{G, S, R_2})$ by definition. We deduce that (G, S, R_2) is positive.

We conclude that the condition that both $C_{G,S,R_1}^{(i_{k-1})} \subset C_{\widehat{H},S,R_1}$ and (G, S, R_1) is positive is independent of the choice of R_1 . Moreover, if this condition holds, then the above argument shows that both $C_{G,S,R_1}^{(i_{k-1})}$ and $C_{\widehat{H},S,R_1}$ are independent of the choice of R_1 . It follows that the condition that both $C_{G,S,R_1}^{(i_{k-1})} = C_{\widehat{H},S,R_1}$ and (G, S, R_1) is positive is also independent of the choice of R_1 , as desired. \square

Proof of Lemma 4.4.30. By Lemma 4.4.27, $\theta(\widehat{H}, S \setminus \{i_k\}, R, i_{k-1})$ equals

$$\sum_{\substack{\widehat{H} \subset H, |H|=s-1 \\ (H, S \setminus \{i_k\}, R) \text{ positive}}} \text{mult}(H, S \setminus \{i_k\}, R) y^H \left(\prod_{i \notin S \setminus \{i_k\}} y^{A_i} \right).$$

Consider a face $\widehat{H} \subset H$ with $|H| = s - 1$ and $(H, S \setminus \{i_k\}, R)$ positive. Let V_H be the unique positive ordering of the vertices of H , indexed by $S \setminus \{i_k\}$. If $H \subset G$ and $|G| = s$, consider the unique ordering $V_{H,G}$ of the vertices of G indexed by S such that $V_{H,G}|_{S \setminus \{i_k\}} = V_H$. The above expression equals

$$\sum_{\substack{\widehat{H} \subset H, |H|=s-1 \\ (H, S \setminus \{i_k\}, R) \text{ positive}}} \det(A_{V_{H,S \setminus \{i_k\}, R}}) y^H \left(\prod_{i \notin S \setminus \{i_k\}} y^{A_i} \right).$$

Since $(\widehat{H}, S, R \cup \{i_k\}, i_k)$ is material, $(0, \dots, 0, 1) \in C_{\widehat{H}, S \cap [i_j], R \cup \{i_k\}} \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$. Hence we can apply Proposition 4.4.18 to obtain that the above expression is equal to

$$\begin{aligned} & - \sum_{\substack{\widehat{H} \subset H, |H|=s-1 \\ (H, S \setminus \{i_k\}, R) \text{ positive}}} \sum_{\substack{H \subset G \\ |G|=s}} \det(A_{V_{H,G,S,R \cup \{i_k\}}}) y^G \left(\prod_{i \notin S} y^{A_i} \right) \\ & = - \sum_{\substack{\widehat{H} \subset G \\ |G|=s}} \sum_{\substack{\widehat{H} \subset H \subset G, |H|=s-1 \\ (H, S \setminus \{i_k\}, R) \text{ positive}}} \det(A_{V_{H,G,S,R \cup \{i_k\}}}) y^G \left(\prod_{i \notin S} y^{A_i} \right). \end{aligned}$$

Consider a face $\widehat{H} \subset G$ with $|G| = s$. Let $C = C_{G,S,R \cup \{i_k\}}$. If C is not full-dimensional, then we have $\det(A_{V_{H,G,S,R \cup \{i_k\}}}) = 0$ for all choices of H . We may assume that C is full-dimensional, and hence simplicial. As above, $(0, \dots, 0, 1) \in C_{\widehat{H}, S \cap [i_j], R \cup \{i_k\}} \subset \text{pr}_{S, S \cap [i_j]}(C) \subset \mathbb{R}^{S \cap [i_j]}$ for $j > k$. Also, $\pi_{R \cup \{i_k\}}(v)_{i_k} = v_{2i_k-1} \geq 0$ for all $v \in G$, and hence $(0, \dots, 0, -1) \notin \text{pr}_{S, S \cap [i_k]}(C)$. Then Proposition 4.4.14 implies that the above expression simplifies to give

$$= - \sum_{\substack{\widehat{H} \subset G, |G|=s \\ (G, S, R \cup \{i_k\}) \text{ positive}}} \text{mult}(G, S, R \cup \{i_k\}) y^G \left(\prod_{i \notin S} y^{A_i} \right).$$

Since $(\widehat{H}, S, R \cup \{i_k\}, i_k)$ is material, Lemma 4.4.27 implies that the latter equals $-\theta(\widehat{H}, S, R \cup \{i_k\}, i_k)$. \square

5. THE LOCAL FORMAL ZETA FUNCTION AND ITS CANDIDATE POLES

5.1. Overview. In this section, we introduce the local formal zeta function (Definition 5.3.1) and develop its fundamental properties. In Section 5.2, we recall the formula for $Z_{\text{mot}}(T)$ in [BN20, Theorem 8.3.5]. In Section 5.3, we define the local formal zeta function and discuss its candidate poles. In Section 5.4, we prove a relation that the local formal zeta function satisfies that will be a crucial tool in the proof of Theorem 1.4.7.

We first introduce some notation for use in this section and in Section 6. We use capital letters to denote elements of the vector space containing $\text{Newt}(f)$ and use lowercase letters to denote elements of the dual space. Let Γ be the union of the proper interior faces of $\text{Newt}(f)$ and their subfaces. That is, Γ is the union

of faces F of $\text{Newt}(f)$ that are visible from the origin in the sense that for every $W \in F$, the intersection of $\text{Newt}(f)$ with the interval from the origin to W equals $\{W\}$. Let $\Sigma = \Sigma_f$ be the dual fan of $\text{Newt}(f)$. For each face F of $\text{Newt}(f)$, let σ_F be the cone of Σ dual to F .

5.2. Formula for the local motivic zeta function. We now recall the formula of Bultot and Nicaise for the local motivic zeta function of a nondegenerate polynomial f .

Consider a nonempty compact face K of Γ . Following [BN20], we associate two classes in $\mathcal{M}^{\hat{\mu}}$ to K . For $i \in \{0, 1\}$, let $Y_K(i)$ be the closed subscheme of $\text{Spec } \mathbb{k}[\text{span}(K) \cap \mathbb{Z}^n]$ cut out by $f_K = i$. When $i = 0$, we endow $Y_K(0)$ with the trivial $\hat{\mu}$ -action and obtain a class $[Y_K(0)] \in \mathcal{M}^{\hat{\mu}}$. We define a $\hat{\mu}$ -action on $Y_K(1)$ as follows. Let ρ_K be the lattice distance of K to the origin, and let $w = w_K := \rho_K \psi_K \in \text{Hom}(\text{span}(K) \cap \mathbb{Z}^n, \mathbb{Z})$. Then w determines a cocharacter $\text{Spec } \mathbb{k}[\mathbb{Z}] \rightarrow \text{Spec } \mathbb{k}[\text{span}(K) \cap \mathbb{Z}^n]$, which we can restrict via $\text{Spec } \mathbb{k}[T]/(T^\rho - 1) \rightarrow \text{Spec } \mathbb{k}[\mathbb{Z}]$ to determine a μ_ρ -action on $\text{Spec } \mathbb{k}[\text{span}(K) \cap \mathbb{Z}^n]$. This induces an action of μ_ρ on $Y_K(1)$. Explicitly, choose a basis for $\text{span}(K) \cap \mathbb{Z}^n$ and write $w = (w_1, \dots, w_r)$ and $f = \sum_{a \in \mathbb{Z}^r} \lambda_a x^a$. Then for each $a = (a_1, \dots, a_r)$ with $\lambda_a \neq 0$, $\sum_{i=1}^r a_i w_i$ is divisible by ρ , and the action is

$$\zeta \cdot (x_1, \dots, x_r) = (\zeta^{w_1} x_1, \dots, \zeta^{w_r} x_r).$$

This gives a class $[Y_K(1)]$ in $\mathcal{M}^{\hat{\mu}}$. When K is the empty compact face of Γ , $\text{span}(K) \cap \mathbb{Z}^n = \{0\}$, and we let $Y_K(0)$ be the point $\text{Spec } \mathbb{k}[\text{span}(K) \cap \mathbb{Z}^n]$ and let $Y_K(1) = \emptyset$. Then $[Y_K(0)] = 1$ and $[Y_K(1)] = 0$.

Remark 5.2.1. The above construction differs slightly from that in [BN20]. Explicitly, for $i \in \{0, 1\}$, [BN20] let $X_K(i)$ be the closed subscheme of $\text{Spec } \mathbb{k}[\mathbb{Z}^n]$ cut out by $f_K = i$. Consider $X_K(0)$ with the trivial $\hat{\mu}$ -action. Let w be any linear function in \mathbb{Z}^n that restricts to $\rho \psi_K$, and, as above, consider the corresponding μ_ρ -action on $X_K(1) \subset \text{Spec } \mathbb{k}[\mathbb{Z}^n]$. They consider the classes $[X_K(i)]$ in $\mathcal{M}^{\hat{\mu}}$. It follows from [BN20, Proposition 7.1.1] that $[X_K(i)] = [Y_K(i)](\mathbb{L} - 1)^{n-1-\dim K}$.

The following lemma will be important in the proof of Theorem 6.1.2.

Lemma 5.2.2. *Let $G \subset F$ be an inclusion of compact faces of Γ . Suppose there exists a vertex A of F such that $F = \text{Conv}\{G, A\}$ and $\text{span}(F) \cap \mathbb{Z}^n = \text{span}(G) \cap \mathbb{Z}^n + \mathbb{Z} \cdot A$. Then for $i \in \{0, 1\}$, $[Y_G(i)] + [Y_F(i)] = (\mathbb{L} - 1)^{\dim F} \in \mathcal{M}^{\hat{\mu}}$.*

Proof. Let $r = \dim F$. Let ρ_G and ρ_F be the smallest positive integers such that $w_G = \rho_G \psi_G$ and $w_F = \rho_F \psi_F$ lie in $\text{Hom}(\text{span}(G) \cap \mathbb{Z}^n, \mathbb{Z})$ and $\text{Hom}(\text{span}(F) \cap \mathbb{Z}^n, \mathbb{Z})$ respectively.

Then, we may choose coordinates such that $Y_F(i)$ is defined by $\{f_F(x_0, \dots, x_r) = i\}$ in $\text{Spec } \mathbb{k}[\text{span}(F) \cap \mathbb{Z}^n]$, $Y_G(i)$ is defined by $\{f_G(x_1, \dots, x_r) = i\}$ in $\text{Spec } \mathbb{k}[\text{span}(G) \cap \mathbb{Z}^n]$, and $f_F(x_0, \dots, x_r) = x_0 + f_G(x_1, \dots, x_r)$. Also, we may set $\rho = \rho_F = \rho_G$ and write $w_G = (w_1, \dots, w_r)$ and $w_F = (1, w_1, \dots, w_r)$. As above, w_G and w_F induce μ_ρ -actions on $\text{Spec } \mathbb{k}[\text{span}(G) \cap \mathbb{Z}^n]$ and $\text{Spec } \mathbb{k}[\text{span}(F) \cap \mathbb{Z}^n]$ respectively. Consider the μ_ρ -equivariant map

$$\phi: \text{Spec } \mathbb{k}[\text{span}(G) \cap \mathbb{Z}^n] \setminus Y_G(i) \rightarrow Y_F(i),$$

$$\phi(x_1, \dots, x_r) = (i - f_G(x_1, \dots, x_r), x_1, \dots, x_r).$$

Then ϕ is an isomorphism, with inverse $\phi^{-1}(x_0, x_1, \dots, x_r) = (x_1, \dots, x_r)$. By [BN20, Lemma 7.1.1], the class of any r -dimensional torus in $\mathcal{M}^{\hat{\mu}}$ is $(\mathbb{L} - 1)^r$, and the result follows. \square

Example 5.2.3. Let A be a primitive vertex of Γ . Then Lemma 5.2.2, with $F = \{A\}$ and $G = \emptyset$, implies that $[Y_A(0)] = 0$ and $[Y_A(1)] = 1$.

Example 5.2.4. Let F be a compact B_1 -face of Γ with nonempty base G . Then Lemma 5.2.2 implies that $[Y_G(0)] \frac{\mathbb{L}^{-1}T}{1-\mathbb{L}^{-1}T} + [Y_G(1)] + [Y_F(0)] \frac{\mathbb{L}^{-1}T}{1-\mathbb{L}^{-1}T} + [Y_F(1)] = \frac{(\mathbb{L}-1)^{\dim F}}{1-\mathbb{L}^{-1}T}$.

We now discuss two results on lattice point enumeration. The first result is standard. Let C be a nonzero rational polyhedral cone in $\mathbb{R}_{\geq 0}^n$ with rays spanned by primitive integer vectors u_1, \dots, u_r . Let $\text{Box}^+(C) = \{u \in \mathbb{N}^n : u = \sum_{i=1}^r \lambda_i u_i \text{ for some } 0 < \lambda_i \leq 1\}$. Then

$$(21) \quad \sum_{u \in C^\circ \cap \mathbb{N}^n} x^u = \frac{\sum_{u \in \text{Box}^+(C)} x^u}{\prod_{i=1}^r (1 - x^{u_i})} \in \mathbb{Z}[[x_1, \dots, x_n]].$$

We also need the following lemma.

Lemma 5.2.5. *Let C be a rational polyhedral cone of dimension d contained in $\mathbb{R}_{\geq 0}^n$, and let Y be a \mathbb{Z} -linear function that takes nonnegative values on C and is not identically zero on C . Let u_1, \dots, u_r be the primitive generators of the rays of C . Let $I = \{i \in [r] : \langle u_i, Y \rangle \neq 0\}$. Assume that $\langle u_j, \mathbf{1} \rangle = 1$ for $j \notin I$. Then*

$$(L-1)^{d-1} \sum_{u \in C^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, Y \rangle}$$

lies in the subring

$$\mathbb{Z}[L, L^{-1}, T] \left[\frac{1}{1 - L^{-\langle u_i, \mathbf{1} \rangle} T^{\langle u_i, Y \rangle}} \right]_{i \in I} \subset \mathbb{Z}[L][[L^{-1}, T]].$$

Proof. This is essentially [BN20, Lemma 5.1.1]; the point is that we may reduce to the case when C is simplicial and then apply (21). The $1/(1 - L^{-1})$ terms that arise from $j \notin I$ are cancelled by the $(L-1)^{d-1}$ factor. \square

Define a piecewise linear function N on Σ by

$$N(u) = \min\{\langle u, W \rangle : W \in \text{Newt}(f)\}.$$

Remark 5.2.6. If u is a primitive generator of a ray in the dual fan, corresponding to a facet F of $\text{Newt}(f)$, then $N(u)$ is the lattice distance of F to the origin. If $N(u) = 0$, then $u = e_i^*$ for some $1 \leq i \leq n$, and hence $\langle u, \mathbf{1} \rangle = 1$.

Lemma 5.2.7. [BN20, proof of Theorem 8.3.5] *Let u_1, \dots, u_r be the primitive generators of the rays of σ_K . The element*

$$(L-1)^{n-\dim K} \sum_{u \in \sigma_K^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)}$$

lies in the subring $\mathbb{Z}[L, L^{-1}, T] \left[\frac{1}{1 - L^{-\langle u_i, \mathbf{1} \rangle} T^{N(u_i)}} \right]_{\{i \in [r] : N(u_i) \neq 0\}}$ of $\mathbb{Z}[L][[L^{-1}, T]]$.

Proof. Observe that the restriction of N to σ_K is a nonnegative linear function. The result follows from Lemma 5.2.5 and Remark 5.2.6. \square

In [BN20], they define

$$(\mathbb{L}-1)^{n-\dim K} \sum_{u \in \sigma_K^\circ \cap \mathbb{N}^n} \mathbb{L}^{-\langle u, \mathbf{1} \rangle} T^{N(u)} \in \mathcal{M}^{\hat{\mu}}[[T]]$$

to be the image of the expression in Lemma 5.2.7 under the specialization map $\mathbb{Z}[L, L^{-1}][[T]] \rightarrow \mathcal{M}^{\hat{\mu}}[[T]]$ that sends L to \mathbb{L} .

Theorem 5.2.8. [BN20, Theorem 8.3.5] *Suppose f is nondegenerate. Then*

$$(22) \quad Z_{\text{mot}}(T) = \sum_K \left([Y_K(0)] \frac{\mathbb{L}^{-1} T}{1 - \mathbb{L}^{-1} T} + [Y_K(1)] \right) \left((\mathbb{L}-1)^{n-\dim K} \sum_{u \in \sigma_K^\circ \cap \mathbb{N}^n} \mathbb{L}^{-\langle u, \mathbf{1} \rangle} T^{N(u)} \right) \in \mathcal{M}^{\hat{\mu}}[[T]],$$

where the sum is over nonempty compact faces $K \in \Gamma$.

Remark 5.2.9. There is an extra factor of $(\mathbb{L} - 1)$ in (22) that does not appear in [BN20, Theorem 8.3.5] for consistency with our choice of normalization of the local motivic zeta function, cf. Remark 1.2.6.

5.3. The local formal zeta function. We now introduce the *local formal zeta function* of f , denoted $Z_{\text{for}}(T)$, which is a power series over a polynomial ring that specializes to $Z_{\text{mot}}(T)$. The key advantage of $Z_{\text{for}}(T)$ is that it lies in a power series ring over an integral domain, so it is easier to understand sets of candidate poles of $Z_{\text{for}}(T)$. Also, $Z_{\text{for}}(T)$ depends only on $\text{Newt}(f)$, as opposed to $Z_{\text{mot}}(T)$ which depends on f .

Let D be a ring containing $\mathbb{Z}[L, L^{-1}, T, \frac{1}{1-L^{-1}T}]$ as a subring. Let

$$R_D = D[Y_K : \emptyset \neq K \in \Gamma, K \text{ compact}] / (\mathcal{I}_1 + \mathcal{I}_2), \text{ where}$$

$$\mathcal{I}_1 = (Y_V - 1 : V \text{ primitive vertex of } \Gamma), \text{ and}$$

$$\mathcal{I}_2 = (Y_G + Y_F - \frac{(L-1)^{\dim F}}{1-L^{-1}T} : F \text{ compact } B_1\text{-face with nonempty base } G).$$

When $D = \mathbb{Z}[L, L^{-1}][[T]]$, we write $R := R_D$. It follows from Example 5.2.3 and Example 5.2.4 that we have a well-defined $\mathbb{Z}[[T]]$ -algebra homomorphism

$$\text{sp}: R \rightarrow \mathcal{M}^{\mu}[[T]], \text{ given by } \text{sp}(L) = \mathbb{L}, \text{ sp}(Y_K) = [Y_K(0)] \frac{\mathbb{L}^{-1}T}{1-\mathbb{L}^{-1}T} + [Y_K(1)].$$

Definition 5.3.1. *With the notation above,*

$$Z_{\text{for}}(T) := \sum_{\substack{\emptyset \neq K \in \Gamma \\ K \text{ compact}}} Y_K \left((L-1)^{n-\dim K} \sum_{u \in \sigma_K^{\circ} \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)} \right) \in R.$$

Above, the fact that the right-hand side lies in R follows from Lemma 5.2.7. By Theorem 5.2.8, $\text{sp}(Z_{\text{for}}(T)) = Z_{\text{mot}}(T)$.

Lemma 5.3.2. *The ring R_D is isomorphic to a polynomial ring over D . Moreover, if D is a subring of D' , then R_D is naturally a subring of $R_{D'}$.*

Proof. Consider the following change of variables. If K is a nonempty compact face of Γ , then let

$$Z_K := (-1)^{\dim K} \left(Y_K - \frac{(L-1)^{\dim F+1}}{1-L^{-1}T} \sum_{i=0}^{n-\dim F-1} (1-L)^i \right).$$

Then $R_D = D[Z_K : \emptyset \neq K \in \Gamma, K \text{ compact}] / (\mathcal{I}_1 + \mathcal{I}_2)$, where

$$\mathcal{I}_1 = (Z_V - 1 + \frac{L-1}{1-L^{-1}T} \sum_{i=0}^{n-1} (1-L)^i : V \text{ primitive vertex of } \Gamma),$$

$$\mathcal{I}_2 = (Z_G - Z_F : F \text{ compact } B_1\text{-face with nonempty base } G).$$

Consider the equivalence relation on nonempty compact faces in Γ generated by $G \sim F$ whenever F is a compact B_1 -face with nonempty base G . Then R is isomorphic to a polynomial ring over D with variables indexed by all equivalence classes that do not contain a primitive vertex of Γ . If D is a subring of D' , then $R_{D'}$ is a polynomial ring over D' in the same variables as above. It follows that the natural map $R_D \rightarrow R_{D'}$ is injective. \square

When $D = \mathbb{Z}[L][[L^{-1}, T]]$, we let $\tilde{R} := R_D$. By Lemma 5.3.2, R is a subring of \tilde{R} . In what follows, we will freely view $Z_{\text{for}}(T)$ as an element of \tilde{R} , in order to ensure relevant infinite sums in L^{-1} and T are well-defined.

We next define the notion of a set of candidate poles for the local formal zeta function. Let \mathcal{P} be a finite set of rational numbers containing -1 . Then \mathcal{P} is a *set of candidate poles* for some power series $Z(T) \in R$ if $Z(T)$ belongs to the subring R_D of R , where

$$D = \mathbb{Z}[L, L^{-1}, T] \left[\frac{1}{1 - L^a T^b} \right]_{(a,b) \in \mathbb{Z} \times \mathbb{Z}_{>0}, a/b \in \mathcal{P}}.$$

By Lemma 5.2.7, $\{\alpha \in \mathbb{Q} : \text{Contrib}(\alpha) \neq \emptyset\} \cup \{-1\}$ is a set of candidate poles for $Z_{\text{for}}(T)$.

Remark 5.3.3. Since $\text{sp}(Z_{\text{for}}(T)) = Z_{\text{mot}}(T)$, any set of candidate poles for $Z_{\text{for}}(T)$ is a set of candidate poles for $Z_{\text{mot}}(T)$.

Remark 5.3.4. Let \mathcal{P}_1 and \mathcal{P}_2 be sets of candidate poles for elements $Z_1(T)$ and $Z_2(T)$ in R respectively. It follows from the definition that $\mathcal{P}_1 \cup \mathcal{P}_2$ is a set of candidate poles for $Z_1(T) + Z_2(T)$.

The main benefit of working with candidate poles of $Z_{\text{for}}(T)$ is that they satisfy the following lemma.

Lemma 5.3.5. *Let \mathcal{P}_1 and \mathcal{P}_2 be sets of candidate poles for $Z(T) \in R$. Then $\mathcal{P}_1 \cap \mathcal{P}_2$ is a set of candidate poles for $Z(T)$.*

Proof. Let $D' = \mathbb{Z}[L, L^{-1}, T, \frac{1}{1-L^{-1}T}]$ and $R' = R_{D'}$. We can write $Z(T) = \frac{F_i(T)}{G_i(t)}$ for $i \in \{1, 2\}$ for some $F_i(T) \in R$ and some $G_i(t)$ a finite product of terms of the form $\{1 - L^a T^b : (a, b) \in \mathbb{Z} \times \mathbb{Z}_{>0}, a/b \in \mathcal{P}_i\}$. Suppose that $G_1(T) = (1 - L^c T^d)G'_1(T)$ for some $(c, d) \in \mathbb{Z} \times \mathbb{Z}_{>0}, c/d \in \mathcal{P}_1 \setminus \mathcal{P}_2$. By induction on $\deg(G_1(T))$, the result will follow if we can show that $F_1(T) = (1 - L^c T^d)F'_1(T)$ for some $F'_1(T) \in R'$.

Since the leading coefficient of $1 - L^c T^d$ is a unit in R' , we may apply the division algorithm to write $F_1(T) = (1 - L^c T^d)F'_1(T) + \tilde{F}_1(T)$ for some $F'_1(T), \tilde{F}_1(T) \in R'$, with $\deg \tilde{F}_1(T) < d$. The equality $F_1(T)G_2(T) = F_2(T)(1 - L^c T^d)G'_1(T)$ in R' implies that $1 - L^c T^d$ divides $\tilde{F}_1(T)G_2(T)$. Over an appropriate choice of ring R_D containing R as a subring, $1 - L^c T^d$ has roots $\{\exp(\frac{2\pi i j}{d})L^{-c/d} : 0 \leq j < d\}$. Similarly, $G_2(T)$ has roots contained in $\{\exp(\frac{2\pi i j}{b})L^{-a/b} : 0 \leq j < b, (a, b) \in \mathbb{Z} \times \mathbb{Z}_{>0}, a/b \in \mathcal{P}_2\}$. Since $c/d \notin \mathcal{P}_2$ and $\deg \tilde{F}_1(T) < d$, we deduce that $\tilde{F}_1(T) = 0$, and $Z(T) = \frac{F'_1(T)}{G'_1(T)}$, as desired. \square

Remark 5.3.6. By Lemma 5.3.5, if $Z(T) \in R$ admits a set of candidate poles, then there exists a minimal set of candidate poles. In particular, there exists a minimal set of candidate poles of $Z_{\text{for}}(T)$.

5.4. Simplifying the local formal zeta function. We now develop some tools to manipulate the local formal zeta function. Given a subset $C \subset \mathbb{R}_{\geq 0}^n$, we write

$$(23) \quad Z_{\text{for}}(T)|_C := \sum_{\substack{\emptyset \neq K \in \Gamma \\ K \text{ compact}}} Y_K \left((L-1)^{n-\dim K} \sum_{u \in \sigma_K^\circ \cap C \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)} \right) \in \tilde{R}.$$

We call $Z_{\text{for}}(T)|_C$ the *contribution* of C to $Z_{\text{for}}(T)$.

We will now prove a key technical tool we will use to manipulate $Z_{\text{for}}(T)$. Lemma 5.4.1 is analogous to [ELT22, Lemma 3.3].

Lemma 5.4.1. *Let F be a compact B_1 -face with nonempty base G and apex A in the direction e_ℓ^* . Let C' be a nonzero rational polyhedral cone with $(C')^\circ \subset \sigma_F^\circ$, and let $C \subset \sigma_G$ be the convex hull of C' and $\mathbb{R}_{\geq 0}e_\ell^*$. Then*

$$Z_{\text{for}}(T)|_{(C^\circ \cup (C')^\circ)} = (L-1)^n \left(\sum_{u \in (C^\circ \cup (C')^\circ) \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A \rangle} \right) \in \tilde{R}.$$

Proof. A simplicial refinement of C' induces a simplicial refinement of C , so we may reduce to the case that C' is simplicial. Let u_1, \dots, u_r be the primitive integer vectors spanning the rays of C' . Then $u_0 = e_\ell^*, u_1, \dots, u_r$ are the primitive integer vectors spanning the rays of C . With the notation of (21), $\text{Box}^+(C') = \{u \in \mathbb{N}^n : u = \sum_{i=1}^r \lambda_i u_i \text{ for some } 0 < \lambda_i \leq 1\}$ and $\text{Box}^+(C) = \{u \in \mathbb{N}^n : u = \sum_{i=0}^r \lambda_i u_i \text{ for some } 0 < \lambda_i \leq 1\}$. We claim that $\text{Box}^+(C) = \{u + e_\ell^* : u \in \text{Box}^+(C')\}$.

Clearly, $\{u + e_\ell^* : u \in \text{Box}^+(C')\} \subset \text{Box}^+(C)$. Conversely, consider an element $u' = \sum_{i=0}^r \lambda_i u_i \in \text{Box}^+(C)$ for some $0 < \lambda_i \leq 1$. Let X be a point in G . For $1 \leq i \leq r$, since $u_i \in \sigma_F$, we have $\langle u_i, A - X \rangle = 0$. Also, $N(e_\ell^*) = \langle e_\ell^*, X \rangle = 0$ and $\langle e_\ell^*, A \rangle = 1$. We compute:

$$\langle u', A - X \rangle = \lambda_0 + \sum_{i=1}^r \lambda_i \langle u_i, A - X \rangle = \lambda_0 \in \mathbb{Z}.$$

Hence $\lambda_0 \in \mathbb{Z} \cap (0, 1] = \{1\}$. Then $u' = \sum_{i=1}^r \lambda_i u_i + e_\ell^* \in \{u + e_\ell^* : u \in \text{Box}^+(C')\}$, which proves the claim.

Observe that if $u \in \text{Box}^+(C')$, then $N(u + e_\ell^*) = N(u) = \langle u, A \rangle$ and $\langle u + e_\ell^*, \mathbf{1} \rangle = \langle u, \mathbf{1} \rangle + 1$. Also, $N(u_i) = \langle u_i, A \rangle$ for $1 \leq i \leq r$. Then using (21) and the relations in \mathcal{I}_2 , we compute the left hand side of the equality in Lemma 5.4.1:

$$\begin{aligned} & (L-1) \left(Y_F (L-1)^{n-1-\dim F} + Y_G (L-1)^{n-1-\dim G} \frac{L^{-1}}{1-L^{-1}} \right) \frac{\sum_{u \in \text{Box}^+(C')} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)}}{\prod_{i=1}^r (1 - L^{-\langle u_i, \mathbf{1} \rangle} T^{N(u_i)})} \\ &= \frac{(L-1)^n \sum_{u \in \text{Box}^+(C')} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A \rangle}}{1 - L^{-1} T \prod_{i=1}^r (1 - L^{-\langle u_i, \mathbf{1} \rangle} T^{\langle u_i, A \rangle})}. \end{aligned}$$

Similarly, using (21), we compute the right-hand side of the equality in Lemma 5.4.1:

$$(L-1)^n \left(1 + \frac{L^{-1} T}{1 - L^{-1} T} \right) \frac{\sum_{u \in \text{Box}^+(C')} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A \rangle}}{\prod_{i=1}^r (1 - L^{-\langle u_i, \mathbf{1} \rangle} T^{\langle u_i, A \rangle})}.$$

The result follows. \square

Remark 5.4.2. We note that a version of Lemma 5.4.1 holds when G is empty, i.e., $F = \{A\}$ is a vertex with some coordinate 1. Then A is a primitive vertex, so the relations in \mathcal{I}_1 imply that for $C \subset \sigma_A^\circ$,

$$Z_{\text{for}}(T)|_C = (L-1)^n \left(\sum_{u \in C \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A \rangle} \right) \in \tilde{\mathcal{R}}.$$

6. FAKE POLES FOR THE LOCAL FORMAL ZETA FUNCTION

6.1. Overview. In this section, we prove Theorem 1.4.7. We first introduce some notation before stating a strengthening of Theorem 1.4.7 and outlining its proof.

We let $\text{Vert}(F)$ denote the set of vertices of F . Given a face F of Γ , recall that C_F is the closure of the cone over F , with distinguished generators $\text{Gen}(C_F)$. Then $\text{span}(F) = \text{span}(C_F)$ and $\text{Gen}(C_F) = \text{Vert}(F) \cup \text{Unb}(C_F)$. Given an inclusion of faces $M \subset F$, let $\text{Gen}(C_F \setminus C_M) = \text{Gen}(C_F) \setminus \text{Gen}(C_M)$. Recall that a face F of $\text{Newt}(f)$ is B_1 if it has an apex A with base direction e_ℓ^* , and $\langle e_\ell^*, A \rangle = 1$. Given a B_1 -face F , let \mathcal{A}_F be the set of all choices of such an apex A .

Definition 6.1.1. We say that the Newton polyhedron $\text{Newt}(f)$ is α -simplicial if for any minimal element M in $\text{Contrib}(\alpha)$ and any face $F \supset M$, $\dim C_F = \dim C_M + |\text{Gen}(C_F \setminus C_M)|$. Equivalently, the images of the elements of $\text{Gen}(C_F \setminus C_M)$ are linearly independent in $\mathbb{R}^n / \text{span}(C_M)$.

For example, if $\text{Newt}(f)$ is simplicial then it is α -simplicial. If all minimal elements in $\text{Contrib}(\alpha)$ are facets, then $\text{Newt}(f)$ is α -simplicial. One key property of α -simplicial Newton polyhedra is that every face of $\text{Contrib}(\alpha)$ contains a unique minimal face of $\text{Contrib}(\alpha)$ (Lemma 6.2.4). We now state our main theorem.

Theorem 6.1.2. *Suppose f is nondegenerate. Let*

$$\mathcal{P} = \{\alpha \in \mathbb{Q} : \text{Contrib}(\alpha) \neq \emptyset\} \cup \{-1\}, \text{ and}$$

$$\mathcal{P}' = \{\alpha \in \mathcal{P} : \alpha \notin \mathbb{Z}, \text{ every face in } \text{Contrib}(\alpha) \text{ is } UB_1 \text{ and } \text{Newt}(f) \text{ is } \alpha\text{-simplicial}\}.$$

Then $\mathcal{P} \setminus \mathcal{P}'$ is a set of candidate poles for $Z_{\text{mot}}(T)$.

Our strategy to prove Theorem 6.1.2 involves repeatedly applying Lemma 5.4.1, which will require us to choose apices and base directions for various B_1 -faces. We will require the following compatibility condition.

Definition 6.1.3. *A locally unique labeling of $\text{Contrib}(\alpha)$ is a choice of an apex A_F and a base direction e_F^* for each $F \in \text{Contrib}(\alpha)$ such that:*

$$(*) \text{ whenever } F \subset F' \text{ and } A_F = A_{F'}, \text{ we have } e_F^* = e_{F'}^*.$$

If every face of $\text{Contrib}(\alpha)$ is UB_1 , then $\text{Contrib}(\alpha)$ has a locally unique labeling (Lemma 6.2.1).

We now summarize the rest of the proof of Theorem 6.1.2. We first establish some notation and basic results in Section 6.2. Then, using Lemma 5.3.5, we reduce to showing that for each candidate pole $\alpha \notin \mathbb{Z}$ that is contributed only by UB_1 -faces and such that $\text{Newt}(f)$ is α -simplicial, there is a set of candidate poles for $Z_{\text{for}}(T)$ not containing α . Fix such an α .

Because $\text{Newt}(f)$ is α -simplicial, every face of $\text{Contrib}(\alpha)$ contains a unique minimal face M of $\text{Contrib}(\alpha)$. In Section 6.3, we develop the tools to argue that we can consider each minimal face M separately. We construct a neighborhood $N_{M, \leq \delta}$ of σ_M . In Lemma 6.3.12, we show that if $Z_{\text{for}}(T)|_{N_{M, \leq \delta}^\circ}$ admits a set of candidate poles not containing α for each minimal face M , then the theorem follows.

To analyze $Z_{\text{for}}(T)|_{N_{M, \leq \delta}^\circ}$, our strategy is to construct a complete fan $\Sigma_{\mathcal{Z}}$ where each maximal cone is labeled by a face containing M , which necessarily lies in $\text{Contrib}(\alpha)$. We construct $\Sigma_{\mathcal{Z}}$ as the normal fan of a polytope where each vertex is labeled by a face containing M . This polytope is determined by an α -compatible pair (see Definition 6.4.11).

Because we have fixed a locally unique labeling, we may associate to each maximal cone of $\Sigma_{\mathcal{Z}}$, which corresponds to a face F containing M , a pair (A_F, e_F^*) consisting of an apex and a base direction. We then associate a face containing M and a pair (A, e_ℓ^*) to all cones in the fan $\Sigma_{\mathcal{Z}} \cap N_{M, \leq \delta}$. See Definition 6.4.3. Consider a nonzero cone C in the fan $\Sigma_{\mathcal{Z}} \cap N_{M, \leq \delta}$ and an associated pair (A, e_ℓ^*) . We arrange that for any face F containing M , if the dual cone of F intersects C , then F is a B_1 -face with apex A and base direction e_ℓ^* . By repeatedly applying Lemma 5.4.1, we show that the contribution of C° to $Z_{\text{for}}(T)$ is equal to $(L-1)^n \sum_{u \in C^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{(u, A)}$. In order to do this, we need to show that C is locally defined by elements “orthogonal to e_ℓ^* ”, see Lemma 6.4.8. Using an additional genericity condition (see Definition 6.4.11), we deduce that the contribution $Z_{\text{for}}(T)|_{C^\circ}$ admits a set of candidate poles not containing α . Using this strategy, in Section 6.4 we prove that the existence of an α -compatible pair implies Theorem 6.1.2.

In Section 6.5, we show that the existence of an α -compatible pair is implied by the existence of a *restricted, weakly α -compatible* pair (Definition 6.4.1, Definition 6.5.1). More precisely, we show that each restricted, weakly α -compatible pair can be “deformed” into an α -compatible pair. In Section 6.6, we give an explicit construction of a restricted, weakly α -compatible pair, which is where we use the locally unique condition. Figure 2 shows an example of a fan corresponding to a restricted, weakly α -compatible pair, and a corresponding deformation.




FIGURE 2. A complete fan with maximal cones indexed by faces containing M and a corresponding deformation. We show the intersection of $\text{span}(\sigma_M)$ with an affine hyperplane. The cone σ_M is shown in black and grey, while the codimension 1 cones of the complete fan appear in red, with their maximal cones labeled in red.

6.2. Combinatorial preliminaries. In this section, we prepare for the constructions that will take the rest of the section. Recall from Definition 1.4.3 that a face G of $\text{Newt}(f)$ is UB_1 if there exists an apex A in G with a unique choice of base direction e_ℓ^* , and $\langle e_\ell^*, A \rangle = 1$.

Lemma 6.2.1. *If every face in $\text{Contrib}(\alpha)$ is UB_1 , then $\text{Contrib}(\alpha)$ admits a locally unique labeling.*

Proof. Suppose that every face F in $\text{Contrib}(\alpha)$ is UB_1 . Then it has an apex A_F with a unique choice of base direction e_F^* . We may choose one such apex, and label F by (A_F, e_F^*) to get a locally unique labeling. Indeed, if $F \subset F'$ and $A_F = A_{F'}$, then $e_{F'}^*$ is a base direction for F with apex A_F . We deduce that $e_F^* = e_{F'}^*$. \square

Given an element $V \in \Gamma$, we say that $u \in \mathbb{R}^n$ is *critical with respect to (α, V)* if $\alpha \langle u, V \rangle + \langle u, \mathbf{1} \rangle = 0$. Recall that we have a piecewise linear function $N(u) = \min\{\langle u, W \rangle : W \in \text{Newt}(f)\}$. We say that $u \in \mathbb{R}_{\geq 0}^n$ is *critical with respect to α* if $\alpha N(u) + \langle u, \mathbf{1} \rangle = 0$. Equivalently, for some/any $G \in \Gamma$ such that $u \in \sigma_G$, and some/any element $V \in G$, u is critical with respect to (α, V) . A set is critical with respect to α (or (α, V)) if every element of the set is critical with respect to α (or (α, V)).

Remark 6.2.2. Unless $V = -(1/\alpha)\mathbf{1}$, then the points $u \in \mathbb{R}^n$ critical with respect to (α, V) form a hyperplane. A special case of this observation is [ELT22, Lemma 3.4].

We conclude this section with three combinatorial observations.

Lemma 6.2.3. *Let M be an element of $\text{Contrib}(\alpha)$, and assume that every face of $\text{Contrib}(\alpha)$ is UB_1 . Assume that $\alpha \neq -|\text{Vert}(M)| \in \mathbb{Z}$. Then there exists a vertex V_M of M such that M is not a B_1 -face with apex V_M .*

Proof. Suppose every vertex of M is an apex. Since $\mathbf{1} \in \text{span}(M)$, it follows that $\mathbf{1} - \sum_{V \in \text{Vert}(M)} V$ is a linear combination of the unbounded directions of M , and hence $\alpha = -\psi_M(\mathbf{1}) = -|\text{Vert}(M)|$, a contradiction. \square

Lemma 6.2.4. *Assume that $\text{Newt}(f)$ is α -simplicial. If M_1, M_2 are distinct minimal elements in $\text{Contrib}(\alpha)$, then $\sigma_{M_1} \cap \sigma_{M_2} = \{0\}$.*

Proof. We argue by contradiction. Suppose $\sigma_{M_1} \cap \sigma_{M_2} \neq \{0\}$. Then there exists a facet F in $\partial \text{Newt}(f)$ such that $\sigma_F \subset \sigma_{M_1} \cap \sigma_{M_2}$. Note that F is interior and hence $F \in \Gamma$. Equivalently, M_1, M_2 are common faces of F . In particular, $M_1 \cap M_2$ is a (possibly empty) face of F , and $C_{M_1 \cap M_2} = C_{M_1} \cap C_{M_2}$. Let $\text{Gen}(C_{M_1} \setminus C_{M_1 \cap M_2}) = \{W_1, \dots, W_s\}$ and $\text{Gen}(C_{M_2} \setminus C_{M_1 \cap M_2}) = \{W'_1, \dots, W'_s\}$. By Definition 6.1.1 applied to $M_2 \subset$

F , $\text{Gen}(C_{M_1} \setminus C_{M_1 \cap M_2})$ is linearly independent in $\mathbb{R}^n / \text{span}(C_{M_2})$. By Definition 6.1.1 applied to $M_1 \subset F$, $\text{Gen}(C_{M_2} \setminus C_{M_1 \cap M_2})$ is linearly independent in $\mathbb{R}^n / \text{span}(C_{M_1})$. We claim that $W_1, \dots, W_s, W'_1, \dots, W'_{s'}$ are linearly independent in $\mathbb{R}^n / \text{span}(C_{M_1 \cap M_2})$. Indeed, if $\sum_i a_i W_i + \sum_j b_j W'_j = 0$ in $\mathbb{R}^n / \text{span}(C_{M_1 \cap M_2})$, then the corresponding equation in $\mathbb{R}^n / \text{span}(C_{M_1})$ implies that $b_j = 0$ for all j , and the corresponding equation in $\mathbb{R}^n / \text{span}(C_{M_2})$ implies that $a_i = 0$ for all i .

By assumption, $\mathbf{1} = a_1 W_1 + \dots + a_s W_s \in \mathbb{R}^n / \text{span}(C_{M_1 \cap M_2})$ for some $a_1, \dots, a_s \in \mathbb{R}$, and $\mathbf{1} = a'_1 W'_1 + \dots + a'_{s'} W'_{s'} \in \mathbb{R}^n / \text{span}(C_{M_1 \cap M_2})$ for some $a'_1, \dots, a'_{s'} \in \mathbb{R}$. Subtracting one equation from the other, and using the linear independence of $W_1, \dots, W_s, W'_1, \dots, W'_{s'}$ in $\mathbb{R}^n / \text{span}(C_{M_1 \cap M_2})$, we deduce that $a_i = a'_j = 0$ for all i, j . Hence $\mathbf{1} \in \text{span}(C_{M_1 \cap M_2})$, and $M_1 \cap M_2 \in \text{Contrib}(\alpha)$. This contradicts the minimality of M_1 and M_2 . \square

Remark 6.2.5. If e_ℓ^* is a base direction of F with apex A , then $e_\ell \notin \text{Unb}(C_F)$. Indeed, if $e_j \in \text{Unb}(C_F)$, then it follows from Definition 1.4.2 that $\langle e_\ell^*, e_j \rangle = 0$.

Assumptions and notation. For the remainder, we will assume that $\alpha \notin \mathbb{Z}$, $\text{Newt}(f)$ is α -simplicial and all faces of $\text{Contrib}(\alpha)$ are UB_1 . Let M be a minimal element of $\text{Contrib}(\alpha)$. Recall that we have chosen a locally unique labeling (A_F, e_F^*) of $\text{Contrib}(\alpha)$. By Lemma 6.2.3, we may fix a vertex V_M of M which is not an apex, and hence satisfies

$$(24) \quad \langle e_F^*, V_M \rangle = 0 \text{ for all } F \supset M.$$

We also fix a point W_M in the relative interior of M . Given a nonzero point W and $\epsilon \in \mathbb{R}$, let $H_{W, \epsilon} = \{u : \langle u, W \rangle = \epsilon\}$, and consider the associated half-spaces $H_{W, \geq \epsilon}, H_{W, > \epsilon}, H_{W, \leq \epsilon}, H_{W, < \epsilon}$. We let $H_W := H_{W, 0}$. Let $S' = \{u \in \mathbb{R}^n : \langle u, \mathbf{1} \rangle = 1\}$, and let $S = \text{Conv}\{e_1^*, \dots, e_n^*\} \subset S'$ be the standard $(n-1)$ -dimensional simplex.

6.3. Covering the critical locus. The goal of this section is to build small neighborhoods covering the locus of $u \in \mathbb{R}_{\geq 0}^n$ that is critical with respect to α . This will allow us to concentrate our attention on a single minimal face in $\text{Contrib}(\alpha)$.

Definition 6.3.1. Let M be a minimal element of $\text{Contrib}(\alpha)$ and let $\delta \in \mathbb{Q}_{\geq 0}$. We define $N_{M, \leq \delta}$ to be the cone over $\{u \in S : \langle u, W_M \rangle - N(u) \leq \delta\}$.

Similarly, we let $N_{M, < \delta}$, $N_{M, \delta}$ and $N_{M, \geq \delta}$ be the cones over $\{u \in S : \langle u, W_M \rangle - N(u) < \delta\}$, $\{u \in S : \langle u, W_M \rangle - N(u) = \delta\}$ and $\{u \in S : \langle u, W_M \rangle - N(u) \geq \delta\}$ respectively.

Here $\langle u, W_M \rangle - N(u)$ is a nonnegative function on $\mathbb{R}_{\geq 0}^n$ that is piecewise linear with respect to Σ . Because $N(u)$ is the support function of a polyhedron and hence convex, $N_{M, \leq \delta}$ is convex. It follows that $N_{M, \leq \delta}$ is a rational polyhedral cone of dimension n . Note that σ_M is the cone over $\{u \in S : \langle u, W_M \rangle - N(u) = 0\}$, and hence $\sigma_M \subset N_{M, \leq \delta}$. We can equivalently write $N_{M, \leq \delta} = \{u \in \mathbb{R}_{\geq 0}^n : \langle u, W_M \rangle - N(u) \leq \delta \langle u, \mathbf{1} \rangle\}$. Also, $N_{M, \leq \delta}^\circ = N_{M, < \delta} \cap \mathbb{R}_{> 0}^n$.

Lemma 6.3.2. Let $C \subset \mathbb{R}_{\geq 0}^n$ be a closed cone such that $C \cap \sigma_M = \{0\}$. Then $C \cap N_{M, \leq \delta} = \{0\}$ for δ sufficiently small.

Proof. We may assume that $C \cap S \neq \emptyset$. Since $C \cap S$ is compact, we may consider the minimal element $b > 0$ of $\{\langle u, W_M \rangle - N(u) : u \in C \cap S\}$. Then $C \cap N_{M, \leq \delta} \cap S = \emptyset$ for $\delta < b$. \square

Lemma 6.3.3. Let K be a face of $\partial \text{Newt}(f)$ and suppose that $K \notin \Gamma$. Let M be a minimal element in $\text{Contrib}(\alpha)$. Then $\sigma_K \cap N_{M, \leq \delta} = \{0\}$ for δ sufficiently small.

Proof. By Lemma 6.3.2, it is enough to show that $\sigma_K \cap \sigma_M = \{0\}$. Suppose that $\sigma_K \cap \sigma_M \neq \{0\}$. Then $\sigma_K \cap \sigma_M = \sigma_{K'}$ for some face K' of $\partial \text{Newt}(f)$ containing both K and M . Since $M \subset K'$, $K' \in \text{Contrib}(\alpha)$. Since $K \subset K' \in \Gamma$, we deduce that $K \in \Gamma$, a contradiction. \square

The following lemma is immediate from Lemma 6.2.4 and Lemma 6.3.2.

Lemma 6.3.4. *If M_1, M_2 are distinct minimal elements in $\text{Contrib}(\alpha)$, then $N_{M_1, \leq \delta} \cap N_{M_2, \leq \delta} = \{0\}$ for δ sufficiently small.*

Lemma 6.3.5. *For δ sufficiently small, if $u \in N_{M, \leq \delta} \setminus \{0\}$, then $N(u) > 0$.*

Proof. After scaling, we may assume that $u \in N_{M, \leq \delta} \cap S$. Since S is compact, we may consider $b = \min\{\langle u, W_M \rangle : u \in S\}$. Since M is interior, $W_M \notin \partial \mathbb{R}_{\geq 0}^n$, and hence $b > 0$. For $\delta < b$ and $u \in N_{M, \leq \delta} \cap S$, $N(u) = \langle u, W_M \rangle - (\langle u, W_M \rangle - N(u)) \geq \langle u, W_M \rangle - \delta \geq b - \delta > 0$. \square

Let Σ' be a fan supported on $\mathbb{R}_{\geq 0}^n$ that refines Σ . We may consider the fan $\Sigma' \cap N_{M, \leq \delta}$ supported on $N_{M, \leq \delta}$ given by all cones of the form $\{C \cap C' : C \in \Sigma', C' \text{ is a face of } N_{M, \leq \delta}\}$. The lemma below gives a more explicit description.

Lemma 6.3.6. *Let Σ' be a fan supported on $\mathbb{R}_{\geq 0}^n$ that refines Σ . Let $C \in \Sigma'$, and fix δ sufficiently small. Then*

- (1) *If $C \cap \sigma_M = \{0\}$, then $C \cap N_{M, \leq \delta} = \{0\}$.*
- (2) *If $C \subset \sigma_M$, then C is a cone in $\Sigma' \cap N_{M, \leq \delta}$.*
- (3) *If $C \cap \sigma_M \neq \{0\}$ and $C \not\subset \sigma_M$, then $\dim(C \cap N_{M, \leq \delta}) = \dim C$. Moreover, $\dim(C \cap N_{M, \delta}) = \dim C - 1$, and $C \cap N_{M, \delta}$ is the only proper face of $C \cap N_{M, \leq \delta}$ that is not contained in a proper face of C .*

Proof. If $C \cap \sigma_M = \{0\}$, then $C \cap N_{M, \leq \delta} = \{0\}$ by Lemma 6.3.2. If $C \subset \sigma_M$, then since $\sigma_M \subset N_{M, \leq \delta}$, $C \cap N_{M, \leq \delta} = C$. This establishes the first two properties. Assume that $C \cap \sigma_M \neq \{0\}$ and $C \not\subset \sigma_M$. Since Σ' refines Σ , $C \subset \sigma_K$ for some face K of $\partial \text{Newt}(f)$. Fix a vertex V of K , and let $P = C \cap S$. Then $P \cap N_{M, \leq \delta} = P \cap \mathbb{H}_{W_M - V, \leq \delta}$. For any $\delta > 0$, the relative interior of P intersects both $\mathbb{H}_{W_M - V, > 0}$ and $\mathbb{H}_{W_M - V, < \delta}$ because $P \not\subset \sigma_M$ and $P \cap \sigma_M \neq \emptyset$. It follows that for δ sufficiently small, $\mathbb{H}_{W_M - V, \delta}$ intersects the relative interior of P . We deduce that $P \cap N_{M, \leq \delta}$ has dimension $\dim P$, and that the only proper face of $P \cap N_{M, \leq \delta}$ that is not contained in a proper face of P is $P \cap \mathbb{H}_{W_M - V, \delta}$, which has dimension $\dim P - 1$. This establishes the result. \square

The following two remarks are corollaries of the Lemma 6.3.6 and its proof.

Remark 6.3.7. Let C be a rational polyhedral cone such that $C \subset \sigma_K$ for some face K of $\partial \text{Newt}(f)$. Then for δ sufficiently small, $C^\circ \cap N_{M, < \delta} = (C \cap N_{M, \leq \delta})^\circ$, and $C^\circ \cap N_{M, \geq \delta} = (C \cap N_{M, \geq \delta})^\circ \cup (C \cap N_{M, \delta})^\circ$.

Remark 6.3.8. Let K be a nonempty face of $\partial \text{Newt}(f)$, and let $\tilde{\sigma}_K = \sigma_K \cap (\cap_i N_{M_i, \geq \delta})$. Assume that δ is chosen sufficiently small. If $K \in \text{Contrib}(\alpha)$, then $M_i \subset K$ for some $1 \leq i \leq r$, and hence $\sigma_K \subset N_{M_i, < \delta}$, and $\tilde{\sigma}_K = \emptyset$. Assume that $K \notin \text{Contrib}(\alpha)$. By Lemma 6.3.4 and Lemma 6.3.6, $\tilde{\sigma}_K$ is a rational polyhedral cone of dimension $\dim \sigma_K$, and $\sigma_K^\circ \cap (\cap_i N_{M_i, \geq \delta}) = \tilde{\sigma}_K^\circ \cup (\cup_i (\tilde{\sigma}_K \cap N_{M_i, \delta})^\circ)$.

Lemma 6.3.9. *Let Σ' be a fan supported on $\mathbb{R}_{\geq 0}^n$ that refines Σ . Let γ be a ray of the fan $\Sigma' \cap N_{M, \leq \delta}$ such that $\gamma \not\subset \sigma_M$, for some δ chosen sufficiently small. Let C be the smallest cone in Σ' containing γ . The following properties hold:*

- (1) *$C \cap \sigma_M \neq \{0\}$,*
- (2) *the smallest cone in Σ containing C has the form σ_K for some K in Γ ,*
- (3) *if γ is critical with respect to (α, V) , for some vertex V of either K or M , then C is critical with respect to (α, V) .*

Proof. It follows from Lemma 6.3.6 that $\gamma = C \cap N_{M, \delta}$, $C \cap \sigma_M \neq \{0\}$, and $\dim(C \cap N_{M, \leq \delta}) = \dim C = 2$. This establishes (1). Lemma 6.3.3 implies (2). Since $\sigma_K \cap \sigma_M \neq \{0\}$, $\sigma_K \cap \sigma_M = \sigma_{K'}$ for some K' in Γ containing both M and K . Fix any vertex V of K' . Since $M \in \text{Contrib}(\alpha)$, we have $\sigma_{K'} \subset \mathbb{H}_{\alpha V + 1}$.

Let $\gamma' \neq \gamma$ be the other ray spanning $C \cap N_{M, \leq \delta}$. Then $\gamma' \subset \sigma_{K'} \subset H_{\alpha V+1}$. Hence, if $\gamma \subset H_{\alpha V+1}$, then $\text{span}(C) = \text{span}(C \cap N_{M, \leq \delta}) = \mathbb{R}\gamma + \mathbb{R}\gamma' \subset H_{\alpha V+1}$, and $C \subset H_{\alpha V+1}$. This establishes (3). \square

Lemma 6.3.10. *Let γ be a ray of the fan $\Sigma \cap N_{M, \leq \delta}$ such that $\gamma \not\subset \sigma_M$, for some δ chosen sufficiently small. Then γ is not critical with respect to α .*

Proof. By Lemma 6.3.9, there exists $K \in \Gamma$ such that σ_K is the smallest cone in Σ containing γ . Suppose that γ is critical with respect to α . Then γ is critical with respect to (α, V) for any vertex V of K . By Lemma 6.3.9, σ_K is critical with respect to (α, V) , and hence $K \in \text{Contrib}(\alpha)$. Since $\sigma_K \cap N_{M, \leq \delta} \neq \{0\}$, Lemma 6.3.4 implies that $M \subset K$, and hence $\gamma \subset \sigma_M$, a contradiction. \square

Lemma 6.3.11. *Let M_1, \dots, M_r be the minimal elements in $\text{Contrib}(\alpha)$, and let K be a nonempty face of $\partial \text{Newt}(f)$. Then for δ sufficiently small, $(L-1)^{n-\dim K} \sum_{u \in \sigma_K^\circ \cap (\cap_i N_{M_i, \geq \delta}) \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)}$ lies in R and admits a set of candidate poles not containing α .*

Proof. Let $\tilde{\sigma}_K = \sigma_K \cap (\cap_i N_{M_i, \geq \delta})$. By Remark 6.3.8, if $K \in \text{Contrib}(\alpha)$, then $\tilde{\sigma}_K = \emptyset$. Assume that $K \notin \text{Contrib}(\alpha)$. By Remark 6.3.8, $\tilde{\sigma}_K$ is a rational polyhedral cone of dimension $\dim \sigma_K$, and $\sigma_K^\circ \cap (\cap_i N_{M_i, \geq \delta}) = \tilde{\sigma}_K^\circ \cup (\cup_i (\tilde{\sigma}_K \cap N_{M_i, \delta})^\circ)$. By Lemma 6.3.2, the rays of $\tilde{\sigma}_K$ are the union of the rays of σ_K that are not critical with respect to α , and the rays of $\sigma_K \cap N_{M_i, \leq \delta}$ that do not lie in σ_{M_i} for $1 \leq i \leq r$. By Lemma 6.3.10, none of the rays of $\tilde{\sigma}_K$ are critical with respect to α . By Remark 5.2.6 and Lemma 6.3.5, if u is a primitive generator of a ray of $\tilde{\sigma}_K$ and $N(u) = 0$, then $u = e_i^*$ for some $1 \leq i \leq n$, and hence $\langle u, \mathbf{1} \rangle = 1$. Since the restriction of N to $\tilde{\sigma}_K \subset \sigma_K$ is linear, Lemma 5.2.5 implies that the following elements of \tilde{R} lie in R and admit sets of candidate poles not containing α :

$$(L-1)^{n-\dim K} \sum_{u \in \tilde{\sigma}_K^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)}, \text{ and } (L-1)^{n-\dim K} \sum_{u \in (\tilde{\sigma}_K \cap N_{M_i, \delta})^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)}$$

for $1 \leq i \leq r$. The result now follows from Remark 5.3.4. \square

The lemma below follows immediately from Definition 5.3.1 and Lemma 6.3.11 and will allow us to reduce our study of $Z_{\text{for}}(T)$ to the study of $Z_{\text{for}}(T)|_{N_{M, < \delta}}$, when M is a minimal element of $\text{Contrib}(\alpha)$.

Lemma 6.3.12. *Let M_1, \dots, M_r be the minimal elements in $\text{Contrib}(\alpha)$. Then for δ sufficiently small, $Z_{\text{for}}(T)|_{\cap_i N_{M_i, \geq \delta}}$ lies in R and admits a set of candidate poles not containing α .*

6.4. Establishing fake poles using α -compatible sets. The goal of this section is to show that the existence of a fan with certain properties implies that there is a set of candidate poles for the local formal zeta function not containing α .

From this point on, we fix a minimal element M in $\text{Contrib}(\alpha)$. Let $\text{Contrib}(\alpha)_M := \{F \in \text{Contrib}(\alpha) : F \supset M\}$. Fix a nonempty finite set \mathcal{S} . Given a finite collection $\mathcal{Z} = (Z_s)_{s \in \mathcal{S}}$ of elements in \mathbb{Q}^n indexed by \mathcal{S} , we let $Q_{\mathcal{Z}}$ denote the convex hull of \mathcal{Z} , and let $\Sigma_{\mathcal{Z}}$ be the corresponding (rational) dual fan supported on \mathbb{R}^n . Given a nonempty face J of $Q_{\mathcal{Z}}$, we write τ_J for the corresponding cone in $\Sigma_{\mathcal{Z}}$. Given an element $s \in \mathcal{S}$, we write J_{Z_s} for the smallest face of $Q_{\mathcal{Z}}$ containing Z_s , and we write $\tau_{Z_s} := \tau_{J_{Z_s}}$. Explicitly, $\tau_{Z_s} := \{u \in \mathbb{R}^n : \langle u, Z_s \rangle \leq \langle u, Z_{s'} \rangle, \text{ for all } s' \in \mathcal{S}\}$. Observe that $\tau_J = \cap_{Z_s \in J} \tau_{Z_s}$. Also, note that we do not require the elements of \mathcal{Z} to be distinct, and that τ_{Z_s} may equal $\tau_{Z_{s'}}$, even if $s \neq s'$.

Before defining α -compatible pairs (Definition 6.4.11), we introduce a weaker notion, which satisfies all of the properties of α -compatible pairs with the exception of a genericity condition.

Definition 6.4.1. *Consider a pair $(\mathcal{Z}, \mathcal{F})$, where $\mathcal{Z} = (Z_s)_{s \in \mathcal{S}}$ and $\mathcal{F} = (F_s)_{s \in \mathcal{S}}$ are collections of elements of \mathbb{Q}^n and $\text{Contrib}(\alpha)_M$ respectively. Then $(\mathcal{Z}, \mathcal{F})$ is weakly α -compatible if it satisfies the following property:*

Whenever $\sigma_K^\circ \cap \tau_{Z_s} \cap \tau_{Z_{s'}} \neq \emptyset$ for some $K \in \text{Contrib}(\alpha)_M$ and $s, s' \in \mathcal{S}$, with possibly $s = s'$, then

- (1) $K \subset F_s$,

- (2) either $F_s \subset F_{s'}$ or $F_{s'} \subset F_s$, and
(3) $\langle e_{F_s}^*, Z_s \rangle = \langle e_{F_{s'}}^*, Z_{s'} \rangle = 0$.

Conditions (1) and (2) are used below to associate a face in $\text{Contrib}(\alpha)_M$ to every cone of $\Sigma_{\mathcal{Z}}$ with nonzero intersection with σ_M . See Definition 6.4.3. Condition (3) is used to control the structure of cones in $\Sigma_{\mathcal{Z}}$ in a way that will allow us to apply Lemma 5.4.1. See Lemma 6.4.8.

Lemma 6.4.2. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, and let J be a nonempty face of $Q_{\mathcal{Z}}$ such that $\sigma_M \cap \tau_J \neq \{0\}$. Then $\{F_s : Z_s \in J\}$ is the set of elements of a chain of faces in Γ . Moreover, if $\sigma_K^\circ \cap \tau_J \neq \emptyset$ for some $K \in \text{Contrib}(\alpha)_M$, then $K \subset F_s$ for all $s \in \mathcal{S}$ such that $Z_s \in J$.*

Proof. There exists $K \in \text{Contrib}(\alpha)_M$ such that $\sigma_K^\circ \cap \tau_J \neq \emptyset$. Hence, for any $s, s' \in \mathcal{S}$ such that $Z_s, Z_{s'} \in J$, $\sigma_K^\circ \cap \tau_{Z_s} \cap \tau_{Z_{s'}} \neq \emptyset$. By property (2) of Definition 6.4.1, F_s and $F_{s'}$ are comparable under inclusion. Hence if we consider the set $\{F_s : Z_s \in J\}$ as a poset under inclusion, then all elements are comparable, and the poset is a chain. The second statement follows from property (1) of Definition 6.4.1. \square

Definition 6.4.3. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, and let J be a nonempty face of $Q_{\mathcal{Z}}$ such that $\sigma_M \cap \tau_J \neq \{0\}$. Then set $F_J := \max\{F_s : Z_s \in J\}$.*

Lemma 6.4.2 implies that the above is well-defined. Also, it follows from Lemma 6.3.2 that we can replace the condition $\sigma_M \cap \tau_J \neq \{0\}$ with the condition that $\tau_J \cap N_{M, \leq \delta} \neq \{0\}$ for some δ chosen sufficiently small.

Remark 6.4.4. Let J be a nonempty face of $Q_{\mathcal{Z}}$ and $K \in \text{Contrib}(\alpha)_M$ such that $\sigma_K^\circ \cap \tau_J \neq \emptyset$. Then Lemma 6.4.2 implies that $K \subset F_J$.

Remark 6.4.5. If $J \subset J'$ is an inclusion of nonempty faces of $Q_{\mathcal{Z}}$ and $\sigma_M \cap \tau_{J'} \neq \{0\}$, then $\sigma_M \cap \tau_J \neq \{0\}$ and $F_J \subset F_{J'}$.

Lemma 6.4.6. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, and let J be a nonempty face of $Q_{\mathcal{Z}}$ and $K \in \Gamma$ such that $\sigma_K \cap \tau_J \cap N_{M, \leq \delta} \neq \{0\}$ for some δ sufficiently small. Then $K \subset F_J$.*

Proof. Since $\sigma_K \cap \tau_J \cap N_{M, \leq \delta} \neq \{0\}$, Lemma 6.3.2 implies that $\sigma_K \cap \tau_J \cap \sigma_M \neq \{0\}$. In particular, F_J is well-defined. Since $\sigma_K \cap \sigma_M \neq \{0\}$, we have that $\sigma_K \cap \sigma_M = \sigma_{K'}$ for some $K' \in \text{Contrib}(\alpha)_M$. Since $\sigma_{K'} \cap \tau_J \neq \{0\}$, there exists $K' \subset K'' \in \text{Contrib}(\alpha)_M$ such that $\sigma_{K''}^\circ \cap \tau_J \neq \emptyset$. By Remark 6.4.4, $K'' \subset F_J$. Since $K \subset K' \subset K''$, the result follows. \square

Let Σ_1, Σ_2 be fans in \mathbb{R}^n dual to polyhedra P_1, P_2 in \mathbb{R}^n respectively. Then the Minkowski sum $P_1 + P_2$ is dual to the intersection $\Sigma_1 \cap \Sigma_2$ of Σ_1 and Σ_2 , where $\Sigma_1 \cap \Sigma_2$ is the fan consisting of all cones $\{\sigma_1 \cap \sigma_2 : \sigma_i \in \Sigma_i\}$. All faces of $P_1 + P_2$ have the form $J_1 + J_2$ for some faces J_i of P_i for $i = 1, 2$. If J_i is dual to σ_i in Σ_i for $i = 1, 2$, then a face of the form $J_1 + J_2$ is dual to $\sigma_1 \cap \sigma_2$. Conversely, every cone C in $\Sigma_1 \cap \Sigma_2$ has the form $C = \sigma_1 \cap \sigma_2$, where σ_i is the smallest face of Σ_i containing C for $i = 1, 2$. Then $C^\circ = \sigma_1^\circ \cap \sigma_2^\circ$, $\text{span}(C) = \text{span}(\sigma_1) \cap \text{span}(\sigma_2)$, and if σ_i is dual to a face J_i of P_i for $i = 1, 2$, then C is dual to $J_1 + J_2$. We will be interested in the polyhedron $\text{Newt}(f)_{\mathcal{Z}} := \text{Newt}(f) + Q_{\mathcal{Z}}$ dual to $\Sigma \cap \Sigma_{\mathcal{Z}}$.

Definition 6.4.7. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, let J be a nonempty face of $Q_{\mathcal{Z}}$, and let C be a cone in $\Sigma \cap \Sigma_{\mathcal{Z}}$. Assume that $C \subset \tau_J$ and $C \cap N_{M, \leq \delta} \neq \{0\}$ for some δ chosen sufficiently small. Let $D(C, J) = D(C, J, M, \delta) := (C \cap \sigma_{A_{F_J}} \cap N_{M, \leq \delta}) + \mathbb{R}_{\geq 0} e_{F_J}^*$.*

The following lemma will allow us to replace contributions to $Z_{\text{for}}(T)$ from certain cones C by contributions from cones $D(C, J)$, whose structure will allow us to apply Lemma 5.4.1.

Lemma 6.4.8. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, let J be a nonempty face of $Q_{\mathcal{Z}}$, and let C be a cone in $\Sigma \cap \Sigma_{\mathcal{Z}}$. Assume that $C \subset \tau_J$ and that $C \cap N_{M, \leq \delta} \neq \{0\}$ for some δ chosen sufficiently small. Then C is dual to a face $K + J'$ of $\text{Newt}(f)_{\mathcal{Z}}$, for some face $K \in \Gamma$ such that $K \subset F_J \subset F_{J'}$, and for some face J' of $Q_{\mathcal{Z}}$ such that $J \subset J'$. Suppose that $C \not\subset \sigma_{A_{F_J}}$. Then $C \cap N_{M, \leq \delta} = D(C, J) \cap N_{M, \leq \delta}$, and $e_{F_J}^* \in \text{span}(C)$.*

Proof. By Lemma 6.3.3, C is dual to a face of $\text{Newt}(f)_{\mathcal{Z}}$ of the form $K + J'$, for some $K \in \Gamma$, and some nonempty face J' of $Q_{\mathcal{Z}}$. Here $C = \sigma_K \cap \tau_{J'}$ and $\tau_{J'}$ is the smallest cone in $\Sigma_{\mathcal{Z}}$ containing C . In particular, $\tau_{J'} \subset \tau_J$ and $J \subset J'$. By Lemma 6.3.2, $\sigma_M \cap \tau_{J'} \neq \{0\}$, and $F_J, F_{J'}$ are well-defined. Also, $\{0\} \neq \sigma_K \cap \tau_J \cap N_{M, \leq \delta}$. Then Lemma 6.4.6 implies that $K \subset F_J$. By Remark 6.4.5, $F_J \subset F_{J'}$.

Fix a vertex V of K . Then

$$(25) \quad \sigma_K = \left(\bigcap_{V \neq V' \in K} H_{V'-V} \right) \cap \left(\bigcap_{V' \notin K} H_{V'-V, \geq 0} \right) \cap \left(\bigcap_{W \in \text{Unb}(C_K)} H_W \right) \cap \left(\bigcap_{W \in \{e_1, \dots, e_n\} \setminus \text{Unb}(C_K)} H_{W, \geq 0} \right),$$

where V' varies over vertices in Γ . Let $\hat{s} \in \mathcal{S}$ such that $Z_{\hat{s}} \in J$ and $F_{\hat{s}} = F_J$. Then $Z_{\hat{s}} \in J'$ and

$$(26) \quad \tau_{J'} = \left(\bigcap_{\substack{\hat{s} \neq s \in \mathcal{S} \\ Z_s \in J'}} H_{Z_s - Z_{\hat{s}}} \right) \cap \left(\bigcap_{\substack{s \in \mathcal{S} \\ Z_s \notin J'}} H_{Z_s - Z_{\hat{s}}, \geq 0} \right).$$

It follows that we may choose a finite collection of nonzero elements $P_C = \{W\} \subset \mathbb{Q}^n \setminus \{0\}$ such that each $W \in P_C$ is of the form either

- (1) $W = Z_s - Z_{\hat{s}}$ for some $s \in \mathcal{S}$,
- (2) $W = V' - V$ for some vertex V' in Γ ,
- (3) $W \in \{e_1, \dots, e_n\}$,

and $C = \sigma_K \cap \tau_{J'}$ is the intersection of half-spaces of the form $H_{W, \geq 0}$ or hyperplanes of the form H_W for various $W \in P_C$.

Suppose that $C \not\subset \sigma_{A_{F_J}}$. Note that $C \not\subset \sigma_{A_{F_J}}$ if and only if $\sigma_K \not\subset \sigma_{A_{F_J}}$, if and only if $A_{F_J} \notin K$. Let $\tilde{\sigma}$ be the intersection of all such half-spaces and hyperplanes appearing in the description (25) of σ_K such that the $W \in P_C$ that defines the hyperplane or half-space has $C \cap N_{M, \leq \delta} \cap H_W \neq \{0\}$, and W doesn't have the form $W = V' - V$ with $V' = A_{F_J}$. Similarly, let $\tilde{\tau}$ be the intersection of all such half-spaces and hyperplanes appearing in the description (26) of $\tau_{J'}$ such that $C \cap N_{M, \leq \delta} \cap H_W \neq \{0\}$. Let $\tilde{C} = \tilde{\sigma} \cap \tilde{\tau}$. Then, by construction, there exists a cone U over a small open neighborhood of $N_{M, \leq \delta} \cap S'$ in $S' = \{u \in \mathbb{R}^n : \langle u, \mathbf{1} \rangle = 1\}$ such that

$$(27) \quad C \cap U = \tilde{C} \cap H_{A_{F_J} - V, \geq 0} \cap U.$$

We claim that $\mathbb{R}e_{F_J}^* \subset \tilde{C}$. Let $H_{W, \geq 0}$ or H_W be a defining half-space or hyperplane of \tilde{C} . We need to show that $e_{F_J}^* \in H_W$. Equivalently, we need to show that $\langle e_{F_J}^*, W \rangle = 0$. By assumption, $C \cap N_{M, \leq \delta} \cap H_W \neq \{0\}$. By Lemma 6.3.2, $C \cap \sigma_M \cap H_W \neq \{0\}$.

First, assume that $W = V' - V$ for some vertex $V' \neq A_{F_J}$ in Γ . Then $\{0\} \neq C \cap N_{M, \leq \delta} \cap H_W \subset \sigma_{V'} \cap \tau_J \cap N_{M, \leq \delta}$, and Lemma 6.4.6 implies that $V' \in F_J$. Then V', V are vertices of F_J that are not equal to A_{F_J} , and hence $\langle e_{F_J}^*, V' \rangle = \langle e_{F_J}^*, V \rangle = 0$, so $\langle e_{F_J}^*, W \rangle = 0$.

Second, assume that $W = Z_s - Z_{\hat{s}}$ for some $s \in \mathcal{S}$. Since $C \subset \tau_{J'}$, we have $\{0\} \neq C \cap \sigma_M \cap H_W \subset \tau_{J'} \cap \sigma_M \cap H_W \subset \sigma_M \cap \tau_{Z_s} \cap \tau_{Z_{\hat{s}}}$. Hence, there exists $K' \in \text{Contrib}(\alpha)_M$ such that $\sigma_{K'}^\circ \cap \tau_{Z_s} \cap \tau_{Z_{\hat{s}}} \neq \emptyset$. By (3) in Definition 6.4.1, $\langle e_{F_J}^*, Z_s \rangle = \langle e_{F_J}^*, Z_{\hat{s}} \rangle = 0$, so $\langle e_{F_J}^*, W \rangle = 0$.

Finally, assume $W \in \{e_1, \dots, e_n\}$. Then $\{0\} \neq C \cap N_{M, \leq \delta} \cap H_W \subset \sigma_K \cap \tau_J \cap N_{M, \leq \delta} \cap H_W$. By Lemma 6.3.2, $\sigma_K \cap \tau_J \cap \sigma_M \cap H_W \neq \{0\}$. It follows that there exists $K' \in \text{Contrib}(\alpha)_M$ such that $K \subset K'$ and $\sigma_{K'}^\circ \cap \tau_J \cap H_W \neq \{0\}$. By Remark 6.4.4, $K' \subset F_J$. Then $W \in \text{Unb}(C_{K'}) \subset \text{Unb}(C_{F_J})$, and hence $\langle e_{F_J}^*, W \rangle = 0$ by Remark 6.2.5.

We conclude that $\mathbb{R}e_{F_J}^* \subset \tilde{C}$. Next, we claim that

$$(28) \quad \tilde{C} \cap H_{A_{F_J} - V, \geq 0} \cap N_{M, \leq \delta} = ((\tilde{C} \cap H_{A_{F_J} - V} \cap N_{M, \leq \delta}) + \mathbb{R}_{\geq 0} e_{F_J}^*) \cap N_{M, \leq \delta}.$$

By (27), the left hand side of (28) is $C \cap N_{M, \leq \delta}$. Let $u \in \tilde{C} \cap \mathbf{H}_{A_{F_J} - V, \geq 0} \cap N_{M, \leq \delta}$. We aim to show that u lies in the right-hand side of (28). It is enough to consider the case when $\langle u, \mathbf{1} \rangle = 1$. Consider the function

$$\phi: \mathbb{R}_{\geq 1} \rightarrow S' \subset \mathbb{R}^n, \text{ defined by } \phi(\lambda) = \lambda u + (1 - \lambda)e_{F_J}^*.$$

Since $\mathbb{R}e_{F_J}^* \subset \tilde{C}$, the image of ϕ is contained in \tilde{C} . It is enough to show that $\phi^{-1}(\mathbf{H}_{A_{F_J} - V} \cap N_{M, \leq \delta}) \neq \emptyset$, since if $\lambda \in \phi^{-1}(\mathbf{H}_{A_{F_J} - V} \cap N_{M, \leq \delta})$, then

$$(29) \quad u = (1/\lambda)(\phi(\lambda) + (\lambda - 1)e_{F_J}^*),$$

and u lies in the right-hand side of (28). Moreover, if we choose $u \in C^\circ$, then $u \notin \sigma_{A_{F_J}}$ and hence $\lambda > 1$ in (29). Then $e_{F_J}^* = (1/(\lambda - 1))(\lambda u - \phi(\lambda)) \in \text{span}(C)$, which establishes the last statement of the lemma.

Consider the linear function

$$f(\lambda) = \langle \phi(\lambda), W_M - V \rangle.$$

Since $u \in S' \cap N_{M, \leq \delta}$, $f(1) \leq \delta$. Since $\langle e_{F_J}^*, V \rangle = 0$, we compute: $f'(\lambda) = f(1) - \langle e_{F_J}^*, W_M \rangle \leq \delta - \langle e_{F_J}^*, W_M \rangle < 0$. The last inequality follows since M is interior implies that $\langle e_{F_J}^*, W_M \rangle > 0$ and δ is chosen sufficiently small. In particular, the image of f is unbounded because it is a non-constant linear function. Hence the image of ϕ is unbounded and so $\phi^{-1}(U \setminus N_{M, \leq \delta}) \neq \emptyset$.

We claim that $\phi^{-1}(\mathbf{H}_{A_{F_J} - V, \geq 0} \cap U) \subset \phi^{-1}(N_{M, \leq \delta})$. Indeed, if $\phi(\lambda) \in \mathbf{H}_{A_{F_J} - V, \geq 0} \cap U$, then $\phi(\lambda) \in C$ by (27). Then $\langle \phi(\lambda), W_M \rangle - N(\phi(\lambda)) = f(\lambda) \leq f(1) \leq \delta$. Since $\langle \phi(\lambda), \mathbf{1} \rangle = 1$, we deduce that $\phi(\lambda) \in N_{M, \leq \delta}$.

It follows that $\emptyset \neq \phi^{-1}(U \setminus N_{M, \leq \delta}) \subset \phi^{-1}(\mathbf{H}_{A_{F_J} - V, < 0} \cap U)$. Since $1 \in \phi^{-1}(\mathbf{H}_{A_{F_J} - V, \geq 0} \cap U)$, we deduce that $\phi^{-1}(\mathbf{H}_{A_{F_J} - V} \cap N_{M, \leq \delta}) = \phi^{-1}(\mathbf{H}_{A_{F_J} - V} \cap U) \neq \emptyset$, so the left-hand side is contained in the right-hand side of (28).

Conversely, since $\mathbb{R}e_{F_J}^* \subset \tilde{C}$ and $e_{F_J}^* \in \mathbf{H}_{A_{F_J} - V, \geq 0}$, the right-hand side of (28) is contained in $\tilde{C} \cap \mathbf{H}_{A_{F_J} - V, \geq 0} \cap N_{M, \leq \delta}$. This establishes (28). By (27),

$$\tilde{C} \cap \mathbf{H}_{A_{F_J} - V} \cap N_{M, \leq \delta} = (\tilde{C} \cap \mathbf{H}_{A_{F_J} - V, \geq 0} \cap N_{M, \leq \delta}) \cap \sigma_{A_{F_J}} = (C \cap N_{M, \leq \delta}) \cap \sigma_{A_{F_J}}.$$

Substituting this expression into the right-hand side of (28) and combining with (27), we deduce that

$$C \cap N_{M, \leq \delta} = \tilde{C} \cap \mathbf{H}_{A_{F_J} - V, \geq 0} \cap N_{M, \leq \delta} = ((C \cap \sigma_{A_{F_J}} \cap N_{M, \leq \delta}) + \mathbb{R}_{\geq 0}e_{F_J}^*) \cap N_{M, \leq \delta}. \quad \square$$

The following lemma is a corollary of the proof of Lemma 6.4.8.

Lemma 6.4.9. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, and let J be a nonempty face of $Q_{\mathcal{Z}}$ such that $\tau_J \cap N_{M, \leq \delta} \neq \{0\}$ for some δ chosen sufficiently small. Then no ray of $\tau_J \cap N_{M, \leq \delta}$ is contained in σ_M .*

Proof. The proof of Lemma 6.4.8, with σ_K replaced by $\mathbb{R}_{\geq 0}^n$ and $\tau_{J'}$ replaced by τ_J , shows that there exists a polyhedral cone τ' and a cone U over a small open neighborhood of $N_{M, \leq \delta} \cap S'$ in S' such that

- (1) $\mathbb{R}e_{F_J}^* \subset \tau'$, and
- (2) $\mathbb{R}_{\geq 0}^n \cap \tau_J \cap U = \tau' \cap U$.

Let u be a generator of a ray in $\tau_J \cap N_{M, \leq \delta}$. We may assume that $\langle u, \mathbf{1} \rangle = 1$. Suppose that $\langle u, W_M \rangle - N(u) < \delta$. Fix $0 < \epsilon \ll 1$ and let $L_\epsilon = \{u + \lambda e_{F_J}^* : |\lambda| < \epsilon\}$. Then $L_\epsilon \subset \tau' \cap U = \mathbb{R}_{\geq 0}^n \cap \tau_J \cap U$. It follows that $L_\epsilon \subset \tau_J \cap N_{M, \leq \delta}$, contradicting the assumption that u generates a ray. We deduce that $\langle u, W_M \rangle - N(u) = \delta$. In particular, $u \notin \sigma_M$. \square

Definition 6.4.10. *Let $(\mathcal{Z}, \mathcal{F})$ be a weakly α -compatible pair, let J be a nonempty face of $Q_{\mathcal{Z}}$, and let C be a cone in $\Sigma \cap \Sigma_{\mathcal{Z}}$. Then we say (C, J) is α -critical if the following properties hold:*

- (1) $C \cap \sigma_M \neq \{0\}$,
- (2) $C \subset \sigma_{A_{F_J}} \cap \tau_J$, and
- (3) C is critical with respect to (α, A_{F_J}) .

Definition 6.4.11. We say a weakly α -compatible pair $(\mathcal{Z}, \mathcal{F})$ is α -compatible if for every α -critical pair (C, J) , $C \subset \sigma_M$.

Note that the notion of an α -compatible pair depends on the choice of a minimal face M . The main technical result required to prove Theorem 6.1.2 is following result on the existence of α -compatible pairs.

Theorem 6.4.12. Let $\alpha \notin \mathbb{Z}$, and assume that all faces of $\text{Contrib}(\alpha)$ are UB_1 and $\text{Newt}(f)$ is α -simplicial. Then for any minimal face $M \in \text{Contrib}(\alpha)$, there exists an α -compatible pair.

Lemma 6.4.13. Consider an α -compatible pair $(\mathcal{Z}, \mathcal{F})$. Let γ be a ray of $\Sigma \cap \Sigma_{\mathcal{Z}} \cap N_{M, \leq \delta}$ for some δ chosen sufficiently small. Assume that $\gamma \not\subset \sigma_M$. Then γ is not critical with respect to α . Moreover, if $\gamma \subset \tau_J$ for some face J of $Q_{\mathcal{Z}}$, then γ is not critical with respect to (α, A_{F_J}) .

Proof. There is a unique face J' of $Q_{\mathcal{Z}}$ such that $\gamma \subset \tau_{J'}$. If $\gamma \subset \tau_J$, then it follows that $J \subset J'$. Let C be the smallest cone in $\Sigma \cap \Sigma_{\mathcal{Z}}$ containing γ . Then $\tau_{J'}$ is the smallest cone of $\Sigma_{\mathcal{Z}}$ containing C . By Lemma 6.4.8, C is dual to a face $K + J'$ of $\text{Newt}(f)_{\mathcal{Z}}$, where $K \in \Gamma$ such that $K \subset F_J$. Assume that γ is critical with respect to (α, W) for some vertex W of either K or M . Note that one possible choice for W is A_{F_J} , since $A_{F_J} \in M$. Then Lemma 6.3.9 implies that C is critical with respect to (α, W) , and $C \cap \sigma_M \neq \{0\}$.

First, assume that $\gamma \subset \sigma_{A_{F_J}}$. Then $C \subset \sigma_{A_{F_J}}$. If $W = A_{F_J}$, then (C, J) is α -critical, and Definition 6.4.11 implies that $\gamma \subset C \subset \sigma_M$, a contradiction. We conclude that $W \neq A_{F_J}$. That is, γ is not critical with respect to (α, A_{F_J}) . Equivalently, in this case, γ is not critical with respect to α .

Second, assume that $\gamma \not\subset \sigma_{A_{F_J}}$. Then $C \not\subset \sigma_{A_{F_J}}$. Equivalently, $A_{F_J} \notin K$. By Lemma 6.4.8, $e_{F_J}^* \in \text{span}(C)$. Let V be a vertex of $K \subset F_J$. Since $V \neq A_{F_J}$, $\langle e_{F_J}^*, V \rangle = 0$, and hence $\langle e_{F_J}^*, \alpha V + \mathbf{1} \rangle = 1 \neq 0$. We deduce that C is not critical with respect to (α, V) . Hence $W \neq V$, and γ is not critical with respect to α . Similarly, since $\alpha \neq -1$ by assumption, $\langle e_{F_J}^*, \alpha A_{F_J} + \mathbf{1} \rangle = \alpha + 1 \neq 0$, and hence C is not critical with respect to (α, A_{F_J}) . Therefore $W \neq A_{F_J}$, and γ is not critical with respect to (α, A_{F_J}) . \square

Proof of Theorem 6.1.2. Let $\alpha \notin \mathbb{Z}$, and assume that all faces of $\text{Contrib}(\alpha)$ are UB_1 and $\text{Newt}(f)$ is α -simplicial. By Remark 5.3.3 and Remark 5.3.6, it is enough to show that there exists a set of candidate poles for $Z_{\text{for}}(T)$ not containing α . Let M_1, \dots, M_r be the minimal elements in $\text{Contrib}(\alpha)$. Assume that δ is chosen sufficiently small. By Lemma 6.3.4,

$$Z_{\text{for}}(T) = Z_{\text{for}}(T)|_{\cap_i N_{M_i, \geq \delta}} + \sum_{i=1}^r Z_{\text{for}}(T)|_{N_{M_i, < \delta}}.$$

By Lemma 6.3.12, $Z_{\text{for}}(T)|_{\cap_i N_{M_i, \geq \delta}}$ lies in R and admits a set of candidate poles not containing α . Choose a minimal face $M \in \text{Contrib}(\alpha)$. By Theorem 6.4.12, we may fix an α -compatible pair $(\mathcal{Z}, \mathcal{F})$. By definition, $Z_{\text{for}}(T)|_{\partial \mathbb{R}_{\geq 0}^n} = 0$. Recall that $N_{M, \leq \delta}^\circ = N_{M, < \delta} \cap \mathbb{R}_{> 0}^n$. We have

$$Z_{\text{for}}(T)|_{N_{M, < \delta}} = Z_{\text{for}}(T)|_{N_{M, \leq \delta}^\circ} = \sum_{\substack{\emptyset \neq J \subset Q_{\mathcal{Z}} \\ \tau_J \cap N_{M, \leq \delta}^\circ \neq \emptyset}} Z_{\text{for}}(T)|_{\tau_J \cap N_{M, \leq \delta}^\circ}.$$

Let J be a nonempty face of $Q_{\mathcal{Z}}$ such that $\tau_J \cap N_{M, \leq \delta}^\circ \neq \emptyset$. By Remark 5.3.4, to show that $Z_{\text{for}}(T)$ lies in R and admits a set of candidate poles not containing α , it is enough to show that $Z_{\text{for}}(T)|_{\tau_J \cap N_{M, \leq \delta}^\circ}$ lies in R and admits a set of candidate poles not containing α for every J .

Let C be a cone in $\Sigma \cap \Sigma_{\mathcal{Z}}$ such that $C^\circ \subset \tau_J^\circ$ and $C \cap N_{M, \leq \delta}^\circ \neq \emptyset$. By Lemma 6.4.8, $C = \sigma_G \cap \tau_J$ and $C^\circ = \sigma_G^\circ \cap \tau_J^\circ$ for some face $G \in \Gamma$ with $G \subset F_J$. Since $C \not\subset \partial \mathbb{R}_{\geq 0}^n$, G is compact.

Suppose that $C \not\subset \sigma_{A_{F_J}}$. Then $A_{F_J} \notin G$, and G is contained in the base of the (possibly unbounded) B_1 -face F_J . Since F_J is a B_1 -face, we may consider the face $F = \text{Conv}\{G, A_{F_J}\}$ of F_J . Then F is a compact

B_1 -face with apex A_{F_J} and base G in the direction $e_{F_J}^*$. Consider the face $C' = C \cap \sigma_{A_{F_J}} \cap N_{M, \leq \delta}$ of $C \cap N_{M, \leq \delta}$. Note that $C' \subset C \cap \sigma_{A_{F_J}} \subset \sigma_F$. By Lemma 6.4.8,

$$(30) \quad C \cap N_{M, \leq \delta} = D \cap N_{M, \leq \delta},$$

where $D = D(C, J) = C' + \mathbb{R}_{\geq 0} e_{F_J}^* \subset \sigma_G$. By Lemma 6.4.13, the rays of $C \cap N_{M, \leq \delta}$ that are critical with respect to either α or (α, A_{F_J}) are contained σ_M . If V is a vertex of G , then $\langle e_{F_J}^*, V \rangle = 0$, and hence $\langle e_{F_J}^*, \alpha V + \mathbf{1} \rangle = 1 \neq 0$. Also, $\langle e_{F_J}^*, \alpha A_{F_J} + \mathbf{1} \rangle = \alpha + 1 \neq 0$, by assumption. We deduce that $e_{F_J}^*$ is not critical with respect to α or (α, A_{F_J}) . This implies that no ray of $D \cap N_{M, \geq \delta}$ is critical with respect to α or (α, A_{F_J}) . Since $\sigma_G^\circ, \sigma_F^\circ \subset \mathbb{R}_{> 0}^n$, Remark 6.3.7 gives the following equalities:

$$\begin{aligned} (C \cap \sigma_{A_{F_J}})^\circ \cap N_{M, \leq \delta}^\circ &= (C')^\circ, \\ C^\circ \cap N_{M, \leq \delta}^\circ &= (C \cap N_{M, \leq \delta})^\circ = (D \cap N_{M, \leq \delta})^\circ = D^\circ \cap N_{M, \leq \delta}^\circ, \text{ and} \\ D^\circ \cap N_{M, \geq \delta} &= (D \cap N_{M, \geq \delta})^\circ \cup (D \cap N_{M, \delta})^\circ. \end{aligned}$$

Since the restriction of N to σ_G is linear, Lemma 5.2.5 and Lemma 6.3.5, imply that the following elements of \tilde{R} lie in R and admit sets of candidate poles not containing α :

$$(L-1)^{n-\dim G} \sum_{u \in D^\circ \cap N_{M, \geq \delta} \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{N(u)}, \text{ and } (L-1)^n \sum_{u \in D^\circ \cap N_{M, \geq \delta} \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A_{F_J} \rangle}.$$

We claim that $(C')^\circ \subset \sigma_F^\circ$. We have $(C')^\circ \subset \sigma_{F'}^\circ$ for some $F' \subset F_J$. By Lemma 6.4.6, $F' \subset F_J$. If $\sigma_{F'} \subset \partial \mathbb{R}_{\geq 0}^n$, then (30) implies that $C \cap N_{M, \leq \delta} \subset \partial \mathbb{R}_{\geq 0}^n$, a contradiction. Hence F' is a compact face of F_J containing A_{F_J} . It follows that F' is a B_1 -face with apex A_{F_J} and base $G' \supset G$. Then (30) implies that $C \cap N_{M, \leq \delta} \subset \sigma_{G'}$. Then $\emptyset \neq (C \cap N_{M, \leq \delta})^\circ = C^\circ \cap N_{M, \leq \delta}^\circ \subset \sigma_{G'}^\circ$. Since $C^\circ \subset \sigma_G^\circ$, we conclude that $G = G'$, and hence $F = F'$, completing the proof of the claim. We may then apply Lemma 5.4.1 to obtain

$$Z_{\text{for}}(T)|_{(D^\circ \cup (C')^\circ)} = (L-1)^n \left(\sum_{u \in (D^\circ \cup (C')^\circ) \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A_{F_J} \rangle} \right) \in \tilde{R}.$$

Since $D^\circ = (D^\circ \cap N_{M, \geq \delta}) \cup (D^\circ \cap N_{M, \leq \delta}^\circ)$, using the above calculations and Remark 5.3.4, we deduce that

$$(31) \quad Z_{\text{for}}(T)|_{((C \cap \sigma_{A_{F_J}})^\circ \cup C^\circ) \cap N_{M, \leq \delta}^\circ} - (L-1)^n \left(\sum_{u \in (((C \cap \sigma_{A_{F_J}})^\circ \cup C^\circ) \cap N_{M, \leq \delta}^\circ) \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A_{F_J} \rangle} \right)$$

lies in R and admits a set of candidate poles not containing α .

Since $\Sigma \cap \Sigma_{\mathcal{Z}}$ refines $\Sigma_{\mathcal{Z}}$, we may consider the subfan $\Sigma \cap \Sigma_{\mathcal{Z}} \cap N_{M, \leq \delta}|_{\tau_J}$ of $\Sigma \cap \Sigma_{\mathcal{Z}} \cap N_{M, \leq \delta}$. It follows from the description of $\Sigma \cap \Sigma_{\mathcal{Z}} \cap N_{M, \leq \delta}$ in Lemma 6.3.6 and the fact that $\tau_J \cap \mathbb{R}_{> 0}^n \neq \emptyset$ that

$$(32) \quad (\tau_J \cap N_{M, \leq \delta})^\circ = \bigcup_{\substack{C \in \Sigma \cap \Sigma_{\mathcal{Z}} \\ C^\circ \subset \tau_J^\circ \\ C \cap N_{M, \leq \delta}^\circ \neq \emptyset}} (C \cap N_{M, \leq \delta})^\circ = \bigcup_{\substack{C \in \Sigma \cap \Sigma_{\mathcal{Z}} \\ C^\circ \subset \tau_J^\circ \\ C \cap N_{M, \leq \delta}^\circ \neq \emptyset}} C^\circ \cap N_{M, \leq \delta}^\circ = \tau_J^\circ \cap N_{M, \leq \delta}^\circ.$$

It follows from (30) that we may rewrite this as:

$$\tau_J^\circ \cap N_{M, \leq \delta}^\circ = (\sigma_{A_{F_J}}^\circ \cap \tau_J^\circ \cap N_{M, \leq \delta}^\circ) \cup \bigcup_{\substack{C \in \Sigma \cap \Sigma_{\mathcal{Z}} \\ C^\circ \subset \tau_J^\circ \\ C \cap N_{M, \leq \delta}^\circ \neq \emptyset \\ C \not\subset \sigma_{A_{F_J}}} ((C \cap \sigma_{A_{F_J}})^\circ \cup C^\circ) \cap N_{M, \leq \delta}^\circ.$$

We deduce from Remark 5.4.2 and (31) that

$$Z_{\text{for}}(T)|_{\tau_J^\circ \cap N_{M, \leq \delta}^\circ} - (L-1)^n \left(\sum_{u \in \tau_J^\circ \cap N_{M, \leq \delta}^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A_{F_J} \rangle} \right)$$

lies in R and admits a set of candidate poles not containing α . By Remark 5.3.4 and (32), it is enough to show that

$$(L-1)^n \left(\sum_{u \in (\tau_J \cap N_{M, \leq \delta})^\circ \cap \mathbb{N}^n} L^{-\langle u, \mathbf{1} \rangle} T^{\langle u, A_{F_J} \rangle} \right)$$

lies in R and admits a set of candidate poles not containing α . By Lemma 6.4.9 and Lemma 6.4.13, no rays of $\tau_J \cap N_{M, \leq \delta}$ are critical with respect to (α, A_{F_J}) . The result now follows from Lemma 5.2.5 and Lemma 6.3.5. \square

6.5. Existence of α -compatible sets. We continue with the notation of the previous section. Recall that we consider pairs $(\mathcal{Z}, \mathcal{F})$, where $\mathcal{Z} = (Z_s)_{s \in \mathcal{S}}$ and $\mathcal{F} = (F_s)_{s \in \mathcal{S}}$ are collections of elements of \mathbb{Q}^n and $\text{Contrib}(\alpha)_M$ respectively.

Definition 6.5.1. *Consider a pair $(\mathcal{Z}, \mathcal{F})$. Then $(\mathcal{Z}, \mathcal{F})$ is restricted if $Z_s \in \text{span}(\{V_M\} \cup \text{Gen}(C_{F_s} \setminus C_M) \cup \mathcal{A}_M)$ for every $s \in \mathcal{S}$.*

Our goal is to reduce the existence of an α -compatible set to the existence of a restricted, weakly α -compatible set. We will consider sets $\epsilon = \{\epsilon_s\}_{s \in \mathcal{S}} \in \mathbb{R}^{\mathcal{S}}$; note that we allow ϵ_s to be negative. We say that ϵ is chosen to be sufficiently small if $|\epsilon_s|$ is chosen to be sufficiently small for all $s \in \mathcal{S}$. Explicitly, a property holds for ϵ sufficiently small if there exists $\delta > 0$ such that the property holds for all ϵ such that $|\epsilon_s| < \delta$ for all $s \in \mathcal{S}$. Given a sequence of sets $\{\epsilon_m\}_{m \in \mathbb{Z}_{\geq 0}}$, where $\epsilon_m = \{\epsilon_{m,s}\}_{s \in \mathcal{S}}$, for some $\epsilon_{m,s} \in \mathbb{R}$, we write $\lim_{m \rightarrow \infty} \epsilon_m = 0$ if $\lim_{m \rightarrow \infty} \epsilon_{m,s} = 0$ for all $s \in \mathcal{S}$. Given a set $\mathcal{Z} = \{Z_s\}_{s \in \mathcal{S}}$ of elements in \mathbb{R}^n , we let $Z_s(\epsilon_s) := Z_s + \epsilon_s V_M$, and let $\mathcal{Z}(\epsilon) := \{Z_s(\epsilon_s)\}_{s \in \mathcal{S}}$. Then $Q_{\mathcal{Z}(\epsilon)}$ is the convex hull of the elements of $\mathcal{Z}(\epsilon)$, and is dual to the fan $\Sigma_{\mathcal{Z}(\epsilon)}$. Also, $\text{Newt}(f)_{\mathcal{Z}(\epsilon)} = \text{Newt}(f) + Q_{\mathcal{Z}(\epsilon)}$ is dual to $\Sigma \cap \Sigma_{\mathcal{Z}(\epsilon)}$.

Lemma 6.5.2. *Consider a pair $(\mathcal{Z}, \mathcal{F})$. For $\epsilon \in \mathbb{Q}^{\mathcal{S}}$ sufficiently small, $(\mathcal{Z}(\epsilon), \mathcal{F})$ is restricted if $(\mathcal{Z}, \mathcal{F})$ is restricted, and $(\mathcal{Z}(\epsilon), \mathcal{F})$ is weakly α -compatible if $(\mathcal{Z}, \mathcal{F})$ is weakly α -compatible.*

Proof. Assume that $(\mathcal{Z}, \mathcal{F})$ is restricted. Since $Z_s(\epsilon_s)$ is a linear combination of Z_s and V_M , it follows that $(\mathcal{Z}(\epsilon), \mathcal{F})$ is restricted.

Assume that $(\mathcal{Z}, \mathcal{F})$ is weakly α -compatible. We want to show that $(\mathcal{Z}(\epsilon), \mathcal{F})$ is weakly α -compatible. Fix a face $K \in \text{Contrib}(\alpha)_M$ and $s, s' \in \mathcal{S}$. There is nothing to show if, after possibly shrinking ϵ , $\sigma_K^\circ \cap \tau_{\mathcal{Z}(\epsilon), s} \cap \tau_{\mathcal{Z}(\epsilon), s'} = \emptyset$. Hence we may assume that there exists a sequence of sets $\{\epsilon_m\}_{m \in \mathbb{Z}_{\geq 0}}$ such that $\lim_{m \rightarrow \infty} \epsilon_m = 0$ and $\sigma_K^\circ \cap \tau_{\mathcal{Z}(\epsilon_m), s} \cap \tau_{\mathcal{Z}(\epsilon_m), s'} \neq \emptyset$. Then Bolzano–Weierstrass implies that $\sigma_K \cap \tau_{\mathcal{Z}, s} \cap \tau_{\mathcal{Z}, s'} \neq \{0\}$. Hence there exists $K \subset K'$ such that $\sigma_{K'}^\circ \cap \tau_{\mathcal{Z}, s} \cap \tau_{\mathcal{Z}, s'} \neq \emptyset$. Since \mathcal{Z} is weakly α -compatible, we deduce that $K' \subset F_s$, either $F_{s'} \subset F_s$ or $F_s \subset F_{s'}$, and $\langle e_{F_s}^*, Z_s \rangle = \langle e_{F_s}^*, Z_{s'} \rangle = 0$. Then $K \subset K' \subset F_s$. By (24), $\langle e_{F_s}^*, Z_s(\epsilon_s) \rangle = \langle e_{F_s}^*, Z_s \rangle = 0$ and $\langle e_{F_s}^*, Z_{s'}(\epsilon_{s'}) \rangle = \langle e_{F_s}^*, Z_{s'} \rangle = 0$. \square

Before proceeding, we need a series of basic lemmas on deforming polyhedra. Let $\text{rec}(P)$ denote the recession cone of a polyhedron P . Let σ^\vee denote the dual cone to a cone σ . Given a face K of P , let τ_K denote the corresponding cone in the dual fan to P . In particular, $\text{rec}(K)$ is a face of $\text{rec}(P)$, $\tau_{\text{rec}(K)}$ is a face of $\text{rec}(P)^\vee$, and $\tau_K^\circ \subset \tau_{\text{rec}(K)}^\circ$.

Fix a nonempty finite set T . Let $P = \text{Conv}\{V_t : t \in T\} + \sigma \subset \mathbb{R}^n$ be a polyhedron, for some $V_t \in \mathbb{R}^n$, and some pointed (polyhedral) recession cone $\sigma = \text{rec}(P)$. Let $\{P(\epsilon) = \text{Conv}\{V_t(\epsilon) : t \in T\} + \sigma\}_\epsilon$ be a set of polyhedra with the same recession fan indexed by $\epsilon \in \mathbb{R}^\ell$ for some $\ell \geq 1$. Assume that $V_t(\epsilon) \in \mathbb{R}^n$ is a

continuous function of $\epsilon \in \mathbb{R}^\ell$, for all t in T . If $J(\epsilon)$ is a nonempty face of $P(\epsilon)$ and J is a nonempty face of P , we write $T(J(\epsilon)) := \{t \in T : V_t(\epsilon) \in J(\epsilon)\}$ and $T(J) := \{t \in T : V_t \in J\}$. We may also consider the recession cones $\text{rec}(J(\epsilon))$ and $\text{rec}(J)$, which are both faces of σ .

Definition 6.5.3. *For fixed $\epsilon \in \mathbb{R}^\ell$, we say $P(\epsilon)$ refines P if for any proper nonempty face $J(\epsilon)$ of $P(\epsilon)$, there exists a proper nonempty face J of P such that $T(J(\epsilon)) \subset T(J)$ and $\text{rec}(J(\epsilon)) \subset \text{rec}(J)$.*

Lemma 6.5.4. *For $\epsilon \in \mathbb{R}^\ell$ sufficiently small, $P(\epsilon)$ refines P .*

Proof. Assume the conclusion fails. Then there exists a sequence $\{\epsilon_m\}_{m \in \mathbb{Z}_{\geq 0}}$ such that $\lim_{m \rightarrow \infty} \epsilon_m = 0$, and a sequence of proper nonempty faces $J(\epsilon_m)$ of $P(\epsilon_m)$, such that, for any m and any proper nonempty face J of P , either $T(J(\epsilon_m)) \not\subset T(J)$ or $\text{rec}(J(\epsilon_m)) \not\subset \text{rec}(J)$. Since T is finite and σ has finitely many faces, after possibly replacing $\{\epsilon_m\}_{m \in \mathbb{Z}_{\geq 0}}$ by a subsequence, we may assume that $T(J(\epsilon_m))$ and $\text{rec}(J(\epsilon_m))$ are independent of m . Denote these by $\tilde{T} = T(J(\epsilon_m))$ and $\tilde{R} = \text{rec}(J(\epsilon_m))$ respectively. Consider a sequence u_m of elements in $\tau_{J(\epsilon_m)}^\circ \subset \tau_{\tilde{R}}^\circ$ such that $\|u_m\| = 1$. After possibly replacing $\{\epsilon_m\}_{m \in \mathbb{Z}_{\geq 0}}$ by a subsequence, we may assume that $\lim_{m \rightarrow \infty} u_m = u \in \tau_J^\circ \subset \tau_{\text{rec}(J)}^\circ$ exists for some nonempty face J of P . Since $\tau_{\tilde{R}}$ is closed, $u \in \tau_{\tilde{R}}$, and hence $\tau_{\text{rec}(J)} \subset \tau_{\tilde{R}}$ and $\tilde{R} \subset \text{rec}(J)$. For any $t \in \tilde{T}$ and $t' \in T$,

$$(33) \quad \langle u, V_t \rangle = \lim_{m \rightarrow \infty} \langle u_m, V_t \rangle = \lim_{m \rightarrow \infty} \langle u_m, V_t(\epsilon_m) \rangle \leq \lim_{m \rightarrow \infty} \langle u_m, V_{t'}(\epsilon_m) \rangle = \lim_{m \rightarrow \infty} \langle u_m, V_{t'} \rangle = \langle u, V_{t'} \rangle,$$

and hence $V_t \in J$. We deduce that $\tilde{T} \subset T(J)$, a contradiction. \square

Definition 6.5.5. *We say that $\{P(\epsilon)\}_\epsilon$ is locally combinatorially constant if for any ϵ sufficiently small, and for any nonempty face $J(\epsilon)$ of $P(\epsilon)$, there exists a (unique) nonempty face J of P such that $T(J(\epsilon)) = T(J)$ and $\text{rec}(J(\epsilon)) = \text{rec}(J)$, and, moreover, every nonempty face J of P appears in this way.*

Lemma 6.5.6. *After possibly replacing P with $P(\epsilon)$ for some $\epsilon \in \mathbb{Q}^\ell$, $\{P(\epsilon)\}_\epsilon$ is locally combinatorially constant.*

Proof. Lemma 6.5.4 implies that we may order $\{P(\epsilon)\}_\epsilon$ by refinement. Since T is finite and σ has finitely many faces, there exists an $\epsilon \in \mathbb{R}^\ell$ such that $Q = P(\epsilon)$ is minimal under this ordering. Then $\{Q(\epsilon)\}_\epsilon$ is locally combinatorially constant. Consider $\epsilon' \in \mathbb{R}^\ell$ such that $\epsilon + \epsilon' \in \mathbb{Q}^\ell$, and let $Q' = Q(\epsilon')$. Then for ϵ' sufficiently small, $\{Q'(\epsilon)\}_\epsilon$ is locally combinatorially constant. \square

Given a family of cones $\{C_k\}_{k=1}^\infty$, define $\limsup C_k$ to be the cone of points $u \in \mathbb{R}^n$ such that u is a limit point of a sequence of points $u_k \in C_k$, i.e., there exists a subsequence of u_k converging to u .

Lemma 6.5.7. *Assume that $\{P(\epsilon)\}_\epsilon$ is locally combinatorially constant. Fix a nonempty face J of P . For ϵ sufficiently small, let $J(\epsilon)$ be the nonempty face of $P(\epsilon)$ such that $T(J(\epsilon)) = T(J)$ and $\text{rec}(J(\epsilon)) = \text{rec}(J)$. Consider any $\{\epsilon_k\}_{k \in \mathbb{Z}_{\geq 0}}$ such that $\lim_{m \rightarrow \infty} \epsilon_k = 0$. Then $\limsup_k \tau_{J(\epsilon_k)} \subset \tau_J$, and, if we assume that $\dim P = n$, then $\limsup_k \tau_{J(\epsilon_k)} = \tau_J$.*

Proof. We first show that $\limsup_k \tau_{J(\epsilon_k)} \subset \tau_J$. Suppose that $u_k \in \tau_{J(\epsilon_k)}$, and that, after possibly replacing $\{\epsilon_k\}_{k \in \mathbb{Z}_{\geq 0}}$ with a subsequence, $\lim_{k \rightarrow \infty} u_k = u \in \mathbb{R}^n$ exists. Then $u \in \tau_{J'}$, for some nonempty face J' of P . Then the calculation in (33) implies that $T(J) = T(J(\epsilon_k)) \subset T(J')$, and hence $J \subset J'$ and $u \in \tau_{J'} \subset \tau_J$.

We need to prove the converse statement. Assume that $\dim P = n$. Then τ_J is generated by its rays $\{\gamma_m\}_{1 \leq m \leq p}$. For any ϵ sufficiently small, let $\{\gamma_m(\epsilon)\}_{1 \leq m \leq p}$ denote the corresponding rays in the dual fan to $P(\epsilon)$. Consider elements $\{u_{m,k} \in \gamma_m(\epsilon_k) : 1 \leq m \leq p, k \geq 0\}$ such that $\|u_{m,k}\| = 1$. Then, after possibly replacing $\{\epsilon_k\}_{k \in \mathbb{Z}_{\geq 0}}$ with a subsequence, we may assume that $\lim_{k \rightarrow \infty} u_{m,k} = u_m$ exists for $1 \leq m \leq p$. Then $u_m \in \limsup_k \gamma_m(\epsilon_k) \subset \gamma_m$ and $\|u_m\| = 1$. Given an element $u \in \tau_J$, there exists $a_m \in \mathbb{R}_{\geq 0}^n$ such that $u = \sum_m a_m u_m$. Then $\lim_{k \rightarrow \infty} \sum_m a_m u_{m,k} = u \in \limsup_k \tau_{J(\epsilon_k)}$, as desired. \square

With these lemmas in hand, we now return to our problem. Fix a restricted, weakly α -compatible pair $(\mathcal{Z}, \mathcal{F})$, where $\mathcal{Z} = (Z_s)_{s \in \mathcal{S}}$ and $\mathcal{F} = (F_s)_{s \in \mathcal{S}}$. We may apply Lemma 6.5.6 to both $P(\epsilon) = \text{Newt}(f)_{\mathcal{Z}(\epsilon)}$ and $Q(\epsilon) = Q_{\mathcal{Z}(\epsilon)}$. Hence, by Lemma 6.5.2, we may replace \mathcal{Z} by $\mathcal{Z}(\epsilon)$ so that $\{Q_{\mathcal{Z}(\epsilon)}\}_\epsilon$ and $\{\text{Newt}(f)_{\mathcal{Z}(\epsilon)}\}_\epsilon$ are locally combinatorially constant. Consider a nonempty face J of $Q_{\mathcal{Z}}$, dual to a cone τ_J in $\Sigma_{\mathcal{Z}}$. For ϵ sufficiently small, let $J(\epsilon)$ denote the corresponding nonempty face of $Q_{\mathcal{Z}(\epsilon)}$, dual to the cone $\tau_{J(\epsilon)}$ of $\Sigma_{\mathcal{Z}(\epsilon)}$. Similarly, given a cone C in $\Sigma \cap \Sigma_{\mathcal{Z}}$, we let $C(\epsilon)$ denote the corresponding cone in $\Sigma \cap \Sigma_{\mathcal{Z}(\epsilon)}$. If C is dual to a face of $\text{Newt}(f)_{\mathcal{Z}}$ of the form $K' + J'$ for some $K' \in \Gamma$ and some nonempty face $J' \subset Q_{\mathcal{Z}(\epsilon)}$, then $C(\epsilon) = \sigma_{K'} \cap \tau_{J'(\epsilon)}$ is dual to the face $K' + J'(\epsilon)$ in $\text{Newt}(f)_{\mathcal{Z}(\epsilon)}$.

Remark 6.5.8. Consider a nonempty face J of $Q_{\mathcal{Z}}$ such that $\sigma_M \cap \tau_J \neq \{0\}$. Since $\{Q_{\mathcal{Z}(\epsilon)}\}_\epsilon$ is locally combinatorially constant, for any $s \in \mathcal{S}$, $Z_s \in J$ if and only if $Z_s(\epsilon_s) \in J(\epsilon)$. In particular, if $\sigma_M \cap \tau_{J(\epsilon)} \neq \{0\}$, then $F_J = \max\{F_s : Z_s \in J\} = \max\{F_s : Z_s(\epsilon_s) \in J(\epsilon)\} = F_{J(\epsilon)}$.

The lemma below says that a pair (C, J) not being α -critical is an open condition.

Lemma 6.5.9. *Let J be a nonempty face of $Q_{\mathcal{Z}}$, and consider a cone $C \in \Sigma \cap \Sigma_{\mathcal{Z}}$. Suppose there exists a sequence $\{\epsilon_m\}_{m \in \mathbb{Z}_{>0}}$ such that $\epsilon_m \in \mathbb{Q}^{\mathcal{S}}$, $\lim_{m \rightarrow \infty} \epsilon_m = 0$ and $(C(\epsilon_m), J(\epsilon_m))$ is α -critical for all m . Then (C, J) is α -critical.*

Proof. By Lemma 6.4.8 and since $\{\text{Newt}(f)_{\mathcal{Z}(\epsilon)}\}_\epsilon$ is locally combinatorially constant, $C(\epsilon_m) \subset \tau_{J(\epsilon_m)}$ is dual to a face of the form $K + J'(\epsilon_m)$ of $\text{Newt}(f)_{\mathcal{Z}(\epsilon_m)}$, where $K \in \Gamma$ and J' is a face of $Q_{\mathcal{Z}}$ such that $J(\epsilon_m) \subset J'(\epsilon_m)$, or, equivalently, $J \subset J'$. Then C is dual to $K + J'$. In particular, $C \subset \tau_{J'} \subset \tau_J$. By hypothesis, $C(\epsilon_m) \cap \sigma_M \neq \{0\}$. It follows from Bolzano–Weierstrass and Lemma 6.5.7 that $C \cap \sigma_M \neq \{0\}$. Then $\sigma_M \cap \tau_J \neq \{0\}$, and, by Remark 6.5.8, $F_J = F_{J(\epsilon_m)}$ for all m . The condition $C(\epsilon_m) \subset \sigma_{A_{F_J}}$ implies that $\sigma_K \subset \sigma_{A_{F_J}}$, and hence $C \subset \sigma_{A_{F_J}}$. By Lemma 6.5.7 and since $H_{\alpha A_{F_J} + 1}$ is closed, $C = \limsup_m C(\epsilon_m) \subset H_{\alpha A_{F_J} + 1}$, and hence C is critical with respect to (α, A_{F_J}) . We conclude that (C, J) is α -critical. \square

We say that ϵ can be chosen to be arbitrarily small if for any $\delta > 0$, there exists a choice of ϵ such that $|\epsilon_s| < \delta$ for all $s \in \mathcal{S}$.

Lemma 6.5.10. *Suppose that (C, J) is α -critical and $C \not\subset \sigma_M$. Then there exists an arbitrarily small choice of $\epsilon \in \mathbb{Q}^{\mathcal{S}}$ such that $(C(\epsilon), J(\epsilon))$ is not α -critical.*

Proof. By Lemma 6.4.8, C is dual to a face $K + J'$ of $\text{Newt}(f)_{\mathcal{Z}}$, where $K \in \Gamma$ and J' is a face of $Q_{\mathcal{Z}}$ such that $K \subset F_{J'}$ and $J \subset J'$. Then $C(\epsilon) \subset \tau_{J(\epsilon)}$ is dual to the face $K + J'(\epsilon)$ of $\text{Newt}(f)_{\mathcal{Z}(\epsilon)}$. Note that $C \subset \sigma_{A_{F_J}}$ implies that $\sigma_K \subset \sigma_{A_{F_J}}$, and hence $A_{F_J} \in K$. Since K, M are faces of Γ , $K \cap M$ is a (possibly empty) face of M , and $C_{K \cap M} = C_K \cap C_M$. Let $B_K = \text{Gen}(C_{K \cap M}) \cup \mathcal{A}_M$.

Assume that $\mathbf{1} \in \text{span}(B_K)$. We can write $\mathbf{1} = \sum_{V \in B_K} \lambda_V V$, for some coefficients λ_V , with $\lambda_V = 1$ for all $V \in \mathcal{A}_M$. Applying ψ_M to both sides gives $-\alpha = \sum_{V \in B_K \setminus \text{Unb}(C_{K \cap M})} \lambda_V$. Hence we may equivalently write

$$\alpha A_{F_J} + \mathbf{1} = \sum_{V \in B_K \setminus \text{Unb}(C_{K \cap M})} \lambda_V (V - A_{F_J}) + \sum_{V \in \text{Unb}(C_{K \cap M})} \lambda_V V.$$

Consider $u \in C^\circ \subset \sigma_K^\circ \cap H_{\alpha A_{F_J} + 1}$. Consider $V \in \text{Gen}(C_K)$. If $V \in \text{Unb}(C_K)$, then $u \in \sigma_K^\circ$ implies that $\langle u, V \rangle = 0$. Since $A_{F_J} \in K$, if $V \in \text{Vert}(K)$, then $u \in \sigma_K^\circ$ implies that $\langle u, V - A_{F_J} \rangle = 0$. We compute:

$$\begin{aligned} 0 &= \langle u, \alpha A_{F_J} + \mathbf{1} \rangle = \sum_{V \in B_K \setminus \text{Unb}(C_{K \cap M})} \lambda_V \langle u, V - A_{F_J} \rangle + \sum_{V \in \text{Unb}(C_{K \cap M})} \lambda_V \langle u, V \rangle \\ &= \sum_{V \in B_K \setminus C_K} \langle u, V - A_{F_J} \rangle. \end{aligned}$$

Since each term in the right-hand sum is positive, we deduce that the sum must be empty. It follows that $B_K \subset C_{K \cap M}$ and $\mathbf{1} \in \text{span}(C_{K \cap M})$. Since M is minimal in $\text{Contrib}(\alpha)$, we deduce that $K \cap M = M$. Then $C \subset \sigma_K \subset \sigma_M$, a contradiction.

We conclude that $\mathbf{1} \notin \text{span}(B_K)$. Since $A_{F_J} \in B_K$, it follows that $\alpha A_{F_J} + \mathbf{1} \notin \text{span}(B_K)$. Since $\alpha A_{F_J} + \mathbf{1} \in \text{span}(M) \cap \mathbb{Q}^n$ and $\text{Newt}(f)$ is α -simplicial, it follows that there exists $u' \in \mathbb{Q}^n$ such that

- (1) $\langle u', \alpha A_{F_J} + \mathbf{1} \rangle = 1$,
- (2) $\langle u', V \rangle = 0$ for all elements $V \in B_K$, and
- (3) $\langle u', V \rangle = 0$ for all $V \in \text{Gen}(C_{F_{J'}} \setminus C_M)$.

Consider an element $u \in C^\circ \cap \mathbb{Q}^n = \sigma_K^\circ \cap (\tau_{J'})^\circ \cap \mathbb{Q}^n \subset H_{\alpha A_{F_J} + \mathbf{1}}$, and let $\hat{u}(\lambda) = u + \lambda u'$ for some choice of $\lambda \neq 0 \in \mathbb{R}$. Then property (1) implies that $\langle \hat{u}(\lambda), \alpha A_{F_J} + \mathbf{1} \rangle = \lambda \neq 0$, and hence $\hat{u}(\lambda) \notin H_{\alpha A_{F_J} + \mathbf{1}}$. Properties (2) and (3) imply that for any V in C_K , $\langle \hat{u}(\lambda), V \rangle = \langle u, V \rangle + \lambda \langle u', V \rangle = \langle u, V \rangle$. It follows that $\hat{u}(\lambda) \in \sigma_K^\circ$ provided $|\lambda|$ is sufficiently small.

Recall that $\tau_{J'} = \cap_{Z_s \in J'} \tau_{Z_s}$. Consider $s \in \mathcal{S}$ such that $Z_s \in J'$. We claim that for a generic choice of $\lambda \in \mathbb{Q}$, we may choose $\epsilon_s \in \mathbb{Q}$ such that $\langle \hat{u}(\lambda), Z_s(\epsilon_s) \rangle = \langle u, Z_s \rangle$. Assume this claim holds. It follows that with this choice of $\epsilon = \{\epsilon_s\}_{s \in \mathcal{S}}$, $\hat{u}(\lambda) \in (\tau_{J'}(\epsilon))^\circ$ provided $|\lambda|$ is chosen sufficiently small. Then $\hat{u}(\lambda) \in \sigma_K^\circ \cap (\tau_{J'}(\epsilon))^\circ = C(\epsilon)^\circ$ and $\hat{u}(\lambda) \notin H_{\alpha A_{F_J} + \mathbf{1}}$. Either $\sigma_M \cap \tau_{J(\epsilon)} = \{0\}$, or $\sigma_M \cap \tau_{J(\epsilon)} \neq \{0\}$ and, by Remark 6.5.8, $F_J = F_{J(\epsilon)}$ and $C(\epsilon) \not\subset H_{\alpha A_{F_{J(\epsilon)}} + \mathbf{1}}$. In either case, $(C(\epsilon), J(\epsilon))$ is not α -critical.

It remains to verify the claim. We compute:

$$\begin{aligned} \langle \hat{u}(\lambda), Z_s(\epsilon_s) \rangle &= \langle \hat{u}(\lambda), Z_s + \epsilon_s V_M \rangle \\ &= \langle u, Z_s \rangle + \lambda \langle u', Z_s \rangle + \epsilon_s (\langle u, V_M \rangle + \lambda \langle u', V_M \rangle). \end{aligned}$$

Assume that $\langle u', V_M \rangle \neq 0$. Then for $\lambda \neq -\frac{\langle u, V_M \rangle}{\langle u', V_M \rangle}$, we may set $\epsilon_s = -\frac{\lambda \langle u', Z_s \rangle}{\langle u, V_M \rangle + \lambda \langle u', V_M \rangle}$, and the above calculation shows that $\langle \hat{u}(\lambda), Z_s(\epsilon_s) \rangle = \langle u, Z_s \rangle$. Assume that $\langle u', V_M \rangle = 0$, and let $\epsilon_s = 0$. Since $(\mathcal{Z}, \mathcal{F})$ is restricted, $Z_s \in \text{span}(\{V_M\} \cup \text{Gen}(C_{F_s} \setminus C_M) \cup \mathcal{A}_M)$ for every $s \in \mathcal{S}$. Since $Z_s \in J'$, Definition 6.4.3 implies that $F_s \subset F_{J'}$. Then properties (2) and (3) imply that $\langle u', Z_s \rangle = 0$, and the above calculation shows that $\langle \hat{u}(\lambda), Z_s(\epsilon_s) \rangle = \langle u, Z_s \rangle$. \square

Lemma 6.5.11. *Suppose there exists a restricted, weakly α -compatible pair $(\mathcal{Z}, \mathcal{F})$. Then there exists an α -compatible pair $(\mathcal{Z}, \mathcal{F})$.*

Proof. Consider the restricted, weakly α -compatible pair $(\mathcal{Z}, \mathcal{F})$ above. Suppose $(\mathcal{Z}, \mathcal{F})$ is not α -compatible. That is, suppose there exists a pair (C, J) that is α -critical and $C \not\subset \sigma_M$. Then Lemma 6.5.9 and Lemma 6.5.10 imply that we can deform $(\mathcal{Z}, \mathcal{F})$ and strictly increase the number of pairs (C, J) that do not have α -critical intersection. Since there are finitely many such pairs, by repeating this procedure we obtain an α -compatible pair. \square

6.6. Existence of restricted, weakly α -compatible sets. In this section, we use the existence of a locally unique labeling to explicitly construct a restricted, weakly α -compatible pair. Recall that $\text{Gen}(C_F) = \text{Vert}(F) \cup \text{Unb}(C_F)$ is the set of distinguished vertices on the rays of C_F . Recall that because $\text{Newt}(f)$ is α -simplicial, there is a bijection between $\{K \in \Gamma : M \subset K \subset F\}$ and subsets of $\text{Gen}(C_F \setminus C_M) = \text{Gen}(C_F) \setminus \text{Gen}(C_M)$.

Consider an element $F \in \text{Contrib}(\alpha)_M$. Given an element V in $\text{Gen}(C_F)$, let $\zeta(V) \in F \subset \Gamma$ be defined by

$$\zeta(V) := \begin{cases} V & \text{if } V \in \text{Vert}(F), \\ V + V_M & \text{if } V \in \text{Unb}(C_F). \end{cases}$$

Lemma 6.6.1. *Suppose that $F, F' \in \text{Contrib}(\alpha)_M$ and $V \in \text{Gen}(C_F)$. Then $\zeta(V) \in F'$ if and only if $V \in \text{Gen}(C_{F'})$.*

Proof. First, suppose that $V \in \text{Vert}(F)$. Then $\zeta(V) = V \in F'$ if and only if $V \in \text{Vert}(F')$. Second, suppose that $V \in \text{Unb}(C_F)$. Consider $u \in \sigma_{F'}$. Then $\langle u, V \rangle = \langle u, \zeta(V) - V_M \rangle$, so $\zeta(V) \in F'$ if and only if $\langle u, \zeta(V) - V_M \rangle = 0$, if and only if $\langle u, V \rangle = 0$, if and only if $V \in \text{Unb}(C_{F'})$. The result follows. \square

Let \mathcal{S} be the set of saturated chains of faces in Γ starting at M , i.e., a chain of faces starting at M where the dimension increases by one at each step. Let $s = F_\bullet$ be an element of \mathcal{S} . Let ℓ_s denote the length of F_\bullet , i.e., the number of elements in F_\bullet minus one. We let $F_{\bullet,i}$ denote the i th element of F_\bullet for $0 \leq i \leq \ell_s$. For example, $F_{\bullet,0} = M$. We write $F \in F_\bullet$ if $F = F_{\bullet,i}$ for some $0 \leq i \leq \ell_s$.

Define $V_{s,0} = V_M$. Since F_\bullet is saturated and $\text{Newt}(f)$ is α -simplicial, we may define $V_{s,i}$ to be the unique element of $\text{Gen}(C_{F_{\bullet,i}} \setminus C_{F_{\bullet,i-1}})$ for $1 \leq i \leq \ell_s$.

Definition 6.6.2. *Let \mathcal{S} be the set of saturated chains of faces in Γ starting at M . We define a pair $(\mathcal{Z}, \mathcal{F})$, where $\mathcal{Z} = (Z_s)_{s \in \mathcal{S}}$ and $\mathcal{F} = (F_s)_{s \in \mathcal{S}}$ are collections of elements of \mathbb{Q}^n and $\text{Contrib}(\alpha)_M$ respectively, as follows: for any element $s = F_\bullet$ of \mathcal{S} , let*

$$Z_s := \sum_{i=0}^{\ell_s} b_{i,\ell_s} \zeta(V_{s,i}),$$

where $\{b_{i,j}\}_{0 \leq i,j \leq r}$, $r = n - 1 - \dim M$, and

$$b_{i,j} = b_{i,j}(\mu) = \begin{cases} 2^{-i} - 2i\mu, & \text{if } i = j \\ 2^{-(i+1)} - (i+j)\mu, & \text{if } i < j \\ 0, & \text{otherwise} \end{cases}$$

for some $\mu \in \mathbb{Q}$ such that $0 < \mu \ll 1$. Let $F_s := F_{\bullet,\ell_s}$ be the maximal element of F_\bullet .

For example, $b_{0,0} = 1$ and if $s = F_\bullet$, where $F_\bullet = \{M\}$, then $\ell_s = 0$, $Z_s = V_M$, and $F_s = M$. Note that we abuse notation above by not indicating the dependence of $(\mathcal{Z}, \mathcal{F})$ on the choice of μ . Below we fix a value of μ sufficiently small. Our goal is to show that we can construct a restricted, weakly α -compatible pair from $(\mathcal{Z}, \mathcal{F})$. Recall that we have fixed a locally unique labeling (A_F, e_F^*) of $\text{Contrib}(\alpha)_M$.

Definition 6.6.3. *Let $s = F_\bullet \in \mathcal{S}$. Let $\mathcal{A}_s = \{A_F : F \in F_\bullet\}$. Given an element A in \mathcal{A}_s , we define a base direction $e_{s,A}^*$ as follows: if $A = A_F$ for some $F \in F_\bullet$, then $e_{s,A}^* := e_F^*$. We define a linear function $\Phi_s : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by*

$$\Phi_s(X) = X - \sum_{A \in \mathcal{A}_s} \langle e_{s,A}^*, X \rangle (A - V_M).$$

The fact that $e_{s,A}^*$ is well-defined in Definition 6.6.3 is an immediate consequence of the locally unique labeling condition. Explicitly, if $A = A_F = A_{F'}$ for some $F, F' \in F_\bullet$, then either $F \subset F'$ or $F' \subset F$, and $(*)$ implies that $e_F^* = e_{F'}^*$.

Remark 6.6.4. For $s \in \mathcal{S}$, $u \in \sigma_M$ and $X \in \mathbb{R}^n$, $\langle u, \Phi_s(X) \rangle = \langle u, X \rangle$.

Lemma 6.6.5. *Let $s = F_\bullet \in \mathcal{S}$. Then $\langle e_F^*, \Phi_s(X) \rangle = 0$ for any $F \in F_\bullet$ and any $X \in \mathbb{R}^n$. If $F_s \subset K$ and $\langle e_K^*, X \rangle = 0$ for some $X \in \mathbb{R}^n$, then $\langle e_K^*, \Phi_s(X) \rangle = 0$.*

Proof. Recall from (24) that $\langle e_F^*, V_M \rangle = 0$ for all $F \supset M$. Suppose that $F \in F_\bullet$. Then $e_{s,A_F}^* = e_F^*$, and we compute $\langle e_F^*, \Phi_s(X) \rangle = \langle e_F^*, X - \langle e_{s,A_F}^*, X \rangle (A_F - V_M) \rangle = 0$. Suppose that $F_s \subset K$ and $\langle e_K^*, X \rangle = 0$. If $\langle e_K^*, A \rangle = 0$ for all $A \in \mathcal{A}_s$, then $\langle e_K^*, \Phi_s(X) \rangle = \langle e_K^*, X \rangle = 0$. Suppose that $\langle e_K^*, A_F \rangle \neq 0$ for some $F \in F_\bullet$. Then $A_K = A_F$. Since $F \subset K$, the locally unique labeling condition $(*)$ implies that $e_{s,A_F}^* = e_F^* = e_K^*$. As above, $\langle e_K^*, \Phi_s(X) \rangle = \langle e_K^*, X - \langle e_{s,A_F}^*, X \rangle (A_F - V_M) \rangle = 0$. \square

Lemma 6.6.6. *With the notation of Definition 6.6.2, suppose that $(\mathcal{Z}, \mathcal{F})$ satisfies the the following property: Suppose that $\sigma_K^\circ \cap \tau_{Z_s} \cap \tau_{Z_{s'}} \neq \emptyset$, for some $K \in \text{Contrib}(\alpha)_M$ and $s = F_\bullet, s' = F'_\bullet \in \mathcal{S}$. Then*

- (1) $K \subset F_s$, and
- (2) either $F_s \in F'_\bullet$ or $F_{s'} \in F_\bullet$.

Let $\Phi(\mathcal{Z}) := (\Phi_s(Z_s))_{s \in \mathcal{S}}$. Then $(\Phi(\mathcal{Z}), \mathcal{F})$ is restricted and weakly α -compatible.

Proof. It follows from Definition 6.6.2 and Definition 6.6.3 that $(\Phi(\mathcal{Z}), \mathcal{F})$ is restricted. Suppose that $\sigma_K^\circ \cap \tau_{\Phi_s(Z_s)} \cap \tau_{\Phi_{s'}(Z_{s'})} \neq \emptyset$, for some $K \in \text{Contrib}(\alpha)_M$ and $s = F_\bullet, s' = F'_\bullet \in \mathcal{S}$. By Remark 6.6.4, the restriction of $\Sigma_{\mathcal{Z}}$ to σ_M equals the restriction of $\Sigma_{\Phi(\mathcal{Z})}$ to σ_M . Hence $\sigma_K^\circ \cap \tau_{Z_s} \cap \tau_{Z_{s'}} \neq \emptyset$. We deduce that $K \subset F_s$, and, either $F_s \in F'_\bullet$ or $F_{s'} \in F_\bullet$. The latter condition implies that either $F_s \subset F_{s'}$ or $F_{s'} \subset F_s$.

Applying Lemma 6.6.5 with $s = F_\bullet$ and $F = F_s$, gives $\langle e_{F_s}^*, \Phi_s(Z_s) \rangle = 0$. It remains to show that $\langle e_{F_s}^*, \Phi_{s'}(Z_{s'}) \rangle = 0$. Suppose that $F_s \in F'_\bullet$. Applying Lemma 6.6.5 with $s' = F'_\bullet$ and $F = F_s$, gives $\langle e_{F_s}^*, \Phi_{s'}(Z_{s'}) \rangle = 0$, as desired. Suppose that $F_{s'} \in F_\bullet$. Then $F_{s'} \subset F_s$. Applying Lemma 6.6.5 with $s' = F'_\bullet$ and $K = F_s$, gives $\langle e_{F_s}^*, \Phi_{s'}(Z_{s'}) \rangle = 0$, provided $\langle e_{F_s}^*, Z_{s'} \rangle = 0$. By Definition 6.6.2, $Z_{s'} \in \text{span}(\{V_M\} \cup \text{Gen}(C_{F_{s'}} \setminus C_M))$. Since $F_{s'} \subset F_s$, F_s is a B_1 -face with base direction $e_{F_s}^*$, and $\langle e_{F_s}^*, V_M \rangle = 0$ by (24), it follows from Remark 6.2.5 that $\langle e_{F_s}^*, Z_{s'} \rangle = 0$, as desired. \square

It remains to show that $(\mathcal{Z}, \mathcal{F})$ satisfies conditions (1) and (2) in Lemma 6.6.6. We will prove this through a series of lemmas.

Lemma 6.6.7. *There exists a constant $\lambda_M > 0$ such that for any $K, F, F' \in \text{Contrib}(\alpha)_M$ that are not subfaces of a common face in $\text{Contrib}(\alpha)_M$, and for any nonzero $u \in \sigma_K$, there exists an element $V \in \text{Gen}(C_F \setminus C_M) \cup \text{Gen}(C_{F'} \setminus C_M)$ such that $\langle u, \zeta(V) \rangle / N(u) \geq 1 + \lambda_M$.*

Proof. Fix $K, F, F' \in \text{Contrib}(\alpha)_M$ that are not subfaces of a common face in $\text{Contrib}(\alpha)_M$. Let $\mathcal{V} = \text{Gen}(C_F \setminus C_M) \cup \text{Gen}(C_{F'} \setminus C_M)$, and consider the continuous function $\phi: \sigma_K \setminus \{0\} \rightarrow \mathbb{R}$ defined by

$$\phi(u) = \left(\max_{V \in \mathcal{V}} \langle u, \zeta(V) \rangle / N(u) \right) - 1.$$

We claim that image satisfies $\text{im}(\phi) \subset \mathbb{R}_{>0}$. Indeed, suppose $\phi(u) = 0$. Let F_u be the face of $\partial \text{Newt}(f)$ minimized by u . Then F_u contains K and $\{\zeta(V) : V \in \mathcal{V}\}$. By Lemma 6.6.1, C_{F_u} contains C_K, C_F and $C_{F'}$. Then K, F, F' are common subfaces of F_u , a contradiction.

Note that $\phi(\eta u) = \phi(u)$ for all $\eta \in \mathbb{R}_{>0}$ and $u \in \sigma_K \setminus \{0\}$. Since $\sigma_K \cap S$ is compact, there exists $\lambda = \lambda(K, F, F') > 0$ such that $\phi(\sigma_K \cap S) \subset [\lambda, \infty)$. We let λ_M be the minimum value of $\lambda(K, F, F')$ over the finitely many choices of K, F, F' . \square

Below we fix $\lambda_M > 0$ satisfying Lemma 6.6.7.

Lemma 6.6.8. *Let $r = n - 1 - \dim M$. Assume that $\mu > 0$ is chosen sufficiently small. Then the coefficients $\{b_{i,j} = b_{i,j}(\mu)\}_{0 \leq i, j \leq r}$ satisfy the following properties:*

- (1) $b_{i,j} > 0$ for $i \leq j$,
- (2) $b_{i,j} > b_{i,j+1}$ for $i \leq j < r$,
- (3) $b_{i,j} > b_{i+1,j}$ for $i < j$
- (4) $\sum_{i \geq k} b_{i,j} > \sum_{i \geq k} b_{i,j+1}$, for any $1 \leq k \leq j < r$,
- (5) $\sum_{i \geq 0} b_{i,r} + b_{r,r} \lambda_M > 1$.

Proof. We check the conditions hold by direct computation, for μ sufficiently small. Condition (1) is clear. For condition (2), we compute, for $i = j < r$,

$$b_{i,i} = 2^{-i} - 2i\mu > b_{i,i+1} = 2^{-(i+1)} - (2i+1)\mu,$$

and, for $i < j < r$,

$$b_{i,j} = 2^{-(i+1)} - (i+j)\mu > b_{i,j+1} = 2^{-(i+1)} - (i+j+1)\mu.$$

For condition (3), we compute, for $i+1 < j$,

$$b_{i,j} = 2^{-(i+1)} - (i+j)\mu > b_{i+1,j} = 2^{-(i+2)} - (i+j+1)\mu,$$

and, for $i+1 = j$,

$$b_{i,i+1} = 2^{-(i+1)} - (2i+1)\mu > b_{i+1,i+1} = 2^{-(i+1)} - (2i+2)\mu.$$

For condition (4), define $c_{k,j} = \sum_{i \geq k} b_{i,j}$ for $k \leq j$. Then

$$\begin{aligned} c_{k,j} &= 2^{-k} - \sum_{i=k}^j (i+j)\mu \\ &= 2^{-k} - (j-k+1)(3j+k)\mu/2. \end{aligned}$$

For $j < r$, $c_{k,j} > c_{k,j+1}$, as desired. For condition (5), we compute

$$\begin{aligned} \sum_{i \geq 0} b_{i,r} + b_{r,r}\lambda_M - 1 &= c_{0,r} + b_{r,r}\lambda_M - 1 \\ &= -3r(r+1)\mu/2 + b_{r,r}\lambda_M \\ &= -3r(r+1)\mu/2 + (2^{-r} - 2r\mu)\lambda_M \\ &= 2^{-r}\lambda_M - \mu r(3(r+1)/2 + 2\lambda_M). \end{aligned}$$

The latter expression is positive for μ sufficiently small. \square

Lemma 6.6.9. *Let $s = F_\bullet \in \mathcal{S}$ and suppose $u \in \sigma_M \cap \tau_{Z_s}$ is nonzero. Then $\langle u, \zeta(V_{s,i}) \rangle \leq \langle u, \zeta(V_{s,j}) \rangle$ for any $0 \leq i \leq j \leq \ell_s$. Moreover, if $F_s \subset F \in \Gamma$, then there exists a constant $0 \leq m < \lambda_M$ such that $\text{Gen}(C_{F_s} \setminus C_M) = \{V \in \text{Gen}(C_F \setminus C_M) : \langle u, \zeta(V) \rangle / N(u) \leq 1 + m\}$.*

Proof. Since $u \in \sigma_M$, $\zeta(V_{s,0}) = V_M \in M$, and $\zeta(V_{s,j}) \in \text{Newt}(f)$, it follows that $\langle u, \zeta(V_{s,0}) \rangle \leq \langle u, \zeta(V_{s,j}) \rangle$ for any $0 \leq j \leq \ell_s$. Suppose that $\langle u, \zeta(V_{s,i}) \rangle > \langle u, \zeta(V_{s,j}) \rangle$ for some $0 < i < j \leq \ell_s$. Let $\pi = (i, j) \in \text{Sym}_{\ell_s}$ be the permutation of $[\ell_s]$ switching i and j . Let $\pi(s)$ be the unique element in \mathcal{S} such that $\ell_{\pi(s)} = \ell_s$ and $V_{\pi(s),i} = V_{s,\pi(i)}$ for $1 \leq i \leq \ell_s$. Using (3) in Lemma 6.6.8, we compute:

$$\begin{aligned} \langle u, Z_s - Z_{\pi(s)} \rangle &= b_{i,\ell_s} \langle u, \zeta(V_{s,i}) - \zeta(V_{\pi(s),i}) \rangle + b_{j,\ell_s} \langle u, \zeta(V_{s,j}) - \zeta(V_{\pi(s),j}) \rangle \\ &= (b_{i,\ell_s} - b_{j,\ell_s}) \langle u, \zeta(V_{s,i}) - \zeta(V_{s,j}) \rangle > 0. \end{aligned}$$

The latter contradicts the assumption that $u \in \tau_{Z_s}$. This completes the proof of the first statement.

Since M is interior and $u \in \sigma_M$, $N(u) > 0$. Let $m = (\langle u, \zeta(V_{s,\ell_s}) \rangle / N(u)) - 1 \geq 0$. Assume that $m \geq \lambda_M$. Using all the statements of Lemma 6.6.8, we compute

$$\begin{aligned} \langle u, Z_s \rangle / N(u) &= \sum_{i=0}^{\ell_s} b_{i,\ell_s} \langle u, \zeta(V_{s,i}) \rangle / N(u) \\ &\geq \sum_{i=0}^{\ell_s} b_{i,\ell_s} + b_{\ell_s,\ell_s} \lambda_M \\ &\geq \sum_{i=0}^r b_{i,r} + b_{r,r} \lambda_M > 1. \end{aligned}$$

On the other hand, if \tilde{s} is the unique element in \mathcal{S} with $\ell_{\tilde{s}} = 0$, then, since $u \in \sigma_M$,

$$\langle u, Z_{\tilde{s}} \rangle / N(u) = \langle u, V_M \rangle / N(u) = 1 < \langle u, Z_s \rangle / N(u).$$

This contradicts the assumption that $u \in \tau_{Z_s}$. We conclude that $m < \lambda_M$.

Since $\langle u, \zeta(V_{s, \ell_s}) \rangle = \max_{0 \leq i \leq \ell_s} \langle u, \zeta(V_{s, i}) \rangle$, we have

$$\text{Gen}(C_{F_s} \setminus C_M) \subset \{V \in \text{Gen}(C_F \setminus C_M) : \langle u, \zeta(V) \rangle / N(u) \leq 1 + m\}.$$

It remains to prove the reverse inclusion. Suppose that $V \in \text{Gen}(C_F \setminus C_M)$ and $\langle u, \zeta(V) \rangle / N(u) \leq 1 + m$. Equivalently, we assume that $\langle u, \zeta(V_{s, \ell_s}) - \zeta(V) \rangle \geq 0$. We argue by contradiction. Assume that $V \notin C_{F_s}$. Let s' be the unique element in \mathcal{S} such that $\ell_{s'} = \ell_s + 1$, $V_{s', i} = V_{s, i}$ for $0 \leq i \leq \ell_s$, and $V_{s', \ell_{s'}} = V$. If we let $V_{s', -1} = V_{s', -1} = 0$, then we can write

$$\begin{aligned} \langle u, Z_s \rangle &= \sum_{k=0}^{\ell_s} b_{k, \ell_s} \langle u, \zeta(V_{s, k}) \rangle = \sum_{k=0}^{\ell_s} \left(\sum_{i=k}^{\ell_s} b_{i, \ell_s} \right) \langle u, \zeta(V_{s, k}) - \zeta(V_{s, k-1}) \rangle, \\ \langle u, Z_{s'} \rangle &= \sum_{k=0}^{\ell_s+1} b_{k, \ell_s+1} \langle u, \zeta(V_{s', k}) \rangle = \sum_{k=0}^{\ell_s+1} \left(\sum_{i=k}^{\ell_s+1} b_{i, \ell_s+1} \right) \langle u, \zeta(V_{s', k}) - \zeta(V_{s', k-1}) \rangle, \text{ and} \\ \langle u, Z_s - Z_{s'} \rangle &= \sum_{k=0}^{\ell_s} \left(\sum_{i=k}^{\ell_s} b_{i, \ell_s} - \sum_{i=k}^{\ell_s+1} b_{i, \ell_s+1} \right) \langle u, \zeta(V_{s, k}) - \zeta(V_{s, k-1}) \rangle + b_{\ell_s+1, \ell_s+1} \langle u, \zeta(V_{s, \ell_s}) - \zeta(V) \rangle. \end{aligned}$$

Since $u \in \sigma_M$ and $V_{s, 0} = V_M \in M$, we have $\langle u, \zeta(V_{s, 0}) \rangle = N(u) > 0$. By conditions (1) and (4) in Lemma 6.6.8, it follows that all terms above are nonnegative, and at least one term is positive. This contradicts the assumption that $u \in \tau_{Z_s}$. \square

The following lemma completes our proof.

Lemma 6.6.10. *With the notation of Definition 6.6.2, $(\mathcal{Z}, \mathcal{F})$ satisfies the the following property:*

Suppose that $\sigma_K^\circ \cap \tau_{Z_s} \cap \tau_{Z_{s'}} \neq \emptyset$, for some $K \in \text{Contrib}(\alpha)_M$ and $s = F_\bullet, s' = F'_\bullet \in \mathcal{S}$. Then

- (1) $K \subset F_s$, and
- (2) either $F_s \in F'_\bullet$ or $F_{s'} \in F_\bullet$.

Proof. Fix $u \in \sigma_K^\circ \cap \tau_{Z_s} \cap \tau_{Z_{s'}}$. By Lemma 6.6.9, $\langle u, \zeta(V) \rangle / N(u) < 1 + \lambda_M$ for all $V \in \text{Gen}(C_{F_s} \setminus C_M) \cup \text{Gen}(C_{F_{s'}} \setminus C_M)$. By Lemma 6.6.7, there exists a face $F \in \text{Contrib}(\alpha)$ such that $K, F_s, F_{s'} \subset F$. By Lemma 6.6.9, there exists $m, m' \geq 0$ such that

$$\text{Gen}(C_{F_s} \setminus C_M) = \{V \in \text{Gen}(C_F \setminus C_M) : \langle u, \zeta(V) \rangle / N(u) \leq 1 + m\}, \text{ and}$$

$$\text{Gen}(C_{F_{s'}} \setminus C_M) = \{V \in \text{Gen}(C_F \setminus C_M) : \langle u, \zeta(V) \rangle / N(u) \leq 1 + m'\}.$$

Since $u \in \sigma_K^\circ$, it follows from Lemma 6.6.1 that $\text{Gen}(C_K \setminus C_M) = \{V \in \text{Gen}(C_F \setminus C_M) : \langle u, \zeta(V) \rangle / N(u) = 1\}$, and hence $K \subset F_s$. It remains to establish (2). Without loss of generality, we may assume that $m \leq m'$. Then $\text{Gen}(C_{F_s} \setminus C_M) = \{V \in \text{Gen}(C_{F_{s'}} \setminus C_M) : \langle u, \zeta(V) \rangle / N(u) \leq 1 + m\}$. By Lemma 6.6.9, $\langle u, \zeta(V_{s', i}) \rangle \leq \langle u, \zeta(V_{s', j}) \rangle$ for $0 \leq i \leq j \leq \ell_{s'}$. We deduce that $F_s \in F'_\bullet$. \square

A corollary of the proof of Lemma 6.6.10 is that $K \in F_\bullet$.

Proof of Theorem 6.4.12. Let $\alpha \notin \mathbb{Z}$, and assume that all faces of $\text{Contrib}(\alpha)$ are UB_1 and $\text{Newt}(f)$ is α -simplicial. Let M be a minimal face of $\text{Contrib}(\alpha)$. Then Lemma 6.6.6 and Lemma 6.6.10 imply that there is a restricted weakly α -compatible pair. Lemma 6.5.11 then implies that there is an α -compatible pair. \square

7. BEYOND THE SIMPLICIAL CASE

Our techniques are capable of proving the local motivic monodromy conjecture for certain nondegenerate singularities whose Newton polyhedra are not simplicial. In this section, we prove our strongest result on the local motivic monodromy conjecture, explain the remaining cases needed to prove the local motivic monodromy conjecture for nondegenerate singularities, and prove the local motivic monodromy conjecture for 3-dimensional nondegenerate singularities.

7.1. Local motivic monodromy conjecture. We first state our strongest theorem on the local motivic monodromy conjecture for nondegenerate singularities. Recall that given $\beta \in \mathbb{Q}$, $D(\beta) \in \mathbb{Z}_{>0}$ is the denominator of β , written as a reduced fraction.

Theorem 7.1.1. *Suppose f is nondegenerate, and suppose that, for every $\alpha \in \mathbb{Q} \setminus \mathbb{Z}$, either:*

- (1) *Newt(f) has $D(\alpha)$ -good projection and there is a face in $\text{Contrib}(\alpha)$ that is not pseudo- UB_1 ,*
- (2) *Newt(f) is α -simplicial and every face of $\text{Contrib}(\alpha)$ is UB_1 , or*
- (3) *there is $\beta \in \mathbb{Q}$ with $D(\alpha)$ dividing $D(\beta)$ and a face F of $\text{Contrib}(\beta)$ with $|\text{Unb}(C_F)| = n - 1$.*

Then there is a set of candidate poles $\mathcal{P} \subset \mathbb{Q}$ for $Z_{\text{mot}}(T)$ such that for all $\alpha \in \mathcal{P}$, $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy.

To complete the proof of Theorem 7.1.1 we need the following lemma.

Lemma 7.1.2. *Let $\alpha \in \mathbb{Q}$. Suppose there is $\beta \in \mathbb{Q}$ with $D(\alpha)$ dividing $D(\beta)$ and a face $F \in \text{Contrib}(\beta)$ with $|\text{Unb}(C_F)| = n - 1$. Then $\exp(2\pi i \alpha)$ is a nearby eigenvalue of monodromy.*

Proof. Let F be a face in $\text{Contrib}(\beta)$ with $|\text{Unb}(C_F)| = n - 1$. Recall that we may write $\langle \text{Unb}(C_F) \rangle = \mathbb{R}_{\geq 0}^{I_F}$ for some $I_F \subset [n]$, and \bar{F} denotes the image of F under the projection $\mathbb{R}^n \rightarrow \mathbb{R}^n / \langle \text{Unb}(C_F) \rangle$. Then $\bar{F} = \{\rho_F\} \subset \mathbb{R}$, where ρ_F is the lattice distance of F to the origin. Observe that $\beta = -1/\rho_F$ and $D(\beta) = \rho_F$. Let $x = x_{I_F}$ be a general point in $\mathbb{A}^{I_F} \subset X_f$. By either Varchenko's theorem (see (10)) or Theorem 3.2.1, we compute

$$\tilde{E}(\mathcal{F}_x) + 1 = \sum_{i=0}^{\rho_F-1} [i/D(\beta)]. \quad \square$$

Proof of Theorem 7.1.1. The result is an immediate consequence of Theorem 3.4.6, Theorem 6.1.2 and Lemma 7.1.2. □

On the other hand, there are three major classes of Newton polyhedra that Theorem 7.1.1 does not cover.

- (1) All faces of $\text{Contrib}(\alpha)$ are UB_1 , but $\text{Newt}(f)$ is not α -simplicial.
- (2) Every face of $\text{Contrib}(\alpha)$ is pseudo- UB_1 , and at least one face of $\text{Contrib}(\alpha)$ is not UB_1 .
- (3) There is a face of $\text{Contrib}(\alpha)$ that is not UB_1 , and $\text{Newt}(f)$ does not have $D(\alpha)$ -good projection.

For (1), see [ELT22, Theorem 4.3] and [Que, Theorem A] for results that produce “fake poles” of topological and naive motivic zeta function under certain conditions but without an α -simplicial assumption. For (2), a B_2 facet in the sense of [ELT22, Definition 3.9] is pseudo- UB_1 but not UB_1 . It is known that B_2 facets sometimes do not give rise to poles of the local topological zeta function, see, e.g., [ELT22, Proposition 3.11]. For (3), see [Est21] for one approach to proving that $\exp(2\pi i \alpha)$ is an eigenvalue of monodromy in this situation. See Example 2.2.4 and Example 3.2.5 for explicit examples. Note that (3) does not occur when $\text{Newt}(f)$ is convenient.

7.2. Dimension 3 case. We now use Theorem 7.1.1 to deduce the local motivic monodromy conjecture for nondegenerate singularities when $n = 3$ and prove the theorem below. See Section 1.3.1 for a history of prior results on monodromy conjectures for nondegenerate singularities when $n = 3$.

Theorem 7.2.1. *Suppose that f is a nondegenerate polynomial in three variables. Then there is a set of candidate poles $\mathcal{P} \subset \mathbb{Q}$ for $Z_{\text{mot}}(T)$ such that for all $\alpha \in \mathcal{P}$, $\exp(2\pi i\alpha)$ is a nearby eigenvalue of monodromy.*

Lemma 7.2.2. *Suppose $n = 3$ and F is a B_1 -face. Then C_F is simplicial.*

Proof. Let A be an apex with base direction e_ℓ^* . Then $C_F \cap \{e_\ell^* = 0\}$ is a polyhedral cone of dimension at most 2, and hence is simplicial. Since C_F is spanned by $C_F \cap \{e_\ell^* = 0\}$ and the ray through A , it follows that C_F is simplicial. \square

Proof of Theorem 7.2.1. Let $\alpha \in \mathbb{Q}$ be a candidate pole. If $\alpha \in \mathbb{Z}$, then 1 is an eigenvalue of monodromy for $H^0(\mathcal{F}_0, \mathbb{C})$. Hence, we may assume that $\alpha \notin \mathbb{Z}$. Similarly, we may assume that X_f is not smooth at the origin, else $\{-1\}$ is a set of candidate poles for $Z_{\text{mot}}(T)$. We show that if $\text{Newt}(f)$ does not satisfy condition (1) or (3) of Theorem 7.1.1, then it satisfies condition (2). By Lemma 7.1.2, we may therefore assume that, for all β with $D(\alpha)$ dividing $D(\beta)$ and all $F \in \text{Contrib}(\beta)$, $|\text{Unb}(C_F)| \leq 1$. Then $\text{Newt}(f)$ has $D(\alpha)$ -good projection, so we may assume that for every face F of $\text{Contrib}(\alpha)$, F is pseudo- UB_1 .

We now argue that every face F of $\text{Contrib}(\alpha)$ is UB_1 . Then by Lemma 7.2.2, $\text{Newt}(f)$ is α -simplicial and we have verified that condition (2) of Theorem 7.1.1 is satisfied.

If $\dim F \leq 1$, then F is simplicial and hence is UB_1 . First suppose that $\dim F = 2$ and F is compact. Choose two vertices w_1 and w_2 that lie on a 2-dimensional face of $\mathbb{R}_{\geq 0}^3$ (if they do not exist, then no triangulation contains any UB_1 -facet), say $\{e_1^* = 0\}$. Observe that for $j \in \{2, 3\}$, either $\langle e_j^*, w_1 \rangle > 0$ or $\langle e_j^*, w_2 \rangle > 0$, else one of w_1 or w_2 would be in the upper convex hull of the other.

If F is not UB_1 , there are at least two other vertices, w_3 and w_4 . We now consider two cases.

- (1) First consider the case when w_3 and w_4 both have $\langle e_1^*, w_i \rangle = 1$. Consider the 2-dimensional UB_1 lattice simplex with vertices w_1, w_3, w_4 . We may assume that there is an apex with base direction e_2^* . If the apex is w_1 , then w_3 and w_4 must be of the form $(1, 0, c)$ for some $c \in \mathbb{N}$. But then one of w_3 or w_4 is in the upper convex hull of the other. Hence we may assume that the apex is w_3 . Then, $w_1 = (0, 0, a)$ and $w_3 = (1, 1, b)$ for some $a, b \in \mathbb{N}$. Note that $\langle e_1^*, w_2 \rangle = 0$ implies that $\langle e_2^*, w_2 \rangle > 0$. Now consider the 2-dimensional UB_1 lattice simplex with vertices w_2, w_3, w_4 . This has an apex at height 1 with base direction e_3^* . As above, the apex can not be w_2 , else one of w_3 or w_4 is in the upper convex hull of the other. It also can not be w_3 , else $b = 1$ and $\alpha = -1 \in \mathbb{Z}$. Hence $b = 0$ and the apex is w_4 . Since w_1, w_2, w_3, w_4 all lie on F , we deduce that

$$w_1 = (0, 0, a), w_2 = (0, a, 0), w_3 = (1, 1, 0), w_4 = (1, 0, 1),$$

for some $a \in \mathbb{Z}_{>0}$. Note that $a > 1$, else X_f is smooth at the origin. Then $w = ((a-1)/a)w_1 + (1/a)w_2 = (0, 1, a-1)$ is a lattice point in F , and the 2-dimensional lattice simplex with vertices w, w_3, w_4 is not UB_1 , a contradiction.

- (2) Now suppose that there is some vertex w_3 such that $\langle e_1^*, w_3 \rangle > 1$. As w_1, w_2, w_3 span a 2-dimensional UB_1 lattice simplex and either $\langle e_2^*, w_1 \rangle > 0$ or $\langle e_2^*, w_2 \rangle > 0$, we may assume that w_1 is an apex with base direction e_2^* and $\langle e_2^*, w_1 \rangle = 1$. The fourth vertex w_4 must have $\langle e_2^*, w_4 \rangle > 1$, as otherwise we would be in the previous case. Consider the 2-dimensional UB_1 lattice simplex with vertices w_2, w_3, w_4 . It cannot have an apex at height 1 in either the directions e_1^* or e_3^* . Then we must have $w_2 = e_3$, which implies that X_f is smooth at the origin, a contradiction.

Now suppose that $\dim F = 2$ and $|\text{Unb}(C_F)| = 1$. Then C_F has good projection and \bar{F} is UB_1 . By Remark 3.4.4, F is UB_1 . \square

REFERENCES

- [A’C75] N. A’Campo, *La fonction zêta d’une monodromie*, Comment. Math. Helv. **50** (1975), 233–248.
- [Ath12a] C. Athanasiadis, *Cubical subdivisions and local h -vectors*, Ann. Comb. **16** (2012), no. 3, 421–448.
- [Ath12b] ———, *Flag subdivisions and γ -vectors*, Pacific J. Math. **259** (2012), no. 2, 257–278.
- [BN20] E. Bultot and J. Nicaise, *Computing motivic zeta functions on log smooth models*, Math. Z. **295** (2020), no. 1-2, 427–462.
- [BO16] S. Brzostowski and G. Oleksik, *On combinatorial criteria for non-degenerate singularities*, Kodai Math. J. **39** (2016), no. 2, 455–468.
- [BS66] S. Borewicz and I. Shafarevich, *Zahlentheorie*, Lehrbücher und Monographien aus dem Gebiete der Exakten Wissenschaften, Mathematische Reihe, Band 32, Birkhäuser Verlag, Basel-Stuttgart, 1966, Aus dem Russischen übersetzt von Helmut Koch.
- [BV16] B. Bories and W. Veys, *Igusa’s p -adic local zeta function and the monodromy conjecture for non-degenerate surface singularities*, Mem. Amer. Math. Soc. **242** (2016), no. 1145, vii+131.
- [BvdV22] N. Budur and R. van der Veer, *Monodromy conjecture for log generic polynomials*, Math. Z. **301** (2022), no. 1, 713–737.
- [CLS11] D. Cox, J. Little, and H. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [dCMM18] M. de Cataldo, L. Migliorini, and M. Mustață, *Combinatorics and topology of proper toric maps*, J. Reine Angew. Math. **744** (2018), 133–163.
- [Den84] J. Denef, *The rationality of the Poincaré series associated to the p -adic points on a variety*, Invent. Math. **77** (1984), no. 1, 1–23.
- [Den85] ———, *Poles of p -adic complex powers and Newton polyhedra*, Study group on ultrametric analysis, 12th year, 1984/85, No. 1, Secrétariat Math., Paris, 1985, pp. Exp. No. 17, 3.
- [Den87] ———, *On the degree of Igusa’s local zeta function*, Amer. J. Math. **109** (1987), no. 6, 991–1008.
- [Den91a] ———, *Local zeta functions and Euler characteristics*, Duke Math. J. **63** (1991), no. 3, 713–721.
- [Den91b] ———, *Report on Igusa’s local zeta function*, no. 201-203, 1991, Séminaire Bourbaki, Vol. 1990/91, pp. Exp. No. 741, 359–386 (1992).
- [DH01] J. Denef and K. Hoornaert, *Newton polyhedra and Igusa’s local zeta function*, J. Number Theory **89** (2001), no. 1, 31–64.
- [DL92] J. Denef and F. Loeser, *Caractéristiques d’Euler-Poincaré, fonctions zêta locales et modifications analytiques*, J. Amer. Math. Soc. **5** (1992), no. 4, 705–720.
- [DL98] ———, *Motivic Igusa zeta functions*, J. Algebraic Geom. **7** (1998), no. 3, 505–537.
- [DLRS10] J. De Loera, J. Rambau, and F. Santos, *Triangulations*, Algorithms and Computation in Mathematics, vol. 25, Springer-Verlag, Berlin, 2010, Structures for algorithms and applications.
- [dMGP⁺20] A. de Moura, E. Gunther, S. Payne, J. Schuchardt, and A. Stapledon, *Triangulations of simplices with vanishing local h -polynomial*, Algebr. Comb. **3** (2020), no. 6, 1417–1430.
- [ELT22] A. Esterov, A. Lemahieu, and K. Takeuchi, *On the monodromy conjecture for non-degenerate hypersurfaces*, J. Eur. Math. Soc. (JEMS) **24** (2022), no. 11, 3873–3949. MR 4493616
- [Est21] A. Esterov, *The ring of local tropical fans and tropical nearby monodromy eigenvalues*, preprint arXiv:1807.00609 v3, 2021.
- [GJ00] E. Gawrilow and M. Joswig, *polymake: a framework for analyzing convex polytopes*, Polytopes—combinatorics and computation (Oberwolfach, 1997), DMV Sem., vol. 29, Birkhäuser, Basel, 2000, pp. 43–73.
- [Igu75] J.-i. Igusa, *Complex powers and asymptotic expansions. II. Asymptotic expansions*, J. Reine Angew. Math. **278/279** (1975), 307–321.
- [Igu78] ———, *Forms of higher degree*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics, vol. 59, Tata Institute of Fundamental Research, Bombay; by the Narosa Publishing House, New Delhi, 1978.
- [Igu88] ———, *b -functions and p -adic integrals*, Algebraic analysis, Vol. I, Academic Press, Boston, MA, 1988, pp. 231–241.
- [JKYS19] S.-J. Jung, I.-K. Kim, Y. Yoon, and M. Saito, *Spectrum of nondegenerate functions with simplicial Newton polyhedra*, arXiv:1911.09465v3, 2019.
- [Kou76] A. Kouchnirenko, *Polyèdres de Newton et nombres de Milnor*, Invent. Math. **32** (1976), no. 1, 1–31.
- [KS16] E. Katz and A. Stapledon, *Local h -polynomials, invariants of subdivisions, and mixed Ehrhart theory*, Adv. Math. **286** (2016), 181–239.
- [Loe88] F. Loeser, *Fonctions d’Igusa p -adiques et polynômes de Bernstein*, Amer. J. Math. **110** (1988), no. 1, 1–21.

- [Loe90] ———, *Fonctions d'Igusa p -adiques, polynômes de Bernstein, et polyèdres de Newton*, J. Reine Angew. Math. **412** (1990), 75–96.
- [LPS22] M. Larson, S. Payne, and A. Stapledon, *Resolutions of local face modules, functoriality, and nonvanishing of local h -vectors*, Algebr. Comb. (to appear) (2022), arXiv:2209.03543v2.
- [LVP11] A. Lemahieu and L. Van Proeyen, *Monodromy conjecture for nondegenerate surface singularities*, Trans. Amer. Math. Soc. **363** (2011), no. 9, 4801–4829.
- [Mal74] B. Malgrange, *Intégrales asymptotiques et monodromie*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 405–430 (1975).
- [Mil68] J. Milnor, *Singular points of complex hypersurfaces*, Annals of Mathematics Studies, No. 61, Princeton University Press, Princeton, N.J., 1968.
- [Nic10] J. Nicaise, *An introduction to p -adic and motivic zeta functions and the monodromy conjecture*, Algebraic and analytic aspects of zeta functions and L -functions, MSJ Mem., vol. 21, Math. Soc. Japan, Tokyo, 2010, pp. 141–166.
- [Nil12] B. Nill, *Combinatorial questions related to stringy E -polynomials of Gorenstein polytopes*, in Toric Geometry (K. Altmann et. al., eds.), 2012, pp. 62–64.
- [Poo02] B. Poonen, *The Grothendieck ring of varieties is not a domain*, Math. Res. Lett. **9** (2002), no. 4, 493–497.
- [Que] M. H. Quek, *Around the motivic monodromy conjecture for non-degenerate hypersurfaces*, Manuscripta Math. (to appear), arXiv:2208.02566v4.
- [RV03] B. Rodrigues and W. Veys, *Poles of zeta functions on normal surfaces*, Proc. London Math. Soc. (3) **87** (2003), no. 1, 164–196.
- [Sai19] T. Saito, *Milnor monodromies and mixed Hodge structures for non-isolated hypersurface singularities*, Adv. Math. **342** (2019), 134–164.
- [Sta92] R. Stanley, *Subdivisions and local h -vectors*, J. Amer. Math. Soc. **5** (1992), no. 4, 805–851.
- [Sta17] A. Stapledon, *Formulas for monodromy*, Res. Math. Sci. **4** (2017), Paper No. 8, 42 pp.
- [TT16] K. Takeuchi and M. Tibăr, *Monodromies at infinity of non-tame polynomials*, Bull. Soc. Math. France **144** (2016), no. 3, 477–506.
- [Var76] A. Varchenko, *Zeta-function of monodromy and Newton's diagram*, Invent. Math. **37** (1976), no. 3, 253–262.
- [Vey93] W. Veys, *Poles of Igusa's local zeta function and monodromy*, Bull. Soc. Math. France **121** (1993), no. 4, 545–598.
- [Vey07] ———, *Monodromy eigenvalues and zeta functions with differential forms*, Adv. Math. **213** (2007), no. 1, 341–357.
- [VS12] J. Viu-Sos, *Computing the Igusa and Topological zeta functions of a Newton non-degenerated polynomial*, <https://github.com/jviusos/ZetaFunctionsNewtonND-Sagemath>, 2012.
- [VS22] ———, *An introduction to p -adic and motivic integration, zeta functions and invariants of singularities, p -adic analysis, arithmetic and singularities*, Contemp. Math., vol. 778, Amer. Math. Soc., [Providence], RI, [2022] ©2022, pp. 103–176.

STANFORD U. DEPARTMENT OF MATHEMATICS, 450 JANE STANFORD WAY, STANFORD, CA 94305

Email address: mwl Larson@stanford.edu

UT DEPARTMENT OF MATHEMATICS, 2515 SPEEDWAY, RLM 8.100, AUSTIN, TX 78712

Email address: sampayne@utexas.edu

SYDNEY MATHEMATICS RESEARCH INSTITUTE, L4.42, QUADRANGLE A14, UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA

Email address: astapl dn@gmail.com