

THE BERGMAN FAN OF A POLYMATROID

COLIN CROWLEY, JUNE HUH, MATT LARSON, CONNOR SIMPSON,
AND BOTONG WANG

ABSTRACT. We introduce the Bergman fan of a polymatroid and prove that the Chow ring of the Bergman fan is isomorphic to the Chow ring of the polymatroid. Using the Bergman fan, we establish the Kähler package for the Chow ring of the polymatroid, recovering and strengthening a result of Pagaria–Pezzoli.

1. INTRODUCTION

By definition, a *matroid* on a finite set E is given by a rank function $\text{rk}: 2^E \rightarrow \mathbb{Z}_{\geq 0}$ satisfying the following for all subsets $A_1, A_2 \subseteq E$:

(Submodularity) $\text{rk}(A_1 \cup A_2) + \text{rk}(A_1 \cap A_2) \leq \text{rk}(A_1) + \text{rk}(A_2)$.

(Monotonicity) If $A_1 \subseteq A_2$, then $\text{rk}(A_1) \leq \text{rk}(A_2)$.

(Boundedness) The rank of a subset is at most its cardinality.

(Normalization) The rank of the empty subset is zero.

Sans “boundedness”, the axioms above define a *polymatroid*. Throughout this paper, we assume that the polymatroid is *loopless*:

(Looplessness) The rank of any nonempty subset is nonzero.

If P is a polymatroid on E , then its *rank* is $\text{rk}(P) := \text{rk}(E)$. A *flat* of P is a subset $F \subseteq E$ that is maximal among sets of its rank. Ordered by inclusion, the flats of P form a lattice \mathcal{L}_P .¹ The intersection of two flats is a flat, so any subset A of E is contained in a unique minimal flat $\text{cl}_P(A)$, called the *closure* of A in P , which is obtained by intersecting all flats that contain A .

Matroids are combinatorial abstractions of hyperplane arrangements, and more generally, polymatroids are combinatorial abstractions of subspace arrangements.

Example 1.1. Let V_1, \dots, V_n be linear subspaces of a vector space V over a field \mathbb{F} . There is a polymatroid P on the set $E = \{1, \dots, n\}$ defined by the rank function

$$\text{rk}(A) := \text{codim}_V(\cap_{i \in A} V_i).$$

The polymatroid P is a matroid if and only if every V_i is a hyperplane. The map $F \mapsto \cap_{i \in F} V_i$ is a bijection between the flats of P and the subspaces of V obtained by intersecting some of the V_i 's. A polymatroid arising in this way is said to be *realizable* over \mathbb{F} , and the subspace arrangement is called a *realization* of P over \mathbb{F} .

Much of a hyperplane arrangement's combinatorial data is captured by intersection theory on its wonderful compactification [5]. To extend this from hyperplane

¹Unlike in the case of matroids, the lattice of flats of a polymatroid can fail to be graded or atomic.

arrangements to non-realizable matroids, one must replace the wonderful compactification with a combinatorial object, the Bergman fan of a matroid. The purpose of the present paper is to introduce the *Bergman fan of a polymatroid*, a combinatorial model for the wonderful compactification of a subspace arrangement. As in the case of matroids, the Bergman fan of a polymatroid is a tropical variety of degree one. In Section 4, we show that the Chow ring of a polymatroid satisfies the Kähler package with respect to any strictly convex piecewise linear function on its Bergman fan, recovering and strengthening a result of Pagaria and Pezzoli [15, Theorems 4.7 and 4.21].

Our construction of the Bergman fan is inspired by a geometric observation: over an infinite field, the wonderful compactification of any subspace arrangement can be realized as the wonderful compactification of a hyperplane arrangement, taken with respect to an appropriate building set (Remark 3.5). The construction immediately reveals that the Bergman fan of a polymatroid and the Bergman fan of the associated matroid have the same support. Thus, the Kähler package for the polymatroid follows from that of the associated matroid [1] and the general fact that the validity of the Kähler package for the Chow ring of a fan depends only on the support of the fan [2].

1.1. The Bergman fan of a Boolean polymatroid. An important special case is that of Boolean polymatroids. Let $\pi: \tilde{E} \rightarrow E$ be a surjective map between finite sets. The *Boolean polymatroid* $B(\pi)$ on E is defined by the rank function

$$\mathrm{rk}_{B(\pi)}(A) = |\pi^{-1}(A)| \quad \text{for } A \subseteq E.$$

We write $N_{\tilde{E}}$ for $\mathbb{Z}^{\tilde{E}}/\mathbb{Z}(1, 1, \dots, 1)$, and for $S \subseteq \tilde{E}$, write \mathbf{e}_S for $\sum_{i \in S} \mathbf{e}_i$ in $N_{\tilde{E}} \otimes \mathbb{R}$.

Definition 1.2. The *Bergman fan* $\Sigma_{B(\pi)}$ of the Boolean polymatroid $B(\pi)$ is the fan in $N_{\tilde{E}} \otimes \mathbb{R}$ with cones

$$\sigma_{\mathcal{F}, S} := \mathrm{cone}(\mathbf{e}_{\pi^{-1}(F_1)}, \dots, \mathbf{e}_{\pi^{-1}(F_k)}) + \mathrm{cone}(\mathbf{e}_i)_{i \in S},$$

for every chain $\mathcal{F} = \{\emptyset \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_k \subsetneq E\}$ and subset S of \tilde{E} not containing any fiber of π .

Throughout the paper, we write n for the cardinality of E .

Definition 1.3. An *ordered transversal* of π is a sequence s_1, \dots, s_n of elements of \tilde{E} such that each fiber of π contains exactly one element of the sequence. The *polypermutohedron* $Q(\pi)$ is the convex hull of the vectors $\sum_{i=1}^n i \mathbf{e}_{s_i}$ in $\mathbb{R}^{\tilde{E}}$, where s_1, \dots, s_n range over all ordered transversals of π .

In Appendix A, we show $\Sigma_{B(\pi)}$ is the inner normal fan of the polypermutohedron $Q(\pi)$. Definition 1.2 implies that $\Sigma_{B(\pi)}$ is a complete unimodular fan in $N_{\tilde{E}}$.

Example 1.4. If π is a bijection, an ordered transversal of π is a permutation of \tilde{E} , and $Q(\pi)$ is the standard permutohedron in $\mathbb{R}^{\tilde{E}}$. This recovers a familiar fact: the Bergman fan of the Boolean matroid is the normal fan of the permutohedron.

Example 1.5. When E is a singleton, an ordered transversal of π is an element of \tilde{E} , and $Q(\pi)$ is the standard simplex in $\mathbb{R}^{\tilde{E}}$. Thus, the Bergman fan of a Boolean polymatroid on a singleton is the normal fan of the standard simplex.

Example 1.6. When $|E| = n - 1$ and all fibers of π have size d , the toric variety corresponding to $\Sigma_{B(\pi)}$ is a generalization of the Losev–Manin space of curves which compactifies the moduli space of configurations of n points in \mathbb{A}^d up to translation and scaling [9, Corollary 5.6].

1.2. The Bergman fan of a polymatroid. Let P be a polymatroid on E , and let $\pi: \tilde{E} \rightarrow E$ be a surjective map satisfying $\text{rk}_P(i) = |\pi^{-1}(i)|$ for every i in E .

Definition 1.7. The *Bergman fan* Σ_P is the subfan of $\Sigma_{B(\pi)}$ with cones given by

$$\sigma_{\mathcal{F}, S} := \text{cone}(\mathbf{e}_{\pi^{-1}(F_1)}, \dots, \mathbf{e}_{\pi^{-1}(F_k)}) + \text{cone}(\mathbf{e}_i)_{i \in S},$$

one for every chain of flats $\mathcal{F} = \{\emptyset = F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_k \subsetneq E\}$ of P and a subset S of \tilde{E} such that $\text{rk}_P(F_i \cup \pi(T)) > \text{rk}_P(F_i) + |T|$ for all $0 \leq i \leq k$ and all nonempty $T \subseteq S \setminus \pi^{-1}(F_i)$.

The Bergman fan Σ_P is unimodular with respect to $N_{\tilde{E}}$, defining a smooth toric variety X_P over \mathbb{C} . We write $A(\Sigma_P)$ for the Chow ring of X_P . We relate $A(\Sigma_P)$ to the *Chow ring* of the polymatroid P (Definition 4.1), denoted $\text{DP}(P)$, introduced in [15, Section 4]. Our main result states the following.

Theorem 4.2. *There is a natural isomorphism of graded rings $\text{DP}(P) \cong A(\Sigma_P)$.*

In Corollary 4.3, we use Theorem 4.2 to recover a Gröbner basis for $\text{DP}(P)$ found in [15]. In Corollary 4.7, we prove the Kähler package for $A(\Sigma_P)$ with respect to the cone of strictly convex piecewise linear function on Σ_P , extending the Kähler package for $\text{DP}(P)$ with respect to the σ -cone in [15]. See Remark 4.8 for a comparison of the two cones.

1.3. Building sets. In fact, our results hold for polymatroids P equipped with a *geometric building set* \mathcal{G} (Section 3). The statements above are specializations of our results to the case when \mathcal{G} consists of all nonempty flats of P . In maximal generality, we define the *Bergman fan* of (P, \mathcal{G}) , denoted $\Sigma_{P, \mathcal{G}}$ (Definition 3.6). The Chow ring $\text{DP}(P, \mathcal{G})$ associated to (P, \mathcal{G}) was introduced in [15], and it is isomorphic to $A(\Sigma_{P, \mathcal{G}})$ (Theorem 4.2). All corollaries continue to hold, including the Kähler package for $\text{DP}(P, \mathcal{G})$.

Example 1.8. Let P be the polymatroid on $E = \{a, b, c\}$ with flats (depicted in Fig. 1) satisfying $\text{rk}_P(a) = \text{rk}_P(b) = 1$, $\text{rk}_P(ab) = \text{rk}_P(c) = 2$, and $\text{rk}_P(abc) = 3$. Define $\pi: \tilde{E} = \{a, b, c_1, c_2\} \rightarrow E$ by $\pi(a) = a$, $\pi(b) = b$, and $\pi(c_i) = c$. The Bergman fan Σ_P is a pure 2-dimensional simplicial fan in $\mathbb{R}^{\tilde{E}}/\mathbb{R}(1, 1, 1, 1)$ with rays generated by the vectors

$$\mathbf{e}_a, \mathbf{e}_b, \mathbf{e}_{c_1}, \mathbf{e}_{c_2}, \mathbf{e}_{ab}, \mathbf{e}_{c_1 c_2}$$

and 2-dimensional cones

$$\begin{aligned} \sigma_{\{c\}, c_1} &= \text{cone}(\mathbf{e}_{c_1 c_2}, \mathbf{e}_{c_1}), & \sigma_{\{c\}, c_2} &= \text{cone}(\mathbf{e}_{c_1 c_2}, \mathbf{e}_{c_2}) \\ \sigma_{\{a\}, c_1} &= \text{cone}(\mathbf{e}_a, \mathbf{e}_{c_1}), & \sigma_{\{a\}, c_2} &= \text{cone}(\mathbf{e}_a, \mathbf{e}_{c_2}) \\ \sigma_{\{b\}, c_1} &= \text{cone}(\mathbf{e}_b, \mathbf{e}_{c_1}), & \sigma_{\{b\}, c_2} &= \text{cone}(\mathbf{e}_b, \mathbf{e}_{c_2}) \\ \sigma_{\{a \subset ab\}, \emptyset} &= \text{cone}(\mathbf{e}_a, \mathbf{e}_{ab}), & \sigma_{\{b \subset ab\}, \emptyset} &= \text{cone}(\mathbf{e}_b, \mathbf{e}_{ab}) \end{aligned}$$

In Section 3, we will come to understand Σ_P as the *nested set fan* (Section 3) of a matroid \tilde{P} , called the *multisymmetric lift* of P (Definition-Proposition 2.11), with respect to a building set $\tilde{\mathcal{G}}$ constructed from the maximal building set of P .




FIGURE 1. The lattice of flats of P (left), and of its multisymmetric lift \tilde{P} (right). The symmetric group of order 2 acts on \tilde{P} by swapping c_1 and c_2 , and the “geometric” flats invariant under this action (drawn in blue) are precisely the sets of the form $\pi^{-1}(F)$ with F a flat of P . Elements of the building set $\tilde{\mathcal{G}}$ for \tilde{P} are underlined.

Organization. In Section 2, we develop the combinatorics of multisymmetric matroids and lifts, a key tool throughout this paper. We use lifts to define the Bergman fan of a polymatroid (with respect to a geometric building set) in Section 3. Finally, in Section 4, we show that the Chow ring of the Bergman fan agrees with the polymatroid Chow ring of [15] and derive consequences. Examples 1.1 and 2.12, and Remarks 3.5, 3.12, and 4.6 explain the geometry underlying this work.

Acknowledgements. We thank Spencer Backman for comments and conversations. The second author is partially supported by a Simons Investigator Grant and NSF Grant DMS-2053308, the third is supported by an NDSEG fellowship, and the last is supported by a Sloan fellowship.

2. MULTISYMMETRIC MATROIDS

Many proofs in Section 4 reduce statements about polymatroids to known statements about matroids. The key tool for this reduction is multisymmetric matroids, a new cryptomorphic formulation of polymatroids. Let $\mathfrak{S}_{\tilde{E}}$ denote the symmetric group on a finite set \tilde{E} . We continue to assume that all (poly)matroids are loopless.

Definition 2.1. A *multisymmetric matroid* is a matroid M on \tilde{E} equipped with a partition $\tilde{E} = \tilde{E}_1 \sqcup \cdots \sqcup \tilde{E}_n$ such that the action of $\Gamma = \mathfrak{S}_{\tilde{E}_1} \times \cdots \times \mathfrak{S}_{\tilde{E}_n}$ on \tilde{E} takes flats to flats. The *geometric part* of a subset $S \subseteq \tilde{E}$ is $S^{\text{geo}} := \bigcap_{\gamma \in \Gamma} (\gamma \cdot S)$. We call a subset $S \subseteq \tilde{E}$ *geometric* if $S = S^{\text{geo}}$.

For a multisymmetric matroid M , we write \mathcal{L}_M^Γ for its poset of geometric flats.

Example 2.2. Any matroid M on \tilde{E} can be given the *trivial* multisymmetric structure by setting $\Gamma = \prod_{e \in \tilde{E}} \mathfrak{S}_{\{e\}}$. In this case, $\mathcal{L}_M^\Gamma = \mathcal{L}_M$.

Example 2.3. If M is multisymmetric on $\tilde{E} = \tilde{E}_1$, then M is a uniform matroid, and \mathcal{L}_M^Γ is $\{\emptyset, \tilde{E}\}$.

Example 2.4. Let \tilde{E} be the set of edges of the complete graph K_4 , and let M be the graphic matroid of K_4 . No transposition of $\mathfrak{S}_{\tilde{E}}$ preserves the flats of M , so M has no non-trivial multisymmetric structures.

Closure in a multisymmetric matroid is restricted by the group action.

Lemma 2.5. *If M is multisymmetric and $S \subseteq \tilde{E}$ is geometric, $\text{cl}_M(S)$ is geometric.*

Proof. If $S \subseteq \tilde{E}$ is geometric and $\gamma \in \Gamma$, $\gamma \cdot \text{cl}_M(S) = \text{cl}_M(\gamma \cdot S) = \text{cl}_M(S)$. \square

Corollary 2.6. *If M is multisymmetric, then the geometric flats of M form a sublattice of \mathcal{L}_M .*

Proof. If F and G are two geometric flats, then $\text{cl}_M(F \cup G)$ is geometric by Lemma 2.5. The intersection of two geometric flats is also geometric. In other words, the set of geometric flats is closed under both join and meet, and therefore forms a sublattice of \mathcal{L}_M . \square

Corollary 2.7. *If M is multisymmetric on $\tilde{E} = \tilde{E}_1 \sqcup \dots \sqcup \tilde{E}_n$, then \mathcal{L}_M^Γ is the lattice of flats of the polymatroid P on the set of indices $E = \{1, \dots, n\}$ defined by the rank function $\text{rk}_P(A) := \text{rk}_M(\cup_{i \in A} \tilde{E}_i)$.*

Proof. Define $\pi: \tilde{E} \rightarrow E$ by setting $\pi^{-1}(i) = \tilde{E}_i$. If F is a flat of P , then for all $F \subsetneq A \subseteq \{1, \dots, n\}$,

$$\text{rk}_M(\pi^{-1}(F)) = \text{rk}_P(F) < \text{rk}_P(A) = \text{rk}_M(\pi^{-1}(A)).$$

By Lemma 2.5, $\text{cl}_M(\pi^{-1}(F))$ is geometric, so we conclude that $\text{cl}_M(\pi^{-1}(F)) = \pi^{-1}(F)$. In other words, $\pi^{-1}(F)$ is a flat of M . Conversely, if F is not a flat of P , then $\text{rk}_P(F) = \text{rk}_P(F \cup i)$ for some i not in F . This implies $\text{cl}_M(\pi^{-1}(F)) \supsetneq \pi^{-1}(F \cup i)$, so $\pi^{-1}(F)$ is not a flat of M . Therefore, F is a flat of P if and only if $\pi^{-1}(F)$ is a geometric flat of M , and we have an isomorphism of lattices

$$\mathcal{L}_P \longrightarrow \mathcal{L}_M^\Gamma, \quad F \longmapsto \pi^{-1}(F). \quad \square$$

Lemma 2.8. *Let M be a multisymmetric matroid on $\tilde{E} = \tilde{E}_1 \sqcup \dots \sqcup \tilde{E}_n$.*

(i) *If $S \subseteq \tilde{E}$, then either $\text{cl}_M(S) \cap \tilde{E}_i = \tilde{E}_i$ or $\text{cl}_M(S) \cap \tilde{E}_i = S \cap \tilde{E}_i$.*

(ii) *If F is a flat of M , then $\text{rk}_M(F) = \text{rk}_M(F^{\text{geo}}) + |F \setminus F^{\text{geo}}|$.*

Proof. A permutation of $\tilde{E}_i \setminus S$ induces an automorphism of M that fixes S . Any such automorphism also fixes $\text{cl}_M(S)$ because automorphisms commute with closure. Hence, if $(\tilde{E}_i \setminus S) \cap \text{cl}_M(S)$ is nonempty, then $\tilde{E}_i \subseteq \text{cl}_M(S)$. This proves (i).

To prove (ii), let $\{s_1, \dots, s_k\} = F \setminus F^{\text{geo}}$. If (ii) fails then for some $1 \leq i \leq k$,

$$\text{rk}_M(F^{\text{geo}} \cup \{s_1, \dots, s_i\}) = \text{rk}_M(F^{\text{geo}} \cup \{s_1, \dots, s_{i+1}\}).$$

Therefore if $s_{i+1} \in \tilde{E}_j$ then $\tilde{E}_j \subseteq \text{cl}_M(F^{\text{geo}} \cup \{s_1, \dots, s_i\}) \subseteq F$ by the first part. Consequently, $s_{i+1} \in F^{\text{geo}}$, a contradiction. \square

Lemma 2.9. *A multisymmetric matroid is determined by its geometric sets and their ranks.*

Proof. Let M and M' be two multisymmetric matroids on $\tilde{E} = \tilde{E}_1 \sqcup \dots \sqcup \tilde{E}_n$, and suppose that for all $A \subseteq \{1, \dots, n\}$, $\text{rk}_M(\cup_{i \in A} \tilde{E}_i) = \text{rk}_{M'}(\cup_{i \in A} \tilde{E}_i)$. If F is a flat of M , and F' is the closure of F in M' , then

$$\text{rk}_{M'}(F) \leq \text{rk}_{M'}(F^{\text{geo}}) + \text{rk}_{M'}(F \setminus F^{\text{geo}}) \leq \text{rk}_M(F^{\text{geo}}) + |F \setminus F^{\text{geo}}| = \text{rk}_M(F)$$

by Lemma 2.8(ii). Symmetrically, $\text{rk}_M(F') \leq \text{rk}_{M'}(F')$, so

$$\text{rk}_{M'}(F) \leq \text{rk}_M(F) \leq \text{rk}_M(F') \leq \text{rk}_{M'}(F').$$

The left- and rightmost terms are equal, so $F = F'$ because F is a flat of M . This shows that M and M' have the same flats, and that their flats have the same ranks, so M and M' are equal. \square

Remark 2.10. The closure of a geometric set is a geometric flat by Lemma 2.8(i), so Lemma 2.9 implies that a multisymmetric matroid is also determined by its geometric flats and their ranks.

2.1. Lifts. Let P be a polymatroid on $E = \{1, \dots, n\}$ and M a multisymmetric matroid on $\tilde{E} = \tilde{E}_1 \sqcup \dots \sqcup \tilde{E}_n$. If P is the polymatroid given by Corollary 2.7, then we say that M is a *multisymmetric lift* of P . If $\text{rk}_M(\tilde{E}_i) = |\tilde{E}_i|$ for all $1 \leq i \leq n$, then we say that M is a *minimal* multisymmetric lift of P .

Definition-Proposition 2.11. A polymatroid P has a unique minimal multisymmetric lift \tilde{P} constructed as follows. For $1 \leq i \leq n$, let $\tilde{E}_i = \{1, \dots, \text{rk}_P(i)\}$, and $\tilde{E} = \tilde{E}_1 \sqcup \dots \sqcup \tilde{E}_n$. Define the projection $\pi: \tilde{E} \rightarrow E$ by $\pi^{-1}(i) = \tilde{E}_i$.

The minimal multisymmetric lift of P is the matroid \tilde{P} on \tilde{E} with rank function

$$\text{rk}_{\tilde{P}}(S) = \min\{\text{rk}_P(A) + |S \setminus \pi^{-1}(A)| : A \subseteq E\}.$$

Example 2.12. Suppose P is the polymatroid realized by an arrangement of subspaces V_1, \dots, V_n in V as in Example 1.1. The minimal multisymmetric lift \tilde{P} is realized by any hyperplane arrangement

$$\{V_{i,j} : 1 \leq i \leq n, 1 \leq j \leq \text{rk}_P(i)\},$$

where $V_{i,1}, \dots, V_{i,\text{rk}_P(i)}$ are generic hyperplanes containing V_i . Here, we assume that the vector space is defined over an infinite field. The flats of \tilde{P} are in bijection with the subspaces that can be obtained by intersecting some of the hyperplanes $V_{i,j}$. Under this bijection, the geometric flats correspond to intersections of any collection of V_i 's.

Versions of the construction in Definition-Proposition 2.11 make many independent appearances in the literature, for example, in [10, §2] and [11, Propositions 3.1 and 3.2]. The most complete treatment we are aware of is [13, §2], whose terminology differs from ours.

Notation. We continue to use the notations of Definition-Proposition 2.11 in the remainder of this section. For visual clarity, we often write $M = \tilde{P}$. As usual, Γ stands for the product of symmetric groups that acts on M .

Proposition 2.13. *The minimal lift $M = \tilde{P}$ is multisymmetric. Explicitly,*

- (i) rk_M is a matroid rank function, and
- (ii) the action of Γ preserves flats.

Proof. The proof of the first part is reproduced from [14, Proof of Theorem 11.1.9]. We need to check that rk_M is non-negative, increasing, submodular, and satisfies $\text{rk}_M(S) \leq |S|$ for $S \subseteq \tilde{E}$.

Clearly values of rk_M are non-negative. Let $S \subseteq \tilde{E}$ and $s \in \tilde{E} \setminus S$. For all $A \subseteq E$,

$$\text{rk}_P(A) + |S \setminus \pi^{-1}(A)| \leq \text{rk}_P(A) + |S \cup s \setminus \pi^{-1}(A)|,$$

so rk_M is increasing. Moreover, by induction on $|S|$,

$$\begin{aligned} \text{rk}_M(S \cup s) &= \min\{\text{rk}_P(A) + |S \cup s \setminus \pi^{-1}(A)| : A \subseteq E\} \\ &\leq \min\{\text{rk}_P(A) + |S \setminus \pi^{-1}(A)| : A \subseteq E\} + 1 \\ &= \text{rk}_M(S) + 1 \leq |S| + 1 = |S \cup s|, \end{aligned}$$

so it only remains to check that rk_M is submodular. Let $S_1, S_2 \subseteq \tilde{E}$ and $A_1, A_2 \subseteq E$ such that $\text{rk}_M(S_i) = \text{rk}_P(A_i) + |S_i \setminus \pi^{-1}(A_i)|$. Then

$$\begin{aligned} \text{rk}_M(S_1) + \text{rk}_M(S_2) &= \text{rk}_P(A_1) + |S_1 \setminus \pi^{-1}(A_1)| + \text{rk}_P(A_2) + |S_2 \setminus \pi^{-1}(A_2)| \\ &\geq \text{rk}_P(A_1 \cup A_2) + \text{rk}_P(A_1 \cap A_2) \\ &\quad + |S_1 \setminus \pi^{-1}(A_1)| + |S_2 \setminus \pi^{-1}(A_2)| \quad \text{by submodularity of } \text{rk}_P \\ &\geq \text{rk}_P(A_1 \cup A_2) + \text{rk}_P(A_1 \cap A_2) \\ &\quad + |(S_1 \cup S_2) \setminus \pi^{-1}(A_1 \cup A_2)| + |(S_1 \cap S_2) \setminus \pi^{-1}(A_1 \cap A_2)| \\ &\geq \text{rk}_M(S_1 \cup S_2) + \text{rk}_M(S_1 \cap S_2). \end{aligned}$$

For the second part, let $\gamma \in \Gamma$ and $S \subseteq \tilde{E}$. For any subset $A \subseteq E$, we have that $|S \setminus \pi^{-1}(A)| = |(\gamma \cdot S) \setminus \pi^{-1}(A)|$, and hence $\text{rk}_M(S) = \text{rk}_M(\gamma \cdot S)$. \square

The following lemma implies \tilde{P} is a multisymmetric lift of P .

Lemma 2.14. *Fix notation as in Definition-Proposition 2.11, and set $M := \tilde{P}$. If $S \subseteq \tilde{E}$ is stable under Γ , then $\text{rk}_M(S) = \text{rk}_P(\pi(S))$.*

Proof. Since S is stable under the action of Γ , S is a union of fibers of π . Hence,

$$\begin{aligned} \text{rk}_M(S) &= \min\{\text{rk}_P(A) + |S \setminus \pi^{-1}(A)| : A \subseteq E\} \quad \text{by the definition of } \text{rk}_M \\ &= \min\{\text{rk}_P(A) + |S \setminus \pi^{-1}(A)| : A \subseteq \pi(S)\} \quad \text{because } \text{rk}_P \text{ is increasing} \\ &= \min\left\{\text{rk}_P(A) + \sum_{i \in \pi(S) \setminus A} |\pi^{-1}(i)| : A \subseteq \pi(S)\right\} \quad \text{because } S = \pi^{-1}(\pi(S)) \\ &= \min\left\{\text{rk}_P(A) + \sum_{i \in \pi(S) \setminus A} \text{rk}_P(i) : A \subseteq \pi(S)\right\} \quad \text{by Definition-Proposition 2.11} \\ &\geq \text{rk}_P(\pi(S)) \quad \text{by submodularity of } \text{rk}_P. \end{aligned}$$

On the other hand, taking $A = \pi(S)$ in Definition-Proposition 2.11, we have $\text{rk}_M(S) \leq \text{rk}_P(\pi(S))$. \square

Proof of Definition-Proposition 2.11. Let P be a polymatroid, and let $M = \tilde{P}$ on $\tilde{E} = \tilde{E}_1 \sqcup \cdots \sqcup \tilde{E}_n$, acted upon by Γ , be as in the statement of Definition-Proposition 2.11. By Proposition 2.13 and Lemma 2.14, M is a multisymmetric lift of P . It is minimal because

$$\text{rk}_M(\tilde{E}_i) = \text{rk}_P(\pi(\tilde{E}_i)) = \text{rk}_P(i) = |\tilde{E}_i|.$$

The uniqueness statement follows from Lemma 2.9 and the fact that the ranks of geometric sets in any lift of P are determined by P . \square

2.2. Operations. The formation of minimal multisymmetric lifts commutes with some polymatroid operations. Let F be a flat of a polymatroid P on E . The *restriction* of P to F , denoted $P|_F$, is the polymatroid on F with rank function

$$\mathrm{rk}_{P|_F}(A) := \mathrm{rk}_P(A), \quad A \subseteq F.$$

There is a lattice isomorphism

$$\mathcal{L}_{P|_F} \rightarrow \{G \in \mathcal{L}_P : G \leq F\}, \quad H \mapsto H.$$

If P_1 and P_2 are polymatroids on E^1 and E^2 , respectively, then their *direct sum* is the polymatroid $P_1 \oplus P_2$ on $E^1 \sqcup E^2$ with rank function

$$\mathrm{rk}_{P_1 \oplus P_2}(S) = \mathrm{rk}_{P_1}(S \cap E^1) + \mathrm{rk}_{P_2}(S \cap E^2).$$

There is a lattice isomorphism

$$\mathcal{L}_{P_1} \times \mathcal{L}_{P_2} \rightarrow \mathcal{L}_{P_1 \oplus P_2}, \quad (F, G) \mapsto F \sqcup G.$$

Lemma 2.15. *If P_1 and P_2 are polymatroids, then $\widetilde{P_1 \oplus P_2} = \widetilde{P_1} \oplus \widetilde{P_2}$.*

Proof. From Corollary 2.7 and the definition of $P_1 \oplus P_2$, it follows that $\widetilde{P_1 \oplus P_2}$ lifts $P_1 \oplus P_2$. If Γ_1 and Γ_2 are the groups acting on $\widetilde{P_1}$ and $\widetilde{P_2}$, then $\Gamma_1 \times \Gamma_2$ acts on $\widetilde{P_1 \oplus P_2}$, so it is multisymmetric. It is minimal by minimality of $\widetilde{P_1}$ and $\widetilde{P_2}$, so the lemma holds by the uniqueness statement of Definition-Proposition 2.11. \square

Lemma 2.16. *If F is a flat of a polymatroid P , then $\widetilde{P|_F} = \widetilde{P}|_{\pi^{-1}(F)}$.*

Proof. The rank functions of both sides are obtained by restriction, so $\widetilde{P}|_{\pi^{-1}(F)}$ lifts $P|_F$. If Γ acts on \widetilde{P} , then a subset of Γ 's factors acts on $\widetilde{P}|_{\pi^{-1}(F)}$, so it is multisymmetric. It is minimal because \widetilde{P} is, so Definition-Proposition 2.11 implies the lemma. \square

3. BUILDING SETS AND BERGMAN FANS

Here, we recall the combinatorics of geometric building sets. We then generalize the definition of the Bergman fan of a polymatroid given in the introduction by associating a fan to each polymatroid equipped with a geometric building set.

3.1. Geometric building sets. If P is a polymatroid, $\mathcal{G} \subseteq \mathcal{L}_P$, and $F \in \mathcal{L}_P$, let

$$\mathcal{G}_{\leq F} := \{G \in \mathcal{G} : G \leq F\},$$

and write $\max \mathcal{G}$ for the set of maximal elements of \mathcal{G} .

Definition 3.1. A *geometric building set* of a polymatroid P is a collection \mathcal{G} of nonempty flats such that for all $F \in \mathcal{L}_P \setminus \{\emptyset\}$, the map

$$\prod_{G \in \max \mathcal{G}_{\leq F}} \mathcal{L}_{P|_G} \rightarrow \mathcal{L}_{P|_F}$$

is an isomorphism, and

$$\sum_{G \in \max \mathcal{G}_{\leq F}} \mathrm{rk}(G) = \mathrm{rk}(F).$$

If $E \in \mathcal{G}$, then a *nested set* of P with respect to \mathcal{G} is a subset $\mathcal{N} \subseteq \mathcal{G}$ such that for all $\{F_1, \dots, F_k\} \subseteq \mathcal{N}$ pairwise incomparable with $k \geq 2$, we have that

$$\mathrm{cl}_P(F_1 \cup \dots \cup F_k) \notin \mathcal{G}.$$

With respect to a fixed geometric building set, a subset of a nested set is nested, so nested sets form a simplicial complex. All building sets are henceforth assumed to contain E^2 .

Example 3.2. The maximal geometric building set of P is the collection of all nonempty flats. With respect to this building set, the nested sets are flags of nonempty flats.

Lemma 3.3. *Let P be a polymatroid, \tilde{P} its minimal multisymmetric lift, and π as in Definition-Proposition 2.11. If \mathcal{G} is a geometric building set for P , then*

$$\tilde{\mathcal{G}} = \{\pi^{-1}(G) : G \in \mathcal{G}\} \cup \{\text{atoms of } \mathcal{L}_{\tilde{P}}\}$$

is a geometric building set for \tilde{P} .

Proof. Let F be a flat of $M = \tilde{P}$. By Lemma 2.15 and Lemma 2.16, the map

$$\prod_{G \in \max \tilde{\mathcal{G}}_{\leq F^{\text{geo}}}} \mathcal{L}_{M|_G} \rightarrow \mathcal{L}_{M|_F}$$

factors into a chain of isomorphisms

$$\prod_{G \in \max \tilde{\mathcal{G}}_{\leq F^{\text{geo}}}} \mathcal{L}_{M|_G} \cong \prod_{H \in \max \mathcal{G}_{\leq \pi(F^{\text{geo}})}} \mathcal{L}_{\tilde{P}|_H} \cong \mathcal{L}_{P|\pi(F^{\text{geo}})} \cong \mathcal{L}_{M|_{F^{\text{geo}}}},$$

and by Lemma 2.14,

$$\begin{aligned} \sum_{G \in \max \tilde{\mathcal{G}}_{\leq F^{\text{geo}}}} \text{rk}_M(G) &= \sum_{G \in \max \tilde{\mathcal{G}}_{\leq F^{\text{geo}}}} \text{rk}_P(\pi(G)) \\ &= \sum_{H \in \max \mathcal{G}_{\leq \pi(F^{\text{geo}})}} \text{rk}_P(H) = \text{rk}_P(\pi(F^{\text{geo}})) = \text{rk}_M(F^{\text{geo}}). \end{aligned}$$

Consequently, by Lemma 2.8(ii), the first displayed condition of Definition 3.1 holds:

$$\begin{aligned} \prod_{G \in \max \tilde{\mathcal{G}}_{\leq F}} \mathcal{L}_{M|_G} &= \prod_{G \in \max \tilde{\mathcal{G}}_{\leq F^{\text{geo}}}} \mathcal{L}_{M|_G} \times \prod_{i \in F \setminus F^{\text{geo}}} \mathcal{L}_{M|_{\{i\}}} \\ &\cong \mathcal{L}_{M|_{F^{\text{geo}}}} \times \prod_{i \in F \setminus F^{\text{geo}}} \mathcal{L}_{M|_{\{i\}}} \cong \mathcal{L}_{M|_F}, \end{aligned}$$

Also by Lemma 2.8(ii), the second displayed condition of Definition 3.1 holds:

$$\begin{aligned} \sum_{G \in \max \tilde{\mathcal{G}}_{\leq F}} \text{rk}_M(G) &= \text{rk}_M(F^{\text{geo}}) + \sum_{i \in F \setminus F^{\text{geo}}} \text{rk}_M(i) \\ &= \text{rk}_M(F^{\text{geo}}) + |F \setminus F^{\text{geo}}| = \text{rk}_M(F). \end{aligned}$$

□

Lemma 3.4. *If \mathcal{G} is a geometric building set for P , then $\mathcal{N} \subseteq \mathcal{G}$ is \mathcal{G} -nested if and only if $\tilde{\mathcal{N}} = \{\pi^{-1}(F) : F \in \mathcal{N}\}$ is $\tilde{\mathcal{G}}$ -nested.*

²In the realizable case, this assumption guarantees the associated wonderful compactification is smooth and can be described as an iterated blow-up [5, §4.1]. Combinatorially, we lose nothing by this assumption [15, Remark 4.1].

Proof. Set $M = \tilde{P}$. Let $\{F_1, \dots, F_k\} \subseteq \mathcal{N}$ be a set of pairwise incomparable flats. By Corollary 2.6, the geometric flats of M form a sublattice of \mathcal{L}_M , isomorphic to \mathcal{L}_P by Definition-Proposition 2.11. Hence,

$$\text{cl}_M(\pi^{-1}(F_1) \cup \dots \cup \pi^{-1}(F_k)) \in \tilde{\mathcal{G}} \iff \text{cl}_P(F_1 \cup \dots \cup F_k) \in \mathcal{G}. \quad \square$$

Remark 3.5 (Geometry of building sets). Let $\{V_i\}_i$ be a subspace arrangement in V , defining a polymatroid P . Let \mathcal{G} be a building set for P . By [5, §1.6], the *wonderful compactification* of $\{V_i\}_i$ with respect to \mathcal{G} can be constructed by blowing up $\mathbb{P}(V)$ along all subspaces $\cap_{i \in F} \mathbb{P}(V_i)$ with $F \in \mathcal{G}$, first blowing up those of dimension 0, then those of dimension 1, and so on. Let $\{V_{ij}\}_{ij}$ be a hyperplane arrangement realizing \tilde{P} , as in Example 2.12. Blowing up a codimension 1 subvariety is an isomorphism, so Lemma 3.4 implies that the wonderful compactification of $\{V_i\}_i$ with respect to a building set \mathcal{G} is isomorphic to the wonderful compactification of $\{V_{ij}\}_{ij}$ with respect to $\tilde{\mathcal{G}}$.

3.2. Bergman fans. Let P be a polymatroid on E , with minimal lift \tilde{P} on \tilde{E} . Let $\mathbb{R}^{\tilde{E}}$ be the vector space spanned by \mathbf{e}_i for $i \in \tilde{E}$, and write $\mathbf{e}_S := \sum_{i \in S} \mathbf{e}_i$ for $S \subseteq \tilde{E}$. If $\mathcal{S} \subseteq 2^{\tilde{E}}$ is a collection of subsets, write

$$\sigma_{\mathcal{S}} := \text{cone}(\mathbf{e}_S : S \in \mathcal{S}) \subseteq \mathbb{R}^{\tilde{E}} / \mathbb{R}(1, 1, \dots, 1).$$

Definition 3.6. Let P be a polymatroid on E , and $\mathcal{G} \subseteq \mathcal{L}_P$ a geometric building set of P . The *Bergman fan* associated to (P, \mathcal{G}) is $\Sigma_{P, \mathcal{G}} := \{\sigma_{\mathcal{N}}\}_{\mathcal{N}}$, where \mathcal{N} ranges over all $\tilde{\mathcal{G}}$ -nested sets of \tilde{P} such that $\tilde{E} \notin \mathcal{N}$.

If P is a matroid and \mathcal{G} is the maximal geometric building set of P , then $\tilde{P} = P$ and $\tilde{\mathcal{G}} = \mathcal{G}$. In this case, $\Sigma_{P, \mathcal{G}}$ coincides with the Bergman fan of [1, Definition 3.2].

Lemma 3.7. *Let P be a polymatroid and let \mathcal{G} be the maximal geometric building set of P . Then the $\tilde{\mathcal{G}}$ -nested sets of \tilde{P} are in bijection with the chains of flats $\mathcal{F} = \{\emptyset = F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_k \subsetneq E\}$ of P and a subset S of \tilde{E} such that $\text{rk}_P(F \cup \pi(T)) > \text{rk}_P(F) + |T|$ for all proper $F \in \mathcal{F}$ and nonempty $T \subseteq S \setminus \pi^{-1}(F)$.*

Proof. Let \mathcal{N} be a nested set of $\tilde{\mathcal{G}}$, which consists of some geometric flats and some atoms. As the join of two geometric flats is geometric, we see that the geometric flats must form a chain $\{\emptyset = F_0 \subsetneq \pi^{-1}(F_1) \subsetneq \pi^{-1}(F_2) \subsetneq \dots \subsetneq \pi^{-1}(F_k) \subsetneq \tilde{E}\}$, where F_1, \dots, F_k are flats of P . Let S be the set of atoms in \mathcal{N} . It suffices to check that $S \cup \{\pi^{-1}(F_1), \dots, \pi^{-1}(F_k)\}$ is nested if and only if $\text{rk}_P(F \cup \pi(T)) > \text{rk}_P(F) + |T|$ for all proper $F \in \mathcal{F}$ and nonempty $T \subseteq S \setminus \pi^{-1}(F)$.

If the inequality holds, then

$$\text{rk}_P(F \cup \pi(T)) > \text{rk}_P(F) + |T| \geq \text{rk}_{\tilde{P}}(\pi^{-1}(F) \cup T)$$

for any $F \in \{F_1, \dots, F_k\}$ and all nonempty $T \subseteq S \setminus \pi^{-1}(F)$. The left-hand side is the rank of the smallest geometric flat containing $\pi^{-1}(F) \cup T$, so the closure of $\pi^{-1}(F) \cup T$ cannot be geometric. Therefore $S \cup \{\pi^{-1}(F_1), \dots, \pi^{-1}(F_k)\}$ is nested.

Now suppose that $S \cup \{\pi^{-1}(F_1), \dots, \pi^{-1}(F_k)\}$ is nested, but the inequality fails for some $F \in \{F_1, \dots, F_k\}$ and some $T \subseteq S \setminus \pi^{-1}(F)$. Let G be the smallest geometric flat containing $\pi^{-1}(F) \cup T$. Then

$$\text{rk}_{\tilde{P}}(\pi^{-1}(F) \cup T) \leq \text{rk}_{\tilde{P}}(G) = \text{rk}_P(F \cup \pi(T)) \leq \text{rk}_P(F) + |T|.$$

If the first inequality is an equality, then $\text{rk}_{\tilde{P}}(\pi^{-1}(F) \cup T) = \text{rk}_{\tilde{P}}(G)$, and so the closure of $\pi^{-1}(F) \cup T$ is a geometric flat, contradicting that \mathcal{N} is nested. Therefore $\text{rk}_{\tilde{P}}(\pi^{-1}(F) \cup T) < \text{rk}_P(F) + |T| = \text{rk}_{\tilde{P}}(\pi^{-1}(F)) + |T|$, so there is a circuit $C \subseteq \pi^{-1}(F) \cup T$ with $C \setminus \pi^{-1}(F)$ nonempty. For any $c \in C \setminus \pi^{-1}(F)$, we have $c \in \text{cl}_{\tilde{P}}(\pi^{-1}(F) \cup C \setminus c)$, and so by Lemma 2.8(i),

$$\pi^{-1}(\pi(c)) \subseteq \text{cl}_{\tilde{P}}(\pi^{-1}(F) \cup C \setminus c).$$

Using this for all $c \in C \setminus \pi^{-1}(F)$, we have $\pi^{-1}(\pi(C)) \cup \pi^{-1}(F) \subseteq \text{cl}_{\tilde{P}}(\pi^{-1}(F) \cup C)$, so $\text{cl}_{\tilde{P}}(\pi^{-1}(\pi(C)) \cup \pi^{-1}(F)) = \text{cl}_{\tilde{P}}(\pi^{-1}(F) \cup C)$. Note that $\text{cl}_{\tilde{P}}(\pi^{-1}(\pi(C)) \cup \pi^{-1}(F))$ is geometric. Because $C \setminus \pi^{-1}(F) \subseteq T$, this contradicts that \mathcal{N} is nested. \square

Corollary 3.8. *If P is a polymatroid and \mathcal{G} is the maximal geometric building set, then $\Sigma_{P,\mathcal{G}}$ coincides with the Bergman fan Σ_P defined in Definition 1.7.*

Lemma 3.9. *If P is a polymatroid and \mathcal{G} is a geometric building set of P , then $\Sigma_{P,\mathcal{G}}$ is a subfan of the normal fan of a convex polytope.*

Proof. Let E be the ground set of P and B be the Boolean polymatroid with $\text{rk}_B(i) = \text{rk}_P(i)$ for all $i \in E$. Suppose \mathcal{G} is a building set for P . The definition of geometric building set implies for any flat F of P , $\max \mathcal{G}_{\leq F}$ is a partition of F . Hence, $\tilde{\mathcal{G}}$ is a building set of both \tilde{B} and \tilde{P} , and there is an inclusion of fans $\Sigma_{P,\mathcal{G}} = \Sigma_{\tilde{P},\tilde{\mathcal{G}}} \subseteq \Sigma_{\tilde{B},\tilde{\mathcal{G}}}$ (In fact, if all single-element subsets of E are flats of P , then \mathcal{G} is a geometric building set for B , and we obtain $\Sigma_{P,\mathcal{G}} \subseteq \Sigma_{\tilde{B},\tilde{\mathcal{G}}} = \Sigma_{B,\mathcal{G}}$). The lattice of flats of \tilde{B} is isomorphic to the lattice of subsets of \tilde{E} , so $\Sigma_{\tilde{B},\tilde{\mathcal{G}}}$ is the normal fan of a convex polytope by [16, Theorem 7.4]. \square

By [4], for any polymatroid P and a geometric building set \mathcal{G} , $A(\Sigma_{P,\mathcal{G}})$ has the following presentation.

Proposition 3.10. *The Chow ring of $\Sigma_{P,\mathcal{G}}$ satisfies*

$$A(\Sigma_{P,\mathcal{G}}) = \mathbb{Z}[z_G : G \in \tilde{\mathcal{G}} \setminus \{\tilde{E}\}] / I_{P,\mathcal{G}}$$

where $I_{P,\mathcal{G}}$ is the ideal generated by

$$\begin{aligned} & z_{G_1} \cdots z_{G_k} \quad \text{for any not } \tilde{\mathcal{G}}\text{-nested collection } \{G_1, \dots, G_k\}, \\ & \sum_{i \in G} z_G - \sum_{j \in F} z_F \quad \text{for any } i, j \in \tilde{E}. \end{aligned}$$

In particular, the Chow ring of Σ_P satisfies

$$A(\Sigma_P) = \mathbb{Z}[z_F : F \text{ nonempty proper flat of } P] \otimes \mathbb{Z}[z_i : i \in \tilde{E}] / I_P,$$

where I_P is an ideal generated by the following polynomials where z_\emptyset is replaced by 1 wherever it appears:

$$\begin{aligned} z_F - z_i, \quad \pi^{-1}(F) = \{i\} \text{ is a singleton set,} \\ z_{F_1} z_{F_2}, \quad F_1 \text{ and } F_2 \text{ are incomparable proper flats of } P, \\ z_F \prod_{i \in T} z_i, \quad F \text{ is a proper flat and } T \subseteq \tilde{E} \setminus \pi^{-1}(F) \text{ is nonempty satisfying} \\ \text{rk}_P(F \cup \pi(T)) \leq \text{rk}_P(F) + |T|, \\ \sum_{i \in F} z_F - \sum_{j \in G} z_G, \quad i \text{ and } j \text{ are elements of } \tilde{E}. \end{aligned}$$

We identify $A^1(\Sigma_{P,\mathcal{G}})$ with the space of piecewise linear functions on the support of $\Sigma_{P,\mathcal{G}}$, modulo global linear functions. Explicitly, a piecewise linear function ℓ is a representative of

$$\sum_{\mathbf{u}_F} \ell(\mathbf{u}_F) z_F \in A^1(\Sigma_{P,\mathcal{G}}),$$

where the sum is over all primitive ray generators \mathbf{u}_F of rays of $\Sigma_{P,\mathcal{G}}$.

Remark 3.11. A slightly different presentation of $A(\Sigma_{P,\mathcal{G}})$ is used by [8]. For $F \in \tilde{\mathcal{G}}$,

$$y_F := \begin{cases} -\sum_{i \in G} z_G, & F = \tilde{E} \\ z_F, & \text{otherwise.} \end{cases}$$

In terms of the y_F 's, $A(\Sigma_{P,\mathcal{G}})$ is defined by the ideal I_{FY} generated by

$$\begin{aligned} y_{G_1} \cdots y_{G_k}, \quad \{G_1, \dots, G_k\} \text{ not } \tilde{\mathcal{G}}\text{-nested and} \\ \sum_{i \in G} y_G, \quad i \in \tilde{E}. \end{aligned}$$

Remark 3.12 (Tropicalization and the Bergman fan). Suppose P is the matroid realized by an essential hyperplane arrangement V_1, \dots, V_n in V , that is, the intersection $V_1 \cap \cdots \cap V_n$ is equal to the origin. Denote the defining equation of V_i by $\ell_i = 0$. The inclusion

$$\mathbb{P}(V \setminus \cup_i V_i) \hookrightarrow \mathbb{P}((\mathbb{F}^*)^n), \quad v \mapsto [\ell_1(v) : \dots : \ell_n(v)]$$

shows that $\mathbb{P}(V \setminus \cup_i V_i)$ is a very affine variety. Its tropicalization is the support of Σ_P , by [17, §9.3] and [3, §3] (see also [7, Theorem 4.1]). The corresponding statement for realizable polymatroids does not make sense: the complement of a subspace arrangement may not be very affine, so tropicalization [12, Definition 3.2.1] is not defined. Nevertheless, if $\{V_i\}_i$ is a subspace arrangement (so P is a polymatroid), then generic hyperplanes $\{V_{ij}\}_{ij}$ realizing \tilde{P} as in Example 2.12 define a subtorus of $\mathbb{P}(\prod_i V/V_i)$. Tropicalizing this torus's intersection with $\mathbb{P}(V \setminus \cup_i V_i)$ gives the support of $\Sigma_{\tilde{P}}$, which coincides with that of Σ_P .

4. CHOW RINGS OF POLYMATROIDS

If \mathcal{S} is a collection of sets, write $\cup \mathcal{S}$ for the union of the elements of \mathcal{S} . In [15, Section 4], Pagaria and Pezzoli define the Chow ring of a polymatroid as follows³.

³Unlike [15], we use \mathbb{Z} -coefficients.

Definition 4.1. Let P be a polymatroid and $\mathcal{G} \subseteq \mathcal{L}_P$ a geometric building set. The *Chow ring* of (P, \mathcal{G}) is

$$\mathrm{DP}(P, \mathcal{G}) := \mathbb{Z}[x_F : F \in \mathcal{G}] / I_{\mathrm{DP}},$$

where I_{DP} is the ideal generated by

$$x_{G_1} \cdots x_{G_k} \left(\sum_{\mathcal{G} \ni H \geq G} x_H \right)^b$$

for $G \in \mathcal{G}$, $\mathcal{S} = \{G_1, \dots, G_k\} \subseteq \mathcal{G}$, and $b \geq \mathrm{rk}_P(G) - \mathrm{rk}_P(\cup \mathcal{S}_{<G})$.

When P is realizable by an arrangement of subspaces, $\mathrm{DP}(P, \mathcal{G})$ is the cohomology ring of De Concini & Procesi's wonderful compactification of the arrangement complement. In this section, we make use of the theory of Gröbner bases. For background on this subject, see [6, Chapter 15].

Theorem 4.2. *Let P be a polymatroid on E with lift \tilde{P} on \tilde{E} , and $\pi: \tilde{E} \rightarrow E$ the projection. There is an isomorphism $\mathrm{DP}(P, \mathcal{G}) \cong A(\Sigma_{P, \mathcal{G}})$ sending x_F to $y_{\pi^{-1}(F)}$.*

Proof. Let $I_{\mathrm{DP}} \subseteq \mathbb{Z}[x_F : F \in \mathcal{G}]$ be the defining ideal of $\mathrm{DP}(P, \mathcal{G})$, and let $I_{FY} \subseteq \mathbb{Z}[y_F : F \in \tilde{\mathcal{G}}]$ be the defining ideal of $A(\Sigma_{P, \mathcal{G}})$ as in [8] (See Remark 3.11). We define the following map on polynomial rings.

$$\varphi: \mathbb{Z}[x_F : F \in \mathcal{G}] \rightarrow \mathbb{Z}[y_F : F \in \tilde{\mathcal{G}}], \quad x_F \mapsto y_{\pi^{-1}(F)}$$

First we show $\varphi(I_{\mathrm{DP}}) \subseteq I_{FY}$. Write f for one of the defining relations of I_{DP} :

$$f = \left(\prod_{F \in \mathcal{S}} x_F \right) \left(\sum_{\mathcal{G} \ni H \geq G} x_H \right)^b.$$

By [8, Theorems 1 and 3], I_{FY} contains the following two types of polynomials:

$$\begin{aligned} & \prod_{F \in \mathcal{S}} y_F, \quad \mathcal{S} \text{ not } \tilde{\mathcal{G}}\text{-nested,} \\ & \prod_{F \in \mathcal{N}} y_F \left(\sum_{H \geq G} y_H \right)^d, \quad \mathcal{N} \text{ a nested antichain, } \cup \mathcal{N} < G, \text{ and } d = \mathrm{rk}(G) - \mathrm{rk}(\cup \mathcal{N}). \end{aligned}$$

If \mathcal{S} is not \mathcal{G} -nested, then $\tilde{\mathcal{S}} := \{\pi^{-1}(F) : F \in \mathcal{S}\}$ is not $\tilde{\mathcal{G}}$ -nested by Lemma 3.4. Hence, $\varphi(f)$ is divisible by a relation of the first type. Otherwise, \mathcal{S} is \mathcal{G} -nested, so $\tilde{\mathcal{S}}$ is $\tilde{\mathcal{G}}$ -nested. In this case, $\varphi(f)$ is divisible by a relation of the second type because

$$\begin{aligned} b & \geq \mathrm{rk}_P(G) - \mathrm{rk}_P(\cup \mathcal{S}_{<G}) = \mathrm{rk}_{\tilde{P}}(\pi^{-1}(G)) - \mathrm{rk}_{\tilde{P}}(\cup \tilde{\mathcal{S}}_{<\pi^{-1}(G)}) \\ & = \mathrm{rk}_{\tilde{P}}(\pi^{-1}(G)) - \mathrm{rk}_{\tilde{P}}(\cup \max \tilde{\mathcal{S}}_{<\pi^{-1}(G)}). \end{aligned}$$

This proves that $\varphi(I_{\mathrm{DP}}) \subseteq I_{FY}$, so φ descends to $\tilde{\varphi}: \mathrm{DP}(P, \mathcal{G}) \rightarrow A(\Sigma_{P, \mathcal{G}})$.

If $F \in \tilde{\mathcal{G}}$ is a flat of rank greater than 1, then y_F is in the image of $\tilde{\varphi}$. By the linear relation $\sum_{i \in G} y_G = 0$, it follows that y_i is also in the image of $\tilde{\varphi}$. Therefore, $\tilde{\varphi}$ is surjective. It remains to show that $\tilde{\varphi}$ is injective. By [8, Theorem 2], the generators of I_{FY} in the previous paragraph are a Gröbner basis with respect to any lexicographic monomial order $<$ in which $F_1 \subseteq F_2$ implies $y_{F_1} > y_{F_2}$. Any such order is an elimination order with respect to $\{y_i : i \in \tilde{E}\}$. By [6, Proposition 15.29]⁴, the generators of I_{FY} in the previous paragraph that do not involve any

⁴Eisenbud's proof of this statement works over \mathbb{Z} because all leading coefficients in our Gröbner basis are 1.

$y_i, i \in \tilde{E}$, are a Gröbner basis for $\text{im}(\varphi) \cap I_{FY}$. Any such polynomial is the image of a generator of I_{DP} , so $\varphi^{-1}(I_{FY}) = I_{DP}$. This implies φ is an isomorphism. \square

The proof of Theorem 4.2 also shows the following, originally obtained from [15, Corollary 2.8].

Corollary 4.3. *The defining relations of $\text{DP}(P, \mathcal{G})$ in Definition 4.1 form a Gröbner basis with respect to any lexicographic $<$ such that $x_{F_1} < x_{F_2}$ whenever $F_1 \supsetneq F_2$.*

This recovers the monomial basis of [15].

Corollary 4.4. *The following monomials are a \mathbb{Z} -basis of $\text{DP}(P, \mathcal{G})$:*

$$x_{G_1}^{a_1} \cdots x_{G_k}^{a_k}$$

where $N = \{G_1, \dots, G_k\}$ is a nested set of \mathcal{G} , and

$$1 \leq a_i < \text{rk}(F_i) - \text{rk}(\cup_{N < F_i}) \quad \text{for all } 1 \leq i \leq k.$$

Proof. Immediate from Theorem 4.2 and [8, Corollary 1] (or Corollary 4.3 and [6, Theorem 15.3]). \square

Remark 4.5. In [15, Corollary 2.8], degree-lexicographic order (also called ‘‘homogeneous lexicographic’’ or ‘‘graded lexicographic’’ order) is used. Since the defining relations of $\text{DP}(P, \mathcal{G})$ in Definition 4.1 are all homogeneous, their initial terms with respect to the lex or degree-lex orders are the the same. Thus, they are a Gröbner basis with respect to one order if and only if they are with respect to the other.

Remark 4.6. Suppose P is a polymatroid with building set \mathcal{G} , realized by an arrangement $\{V_i\}_i$ in V . Let $\{V_{ij}\}_{ij}$ realize \tilde{P} as in Example 2.12. In this case, an alternate proof of Theorem 4.2 is possible. By [8, §4], $A(\Sigma_{P, \mathcal{G}})$ is isomorphic to the Chow ring of the wonderful compactification of $\{V_{ij}\}_{ij}$ with respect to $\tilde{\mathcal{G}}$. By Remark 3.5, the wonderful compactifications of $\{V_{ij}\}_{ij}$ with respect to $\tilde{\mathcal{G}}$ and $\{V_i\}_i$ with respect to \mathcal{G} are isomorphic. Hence, their Chow rings are isomorphic. The Chow ring of the latter space is isomorphic to $\text{DP}(P, \mathcal{G})$ by a comparison of the presentations in Definition 4.1 and [5, Theorem 5.2].

In the remainder of this section, we recover [15, Theorem 4.7] and generalize [15, Theorem 4.21] using Theorem 4.2 and the tropical Hodge theory of [2]. If R is a \mathbb{Z} -algebra, define $R_{\mathbb{Q}} := R \otimes_{\mathbb{Z}} \mathbb{Q}$ and $R_{\mathbb{R}}$ likewise.

Corollary 4.7. *Let $\mathbb{K} = \mathbb{Q}, \mathbb{R}$. Let P be a polymatroid of rank r and \mathcal{G} a geometric building set. Let ℓ be any \mathbb{K} -valued strictly convex piecewise linear function on $\Sigma_{P, \mathcal{G}}$, viewed as an element of $A^1(\Sigma_{P, \mathcal{G}})_{\mathbb{K}} \cong \text{DP}^1(P, \mathcal{G})_{\mathbb{K}}$.*

(i) *(Poincaré duality) There is an isomorphism*

$$\text{deg}: \text{DP}(P, \mathcal{G})^{r-1} \rightarrow \mathbb{Z},$$

and for all $0 \leq k < r/2$, the pairing

$$\text{DP}^k(P, \mathcal{G}) \times \text{DP}^{r-k-1}(P, \mathcal{G}) \rightarrow \mathbb{Z}, \quad (a, b) \mapsto \text{deg}(ab)$$

is non-degenerate.

(ii) *(Hard Lefschetz) For every $0 \leq k < r/2$, the multiplication map*

$$\text{DP}^k(P, \mathcal{G})_{\mathbb{K}} \rightarrow \text{DP}^{r-k-1}(P, \mathcal{G})_{\mathbb{K}}, \quad a \mapsto \ell^{r-2k-1}a$$

is an isomorphism.

(iii) (Hodge-Riemann) For every $0 \leq k < r/2$, the bilinear form

$$\mathrm{DP}^k(P, \mathcal{G})_{\mathbb{K}} \times \mathrm{DP}^k(P, \mathcal{G})_{\mathbb{K}} \rightarrow \mathbb{K}, \quad (a, b) \mapsto (-1)^k \deg(\ell^{r-2k-1}ab)$$

is positive definite on the kernel of multiplication by ℓ^{r-2k} .

Proof. Let $M = \tilde{P}$. By [8, Theorem 4], Σ_M is a refinement of $\Sigma_{P, \mathcal{G}}$. In particular, the two fans have the same support. By [1, Proposition 2.4] and Lemma 3.9, both fans are subfans of the normal fans of convex polytopes, so both fans support strictly convex piecewise linear functions. By [8, Proposition 2], $\Sigma_{P, \mathcal{G}}$ is a smooth fan.

The desired statements now follow by applying [2, Theorem 1.6], [1, Proposition 5.2], and [1, Theorems 6.19 and 8.8]. Loosely, [2, Theorem 1.6] says properties (i), (ii), and (iii) hold for $A(\Sigma_{P, \mathcal{G}})_{\mathbb{R}} \cong \mathrm{DP}(P, \mathcal{G})_{\mathbb{R}}$ if and only if they hold for $A(\Sigma_M)_{\mathbb{R}}$, and [1] verifies them for $A(\Sigma_M)_{\mathbb{R}}$.

Properties (ii) and (iii) for $\mathrm{DP}(P, \mathcal{G})_{\mathbb{Q}}$ follow immediately from those for $\mathrm{DP}(P, \mathcal{G})_{\mathbb{R}}$. For (i), note that [2]'s Poincaré duality arguments go through over \mathbb{Z} . (Explicitly, one must check statements 6.6–6.9, and Propositions 6.16, 6.17 of [2].) \square

Remark 4.8. In [15], Corollary 4.7 is proved for ℓ in the σ -cone [15, Definition 4.15], the positive span of

$$-\sum_{G \in \mathcal{G}_{\geq F}} x_G, \quad F \in \mathcal{G}$$

in $\mathrm{DP}(P, \mathcal{G})$. The σ -cone is generally a proper subset of the cone of strictly convex piecewise linear functions on $\Sigma_{P, \mathcal{G}}$. For example, if M is a loopless matroid on E and \mathcal{G} is its maximal building set, then for any $i \in E$,

$$\beta := \sum_{i \notin F} z_F = -\sum_{|G| > 1} (|G| - 1)y_G$$

is in the closure of the cone of strictly convex piecewise linear functions on Σ_M [1, Proposition 4.3, Lemma 9.7]. However, β may not be in the closure of the σ -cone, e.g. when M is Boolean of rank at least 3. For comparison of the σ -cone to the ample cone of the wonderful compactification, see [15, Remark 4.22].

APPENDIX A. COMBINATORICS OF THE BERGMAN FAN OF BOOLEAN POLYMATROIDS

In this appendix we describe the combinatorics of the Bergman fans of Boolean polymatroids, proving in particular that they are the normal fans of polypermutohedra as stated in the introduction. We also give a description of polypermutohedra as a Minkowski sum of simplices. Throughout this appendix, we let $\pi: \tilde{E} \rightarrow E$ be a surjective map of finite sets, with associated Boolean polymatroid $B(\pi)$ on E given by the rank function $\mathrm{rk}_{B(\pi)}(A) = |\pi^{-1}(A)|$ for $A \subseteq E$. We write n for the cardinality of E .

A.1. The Bergman fan as a configuration space.

Definition A.1. Let $\mathbf{w} = (w_i)_i \in \mathbb{R}^{\tilde{E}}$ be a weight on the elements of \tilde{E} . Write $\mathrm{Lowest}_{\pi}(\mathbf{w})$ for the set of $i \in \tilde{E}$ such that i has minimal weight among the elements of $\pi^{-1}(\pi(i))$ with respect to \mathbf{w} . We equip $\mathrm{Lowest}_{\pi}(\mathbf{w})$ with the natural partial preorder given by $i \preceq j$ if $w_i \leq w_j$.

Adding a multiple of the all ones vector to \mathbf{w} does not change $\mathrm{Lowest}_{\pi}(\mathbf{w})$, so $\mathrm{Lowest}_{\pi}(\mathbf{w})$ is well defined for $\mathbf{w} \in \mathbb{R}^{\tilde{E}}/\mathbb{R}(1, 1, \dots, 1)$.

Lemma A.2. *Two points $\mathbf{w}, \mathbf{v} \in \mathbb{R}^{\tilde{E}}/(1, 1, \dots, 1)$ lie in the relative interior of the same cone of the $\Sigma_{B(\pi)}$ if and only if $\text{Lowest}_\pi(\mathbf{v}) = \text{Lowest}_\pi(\mathbf{w})$ as posets.*

Proof. Recall that a cone $\sigma_{\mathcal{F}, S}$ of the fan $\Sigma_{B(\pi)}$ is determined by a chain of sets $\mathcal{F} = \{\emptyset \subsetneq F_1 \subsetneq \dots \subsetneq F_k \subsetneq E\}$ and a set $S \subseteq \tilde{E}$ such that S does not contain a fiber of π . The relative interior of $\sigma_{\mathcal{F}, S}$ contains \mathbf{w} if and only if the underlying set of $\text{Lowest}_\pi(\mathbf{w})$ is equal to $\tilde{E} \setminus S$ and $i \prec j$ whenever there exists r such that $i \notin F_r$ and $j \in F_r$. Therefore $\text{Lowest}_P(\mathbf{w})$ can be recovered from $\sigma_{\mathcal{F}, A}$ and vice versa. \square

A.2. The Bergman fan as the normal fan of a polytope. Recall that an *ordered transversal* of π is a sequence s_1, \dots, s_n of elements of \tilde{E} such that each fiber of π contains exactly one element of the sequence.

Definition A.3. Given real numbers $0 \leq c_1 < c_2 < \dots < c_n$, define the associated polypermutohedron $Q(\pi; c_1, c_2, \dots, c_n)$ as the convex hull of the vectors $\mathbf{v}_{s_1, s_2, \dots, s_n} := c_1 \mathbf{e}_{s_1} + c_2 \mathbf{e}_{s_2} + \dots + c_n \mathbf{e}_{s_n}$, where s_1, s_2, \dots, s_n runs over all ordered transversals of π and $\mathbf{e}_i \in \mathbb{R}^{\tilde{E}}$ is the standard basis vector of $i \in \tilde{E}$.

Lemma A.4. *Let $\mathbf{w} = (w_i)_i \in \mathbb{R}^{\tilde{E}}$ be a weight on the elements of \tilde{E} , and let s_1, s_2, \dots, s_n be an ordered transversal of π . Denote by $\langle -, - \rangle$ the standard dot product on $\mathbb{R}^{\tilde{E}}$. Then the linear functional $\langle \mathbf{w}, - \rangle$ achieves its minimum over $Q(\pi; c_1, c_2, \dots, c_n)$ at the vector $\mathbf{v}_{s_1, s_2, \dots, s_n}$ if and only if*

- (i) s_j has minimum weight among the elements of $\pi(\pi^{-1}(s_j))$ with respect to \mathbf{w} for all j , and
- (ii) $w_{s_1} \leq w_{s_2} \leq \dots \leq w_{s_n}$.

Proof. Suppose that $\mathbf{v}_{s_1, s_2, \dots, s_n}$ minimizes $\langle \mathbf{w}, - \rangle$ over $Q(\pi; c_1, c_2, \dots, c_n)$. Assume contrary to (i) that there is some j and $r \in \pi(\pi^{-1}(s_j))$ such that $w_{s_j} > w_r$. Then replacing s_j with r gives another ordered transversal of π whose corresponding vector has smaller dot product with \mathbf{w} :

$$\begin{aligned} \langle \mathbf{w}, \mathbf{v}_{s_1, \dots, s_j, \dots, s_n} \rangle &= c_1 w_{s_1} + \dots + c_j w_{s_j} + \dots + c_n w_{s_n} \\ &> c_1 w_{s_1} + \dots + c_j w_r + \dots + c_n w_{s_n} = \langle \mathbf{w}, \mathbf{v}_{s_1, \dots, r, \dots, s_n} \rangle. \end{aligned}$$

This proves (i). Now assume contrary to (ii) that there is some j such that $w_{s_j} > w_{s_{j+1}}$. Then switching the order of s_j and s_{j+1} gives another ordered transversal, and thus another vector $\mathbf{v}' \in Q(\pi; c_1, c_2, \dots, c_n)$, where by assumption $\langle \mathbf{w}, \mathbf{v}_{s_1, s_2, \dots, s_n} \rangle \leq \langle \mathbf{w}, \mathbf{v}' \rangle$. Thus $c_j w_{s_j} + c_{j+1} w_{s_{j+1}} \leq c_j w_{s_{j+1}} + c_{j+1} w_{s_j}$, which contradicts the fact that $ab + cd > ac + bd$ whenever $a > c$ and $b > d$. This proves (ii).

For the other direction, assume that \mathbf{w} satisfies the two conditions. If $\langle \mathbf{w}, - \rangle$ achieves its minimum over $Q(\pi; c_1, c_2, \dots, c_n)$ on a vector $\mathbf{v}_{s'_1, s'_2, \dots, s'_n}$, then by the first direction we must have that $w_{s_j} = w_{s'_j}$. Therefore $\langle \mathbf{w}, - \rangle$ also achieves its minimum on $\mathbf{v}_{s_1, s_2, \dots, s_n}$. \square

Proposition A.5. *The inner normal fan of $Q(\pi; c_1, c_2, \dots, c_n)$, modulo the all ones vector, is $\Sigma_{B(\pi)}$.*

Proof. By Lemma A.4, the vertices of $Q(\pi; c_1, c_2, \dots, c_n)$ correspond to ordered transversals of π , and the set of vertices on which a given linear functional $\langle \mathbf{w}, - \rangle$ achieves its minimum is equivalent to the data of $\text{Lowest}_\pi(\mathbf{w})$. Therefore the proposition follows by Lemma A.2. \square

A.3. Minkowski sums of simplices. For $S \subseteq \tilde{E}$, let Δ_S be the convex hull of the vectors \mathbf{e}_i , for $i \in S$.

Proposition A.6. $Q(\pi; 1, 2, \dots, n)$ is the Minkowski sum $\sum_{\{i,j\} \subseteq E} \Delta_{\pi^{-1}(\{i,j\})}$.

In the sum, we allow $i = j$. When π is a bijection, this recovers the description of the usual permutohedron as the graphical zonotope of the complete graph.

Proof. The proof of Proposition A.5 shows that the inner normal fan of $\Delta_{\pi^{-1}(S)}$ is a coarsening of $\Sigma_{B(\pi)}$ for any $S \subseteq E$. In particular, the inner normal fan of the Minkowski sum $\sum_{\{i,j\} \subseteq E} \Delta_{\pi^{-1}(\{i,j\})}$ is a coarsening of $\Sigma_{B(\pi)}$. We may then find all vertices of the Minkowski sum by choosing a maximal cone of $\Sigma_{B(\pi)}$ and finding the vertex of the Minkowski sum on which any vector in the interior of this cone achieves its minimum.

Each maximal cone of the fan $\Sigma_{B(\pi)}$ corresponds to a maximal chain of subsets $\mathcal{F} = \{\emptyset \subsetneq F_1 \subsetneq \dots \subsetneq F_{n-1} \subsetneq E\}$ and a subset $S \subseteq \tilde{E}$ such that $|\pi^{-1}(i) \setminus S| = 1$ for all i . This data is equivalent to the data of an ordered transversal s_1, \dots, s_n of π . Choose a maximal cone of $\Sigma_{B(\pi)}$ corresponding to an ordered transversal s_1, \dots, s_n , and choose a vector in the relative interior of this cone. We can compute the vertex of the Minkowski sum on which this vector achieves its minimum by adding up the minimal vertices of each summand. The minimal vertex of a summand of the form $\Delta_{\pi^{-1}(i)}$ is \mathbf{e}_{s_k} , where k is the unique element of $\tilde{E} \setminus S$ such that $\pi(s_k) = i$. The minimal vertex of a summand of the form $\Delta_{\pi^{-1}(\{i,j\})}$ for $i \neq j$ is \mathbf{e}_{s_ℓ} , where ℓ is the smaller index of the two elements of $\pi^{-1}(\{i,j\}) \cap \{s_1, \dots, s_n\}$. We see that the minimal vertex of the Minkowski sum is $\sum_{i=1}^n i \mathbf{e}_{s_i}$, as desired. \square

Remark A.7. One can deduce from the theory of building sets, e.g., [16, Proposition 7.5], that $\Sigma_{B(\pi)}$ is the normal fan of the Minkowski sum $\sum_{\emptyset \neq S \subseteq E} \Delta_{\pi^{-1}(S)}$.

REFERENCES

- [1] Karim Adiprasito, June Huh, and Eric Katz, *Hodge theory for combinatorial geometries*, Ann. of Math. (2) **188** (2018), no. 2, 381–452. MR 3862944 [2](#), [10](#), [15](#)
- [2] Federico Ardila, Graham Denham, and June Huh, *Lagrangian geometry of matroids*, J. Amer. Math. Soc. **36** (2023), no. 3, 727–794. MR 4583774 [2](#), [14](#), [15](#)
- [3] Federico Ardila and Caroline J. Klivans, *The Bergman complex of a matroid and phylogenetic trees*, J. Combin. Theory Ser. B **96** (2006), no. 1, 38–49. MR 2185977 [12](#)
- [4] Michel Brion, *Piecewise polynomial functions, convex polytopes and enumerative geometry*, Parameter spaces (Warsaw, 1994), Banach Center Publ., vol. 36, Polish Acad. Sci. Inst. Math., Warsaw, 1996, pp. 25–44. MR 1481477 [11](#)
- [5] C. De Concini and C. Procesi, *Wonderful models of subspace arrangements*, Selecta Math. (N.S.) **1** (1995), no. 3, 459–494. MR 1366622 [1](#), [9](#), [10](#), [14](#)
- [6] David Eisenbud, *Commutative algebra*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995, With a view toward algebraic geometry. MR 1322960 [13](#), [14](#)
- [7] Eva Maria Feichtner and Bernd Sturmfels, *Matroid polytopes, nested sets and Bergman fans*, Port. Math. (N.S.) **62** (2005), no. 4, 437–468. MR 2191630 [12](#)
- [8] Eva Maria Feichtner and Sergey Yuzvinsky, *Chow rings of toric varieties defined by atomic lattices*, Invent. Math. **155** (2004), no. 3, 515–536. MR 2038195 [12](#), [13](#), [14](#), [15](#)
- [9] Patricio Gallardo and Evangelos Routis, *Wonderful compactifications of the moduli space of points in affine and projective space*, Eur. J. Math. **3** (2017), no. 3, 520–564. MR 3687430 [3](#)
- [10] Thorkell Helgason, *Aspects of the theory of hypermatroids*, Hypergraph Seminar (Proc. First Working Sem., Ohio State Univ., Columbus, Ohio, 1972; dedicated to Arnold Ross), Lecture Notes in Math., vol. Vol. 411, Springer, Berlin-New York, 1974, pp. 191–213. MR 371691 [6](#)
- [11] L. Lovász, *Flats in matroids and geometric graphs*, Combinatorial surveys (Proc. Sixth British Combinatorial Conf., Royal Holloway Coll., Egham, 1977), 1977, pp. 45–86. MR 0480111 [6](#)

- [12] Diane Maclagan and Bernd Sturmfels, *Introduction to tropical geometry*, Graduate Studies in Mathematics, vol. 161, American Mathematical Society, Providence, RI, 2015. MR 3287221 [12](#)
- [13] Hien Q. Nguyen, *Submodular functions*, Theory of matroids (Neil White, ed.), Cambridge University Press, 1986, pp. 272–297. [6](#)
- [14] James Oxley, *Matroid theory*, second ed., Oxford Graduate Texts in Mathematics, vol. 21, Oxford University Press, Oxford, 2011. MR 2849819 [6](#)
- [15] Roberto Pagaria and Gian Marco Pezzoli, *Hodge Theory for Polymatroids*, Int. Math. Res. Not. IMRN (2023), no. 23, 20118–20168. MR 4675068 [2](#), [3](#), [4](#), [9](#), [12](#), [14](#), [15](#)
- [16] Alexander Postnikov, *Permutohedra, associahedra, and beyond*, Int. Math. Res. Not. IMRN (2009), no. 6, 1026–1106. MR 2487491 [11](#), [17](#)
- [17] Bernd Sturmfels, *Solving systems of polynomial equations*, CBMS Regional Conference Series in Mathematics, vol. 97, Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2002. MR 1925796 [12](#)

UNIVERSITY OF OREGON AND INSTITUTE FOR ADVANCED STUDY

Email address: crowley@uoregon.edu

PRINCETON UNIVERSITY AND KOREA INSTITUTE FOR ADVANCED STUDY

Email address: huh@princeton.edu

INSTITUTE FOR ADVANCED STUDY AND PRINCETON UNIVERSITY

Email address: mattlarson@princeton.edu

INSTITUTE FOR ADVANCED STUDY

Email address: connorgs@connorgs.net

UNIVERSITY OF WISCONSIN–MADISON

Email address: wang@math.wisc.edu